

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 November 2001 (01.11.2001)

PCT

(10) International Publication Number
WO 01/81379 A2

(51) International Patent Classification⁷: **C07K 14/195**

(74) Agents: **POTTER, Jane, E., R.**; Seed Intellectual Property Law Group PLLC, Suite 6300, 701 Fifth Avenue, Seattle, WA 98104-7092 et al. (US).

(21) International Application Number: PCT/US01/13081

(22) International Filing Date: 23 April 2001 (23.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/198,853 21 April 2000 (21.04.2000) US
60/219,752 20 July 2000 (20.07.2000) US

(71) Applicant (for all designated States except US): **CORIXA CORPORATION** [US/US]; 1124 Columbia Street, Suite 200, Seattle, WA 98104 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, FF, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).



(72) Inventors; and

(75) Inventors/Applicants (for US only): **BHATIA, Ajay** [IN/US]; 1805 Bellevue Avenue #204, Seattle, WA 98104 (US). **PROBST, Peter** [DE/US]; 137 N.W. 77th St., Seattle, WA 98117 (US). **STROMBERG, Erika, Jean** [US/US]; 440 N.E. 73rd Street #201, Seattle, WA 98103 (US).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/81379 A2

(54) Title: COMPOUNDS AND METHODS FOR TREATMENT AND DIAGNOSIS OF CHLAMYDIAL INFECTION

(57) Abstract: Compounds and methods for the diagnosis and treatment of Chlamydial infection are disclosed. The compounds provided include polypeptides that contain at least one antigenic portion of a *Chlamydia* antigen and DNA sequences encoding such polypeptides. Pharmaceutical compositions and vaccines comprising such polypeptides or DNA sequences are also provided, together with antibodies directed against such polypeptides. Diagnostic kits containing such polypeptides or DNA sequences and a suitable detection reagent may be used for the detection of Chlamydial infection in patients and in biological samples.

COMPOUNDS AND METHODS FOR TREATMENT AND DIAGNOSIS OF CHLAMYDIAL INFECTION

TECHNICAL FIELD

5 The present invention relates generally to the detection and treatment of Chlamydial infection. In particular, the invention is related to polypeptides comprising a *Chlamydia* antigen and the use of such polypeptides for the serodiagnosis and treatment of Chlamydial infection.

10 BACKGROUND OF THE INVENTION

 Chlamydiae are intracellular bacterial pathogens that are responsible for a wide variety of important human and animal infections. *Chlamydia trachomatis* is one of the most common causes of sexually transmitted diseases and can lead to pelvic inflammatory disease (PID), resulting in tubal obstruction and infertility. *Chlamydia*
15 *trachomatis* may also play a role in male infertility. In 1990, the cost of treating PID in the US was estimated to be \$4 billion. Trachoma, due to ocular infection with *Chlamydia trachomatis*, is the leading cause of preventable blindness worldwide. *Chlamydia pneumonia* is a major cause of acute respiratory tract infections in humans and is also believed to play a role in the pathogenesis of atherosclerosis and, in
20 particular, coronary heart disease. Individuals with a high titer of antibodies to *Chlamydia pneumonia* have been shown to be at least twice as likely to suffer from coronary heart disease as seronegative individuals. Chlamydial infections thus constitute a significant health problem both in the US and worldwide.

 Chlamydial infection is often asymptomatic. For example, by the time a woman
25 seeks medical attention for PID, irreversible damage may have already occurred resulting in infertility. There thus remains a need in the art for improved vaccines and pharmaceutical compositions for the prevention and treatment of *Chlamydia* infections. The present invention fulfills this need and further provides other related advantages.

30 SUMMARY OF THE INVENTION

 The present invention provides compositions and methods for the diagnosis and therapy of *Chlamydia* infection. In one aspect, the present invention provides polypeptides comprising an immunogenic portion of a *Chlamydia* antigen, or a

variant of such an antigen. Certain portions and other variants are immunogenic, such that the ability of the variant to react with antigen-specific antisera is not substantially diminished. Within certain embodiments, the polypeptide comprises an amino acid sequence encoded by a polynucleotide sequence selected from the group consisting of

5 (a) a sequence of SEQ ID NO: 1-48, 114-121, and 125-138; (b) the complements of said sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions. In specific embodiments, the polypeptides of the present invention comprise at least a portion of a *Chlamydial* protein that includes an amino acid sequence selected from the group consisting of sequences recited in SEQ ID NO:122-10 124 and 139-140 and variants thereof.

The present invention further provides polynucleotides that encode a polypeptide as described above, or a portion thereof (such as a portion encoding at least 15 amino acid residues of a *Chlamydial* protein), expression vectors comprising such polynucleotides and host cells transformed or transfected with such expression vectors.

15 In a related aspect, polynucleotide sequences encoding the above polypeptides, recombinant expression vectors comprising one or more of these polynucleotide sequences and host cells transformed or transfected with such expression vectors are also provided.

In another aspect, the present invention provides fusion proteins

20 comprising an inventive polypeptide, or, alternatively, an inventive polypeptide and a known *Chlamydia* antigen, as well as polynucleotides encoding such fusion proteins, in combination with a physiologically acceptable carrier or immunostimulant for use as pharmaceutical compositions and vaccines thereof.

The present invention further provides pharmaceutical compositions that

25 comprise: (a) an antibody, both polyclonal and monoclonal, or antigen-binding fragment thereof that specifically binds to a *Chlamydial* protein; and (b) a physiologically acceptable carrier. Within other aspects, the present invention provides pharmaceutical compositions that comprise one or more *Chlamydia* polypeptides disclosed herein, for example, a polypeptide of SEQ ID NO: 95-109, 122-124 and 139-30 140, or a polynucleotide molecule encoding such a polypeptide, such as a polynucleotide sequence of SEQ ID NO: 1-48, 80-94, 114-121 and 125-138, and a physiologically acceptable carrier. The invention also provides compositions for

prophylactic and therapeutic purposes comprising one or more of the disclosed polynucleotides and/or polypeptides and an immunostimulant, e.g., an adjuvant.

In yet another aspect, methods are provided for stimulating an immune response in a patient, e.g., for inducing protective immunity in a patient, comprising administering to a patient an effective amount of one or more of the above pharmaceutical compositions or vaccines.

In yet a further aspect, methods for the treatment of *Chlamydia* infection in a patient are provided, the methods comprising obtaining peripheral blood mononuclear cells (PBMC) from the patient, incubating the PBMC with a polypeptide of the present invention (or a polynucleotide that encodes such a polypeptide) to provide incubated T cells and administering the incubated T cells to the patient. The present invention additionally provides methods for the treatment of *Chlamydia* infection that comprise incubating antigen presenting cells with a polypeptide of the present invention (or a polynucleotide that encodes such a polypeptide) to provide incubated antigen presenting cells and administering the incubated antigen presenting cells to the patient. Proliferated cells may, but need not, be cloned prior to administration to the patient. In certain embodiments, the antigen presenting cells are selected from the group consisting of dendritic cells, macrophages, monocytes, B-cells, and fibroblasts. Compositions for the treatment of *Chlamydia* infection comprising T cells or antigen presenting cells that have been incubated with a polypeptide or polynucleotide of the present invention are also provided. Within related aspects, vaccines are provided that comprise: (a) an antigen presenting cell that expresses a polypeptide as described above and (b) an immunostimulant.

The present invention further provides, within other aspects, methods for removing *Chlamydial*-infected cells from a biological sample, comprising contacting a biological sample with T cells that specifically react with a *Chlamydial* protein, wherein the step of contacting is performed under conditions and for a time sufficient to permit the removal of cells expressing the protein from the sample.

Within related aspects, methods are provided for inhibiting the development of *Chlamydial* infection in a patient, comprising administering to a patient a biological sample treated as described above. In further aspects of the subject invention, methods and diagnostic kits are provided for detecting *Chlamydia* infection

in a patient. In one embodiment, the method comprises: (a) contacting a biological sample with at least one of the polypeptides or fusion proteins disclosed herein; and (b) detecting in the sample the presence of binding agents that bind to the polypeptide or fusion protein, thereby detecting *Chlamydia* infection in the biological sample. Suitable biological samples include whole blood, sputum, serum, plasma, saliva, cerebrospinal fluid and urine. In one embodiment, the diagnostic kits comprise one or more of the polypeptides or fusion proteins disclosed herein in combination with a detection reagent. In yet another embodiment, the diagnostic kits comprise either a monoclonal antibody or a polyclonal antibody that binds with a polypeptide of the present invention.

The present invention also provides methods for detecting *Chlamydia* infection comprising: (a) obtaining a biological sample from a patient; (b) contacting the sample with at least two oligonucleotide primers in a polymerase chain reaction, at least one of the oligonucleotide primers being specific for a polynucleotide sequence disclosed herein; and (c) detecting in the sample a polynucleotide sequence that amplifies in the presence of the oligonucleotide primers. In one embodiment, the oligonucleotide primer comprises at least about 10 contiguous nucleotides of a polynucleotide sequence disclosed herein, or of a sequence that hybridizes thereto.

In a further aspect, the present invention provides a method for detecting *Chlamydia* infection in a patient comprising: (a) obtaining a biological sample from the patient; (b) contacting the sample with an oligonucleotide probe specific for a polynucleotide sequence disclosed herein; and (c) detecting in the sample a polynucleotide sequence that hybridizes to the oligonucleotide probe. In one embodiment, the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a polynucleotide sequence disclosed herein, or a sequence that hybridizes thereto.

These and other aspects of the present invention will become apparent upon reference to the following detailed description. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

SEQUENCE IDENTIFIERS

SEQ ID NO:1 sets forth a DNA sequence identified for clone E4-A2-39 (CT10 positive) that is 1311 bp and contains the entire ORF for CT460 (SWIB) and a partial ORF for CT461 (yael).

SEQ ID NO:2 sets forth a DNA sequence for clone E2-B10-52 (CT10 positive) that has a 1516 bp insert that contains partial ORFs for genes CT827 (nrda-ribonucleoside reductase large chain) and CT828 (ndrB-ribonucleoside reductase small chain). These genes as were not identified in a Ct L2 library screening.

SEQ ID NO:3 sets forth a DNA sequence for clone E1-B1-80 (CT10 positive) (2397bp) that contains partial ORFs for several genes, CT812 (pmpD), CT015 (phoH ATPase), CT016 (hypothetical protein) and pGp1-D (*C. trachomatis* plasmid gene).

SEQ ID NO:4 sets forth a DNA sequence for clone E4-F9-4 (CT10, CL8, CT1, CT5, CT13, and CHH037 positive) that contains a 1094 bp insert that has a partial ORF for the gene CT316 (L7/L12 ribosomal protein) as well as a partial ORF for gene CT315 (RNA polymerase beta).

SEQ ID NO:5 sets forth a DNA sequence for clone E2-H6-40 (CT3 positive) that has a 2129 bp insert that contains the entire ORF for the gene CT288 and very small fragments of genes CT287 and CT289. Genes in this clone have not been identified in screening with a Ct L2 library.

SEQ ID NO:6 sets forth a DNA sequence for clone E5-D4-2 (CT3, CT10, CT1, CT5, CT12, and CHH037 positive) that has a 1828 bp insert that contains a partial ORF for gene CT378 (pgi), complete ORF for gene CT377 (ItuA) and a complete ORF for the gene CT376 (malate dehydrogenase). In addition, the patient lines CT10, CT1, CT5, CT12, and CHH037 also identified this clone.

SEQ ID NO:7 sets forth a DNA sequence for clone E6-C1-31 (CT3 positive) that has a 861 bp insert that contains a partial ORF for gene CT858.

SEQ ID NO:8 sets forth a DNA sequence for clone E9-E11-76 (CT3 positive) that contains a 763 bp insert that is an amino terminal region of the gene for CT798 (Glycogen synthase). This gene was not identified in a previous screening with a Ct L2 library.

SEQ ID NO:9 sets forth a DNA sequence for clone E2-A9-26 (CT1-positive) that contains part of the gene for ORF-3 which is found on the plasmid in *Chlamydia trachomatis*.

5 SEQ ID NO:10 sets forth a DNA sequence for clone E2-G8-94 (CT1-positive) that has the carboxy terminal end of Lpda gene as well as a partial ORF for CT556.

 SEQ ID NO: 11 sets forth a DNA sequence for clone E1-H1-14 (CT1-positive) that has a 1474 bp insert that contains the amino terminal part of an Lpda ORF on the complementary strand.

10 SEQ ID NO: 12 sets forth a DNA sequence for clone E1-A5-53 (CT1-positive) that contains a 2017 bp insert that has an amino terminal portion of the ORF for dnaK gene on the complementary strand, a partial ORF for the grpE gene (CT395) and a partial ORF for CT166 .

15 SEQ ID NO: 13 sets forth a DNA sequence for clone E3-A1-50 (positive on CT1 line) that is 1199 bp and contains a carboxy terminal portion of the ORF for CT622.

 SEQ ID NO: 14 sets forth a DNA sequence for clone E3-E2-22 that has 877 bp, containing a complete ORF for CT610 on the complementary strand, and was positive on both CT3 and CT10 lines.

20 SEQ ID NO: 15 sets forth the DNA sequence for clone E5-E2-10 (CT10-positive) which is 427 bp and contains a partial ORF for the major outer membrane protein omp1. SEQ ID NO: 16 sets forth the DNA sequence for clone E2-D5-89 (516bp) which is a CT10 positive clone that contains a partial ORF for pmpD gene (CT812).

25 SEQ ID NO: 17 sets forth the DNA sequence for clone E4-G9-75 (CT10-positive) which is 723 bp and contains a partial ORF for the amino terminal region of the pmpH gene (CT872).

 SEQ ID NO: 18 sets forth the DNA sequence for clone E3-F2-37 (CT10, CT3, CT11, and CT13 positive-1377bp insert) which contains a partial ORF for the tRNA-Trp (CT322) gene and a complete ORF for the gene secE (CT321).

30 SEQ ID NO: 19 sets forth the DNA sequence for clone E5-A11-8 (CT10-positive-1736 bp) which contains the complete ORF for groES (CT111) and a majority of the ORF for groEL (CT110).

SEQ ID NO: 20 sets forth the DNA sequence for clone E7-H11-61 (CT3 positive-1135 bp) which has partial inserts for *fliA* (CT061), *tyrS* (CT062), TSA (CT603) and a hypothetical protein (CT602).

5 SEQ ID NO: 21 sets forth a DNA sequence for clone E6-C8-95 which contains a 731 bp insert that was identified using the donor lines CT3, CT1, and CT12 line. This insert has a carboxy terminal half for the gene for the 60 kDa ORF.

 SEQ ID NO: 22 sets forth the DNA sequence for clone E4-D2-79 (CT3 positive) which contains a 1181 bp insert that is a partial ORF for *nrdA* gene. The ORF for this gene was also identified from clone E2-B10-52 (CT10 positive).

10 SEQ ID NO: 23 sets forth the DNA sequence for clone E1-F9-79 (167 bp; CT11 positive) which contains a partial ORF for the gene CT133 on the complementary strand. CT133 is a predicted rRNA methylase.

 SEQ ID NO: 24 sets forth the DNA sequence for clone E2-G12-52 (1265 bp; CT11 positive) which contains a partial ORF for *clpB*, a protease ATPase.

15 SEQ ID NO: 25 sets forth the DNA sequence for clone E4-H3-56 (463 bp insert; CT1 positive) which contains a partial ORF for the TSA gene (CT603) on the complementary strand.

 SEQ ID NO: 26 sets forth the DNA sequence for clone E5-E9-3 (CT1 positive) that contains a 636 bp insert partially encoding the ORF for *dnaK* like gene.
20 Part of this sequence was also identified in clone E1-A5-53.

 SEQ ID NO:27 sets forth the full-length serovar E DNA sequence of CT875.

 SEQ ID NO:28 sets for the full-length serovar E DNA sequence of CT622.

25 SEQ ID NO:29 sets forth the DNA sequence for clone E3-B4-18 (CT1 positive) that contains a 1224 bp insert containing 4 ORFs. The complete ORF for CT772, and the partial ORFs of CT771, CT191, and CT190.

 SEQ ID NO:30 sets forth the DNA sequence for the clone E9-E10-51 (CT10 positive) that contains an 883 bp insert containing two partial ORF, CT680 and
30 CT679.

SEQ ID NO:31 sets forth the DNA sequence of the clone E9-D5-8 (CT10, CTCT1, CT4, and CT11 positive) that contains a 393 bp insert containing the partial ORF for CT680.

5 SEQ ID NO:32 sets forth the DNA sequence of the clone E7-B1-16 (CT10, CT3, CT5, CT11, CT13, and CHH037 positive) that contains a 2577 bp insert containing three ORFs, two full length ORFs for CT694 and CT695 and the third containing the N-terminal portion of CT969.

SEQ ID NO:33 sets forth the DNA sequence of the clone E9-G2-93 (CT10 positive) that contains a 554 bp insert containing a partial ORF for CT178.

10 SEQ ID NO:34 sets forth the DNA sequence of the clone E5-A8-85 (CT1 positive) that contains a 1433 bp insert containing two partial ORFs for CT875 and CT001.

SEQ ID NO:35 sets forth the DNA sequence of the clone E10-C6-45 (CT3 positive) that contains a 196 bp insert containing a partial ORF for CT827.

15 SEQ ID NO:36 sets forth the DNA sequence of the clone E7-H11-10 (CT3 positive) that contains a 1990 bp insert containing the partial ORFs of CT610 and CT613 and the complete ORFs of CT611 and CT612.

SEQ ID NO:37 sets forth the DNA sequence of the clone E2-F7-11 (CT3 and CT10 positive) that contains a 2093 bp insert. It contains a large region of CT609, 20 a complete ORF for CT610 and a partial ORF for CT611.

SEQ ID NO:38 sets forth the DNA sequence of the clone E3-A3-31 (CT1 positive) that contains an 1834 bp insert containing a large region of CT622.

SEQ ID NO:39 sets forth the DNA sequence of the clone E1-G9-23 (CT3 positive) that contains an 1180 bp insert containing almost the entire ORF for 25 CT798.

SEQ ID NO:40 sets forth the DNA sequence of the clone E4-D6-21 (CT3 positive) that contains a 1297 bp insert containing the partial ORFs of CT329 and CT327 and the complete ORF of CT328.

SEQ ID NO:41 sets forth the DNA sequence of the clone E3-F3-18 (CT1 30 positive) that contains an 1141 bp insert containing the partial ORF of CT871.

SEQ ID NO:42 sets forth the DNA sequence of the clone E10-B2-57 (CT10 positive) that contains an 822 bp insert containing the complete ORF of CT066.

SEQ ID NO:43 sets forth the DNA sequence of the clone E3-F3-7 (CT1 positive) that contains a 1643 bp insert containing the partial ORFs of CT869 and CT870.

5 SEQ ID NO:44 sets forth the DNA sequence of the clone E10-H8-1 (CT3 and CT10 positive) that contains an 1862 bp insert containing the partial ORFs of CT871 and CT872.

SEQ ID NO:45 sets forth the DNA sequence of the clone E3-D10-46 (CT1, CT3, CT4, CT11, and CT12 positive) that contains a 1666 bp insert containing the partial ORFs for CT770 and CT773 and the complete ORFs for CT771 and CT722.

10 SEQ ID NO:46 sets forth the DNA sequence of the clone E2-D8-19 (CT1 positive) that contains a 2010 bp insert containing partial ORFs, ORF3 and ORF6, and complete ORFs, ORF4 and ORF5.

SEQ ID NO:47 sets forth the DNA sequence of the clone E4-C3-40 (CT10 positive) that contains a 2044 bp insert containing the partial ORF for CT827
15 and a complete ORF for CT828.

SEQ ID NO:48 sets forth the DNA sequence of the clone E3-H6-10 (CT12 positive) that contains a 3743 bp insert containing the partial ORFs for CT223 and CT229 and the complete ORFs for CT224 and CT224, CT225, CT226, CT227, and CT228.

20 SEQ ID NO:49 sets forth the DNA sequence for the Chlamydia pneumoniae homologue, CPn0454 of the Chlamydia trachomatis gene CT872.

SEQ ID NO:50 sets forth the DNA sequence for the Chlamydia pneumoniae homologue, CPn0187, of the Chlamydia trachomatis gene CT133.

25 SEQ ID NO:51 sets forth the DNA sequence for the Chlamydia pneumoniae homologue, CPn0075 of the Chlamydia trachomatis gene CT321.

SEQ ID NO:52 sets forth the DNA sequence for the Chlamydia pneumoniae homologue, CPn0074, of the Chlamydia trachomatis gene CT322.

SEQ ID NO:53 sets forth the DNA sequence for the Chlamydia pneumoniae homologue, CPn0948, of the Chlamydia trachomatis gene CT798.

30 SEQ ID NO:54 sets forth the DNA sequence for the Chlamydia pneumoniae homologue, CPn0985, of the Chlamydia trachomatis gene CT828.

SEQ ID NO:55 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0984, of the *Chlamydia trachomatis* gene CT827.

SEQ ID NO:56 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0062, of the *Chlamydia trachomatis* gene CT289.

5 SEQ ID NO:57 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn00065, of the *Chlamydia trachomatis* gene CT288.

SEQ ID NO:58 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0438, of the *Chlamydia trachomatis* gene CT287.

10 SEQ ID NO:59 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0963, of the *Chlamydia trachomatis* gene CT812.

SEQ ID NO:60 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0778, of the *Chlamydia trachomatis* gene CT603.

SEQ ID NO:61 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0503, of the *Chlamydia trachomatis* gene CT396.

15 SEQ ID NO:62 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn1016, of the *Chlamydia trachomatis* gene CT858.

SEQ ID NO:63 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0728, of the *Chlamydia trachomatis* gene CT622.

20 SEQ ID NO:64 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0557, of the *Chlamydia trachomatis* gene CT460.

SEQ ID NO:65 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0454, of the *Chlamydia trachomatis* gene CT872.

SEQ ID NO:66 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0187, of the *Chlamydia trachomatis* gene CT133.

25 SEQ ID NO:67 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0075, of the *Chlamydia trachomatis* gene CT321.

SEQ ID NO:68 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0074, of the *Chlamydia trachomatis* gene CT322.

30 SEQ ID NO:69 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0948, of the *Chlamydia trachomatis* gene CT798.

SEQ ID NO:70 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0985, of the *Chlamydia trachomatis* gene CT828.

SEQ ID NO:71 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0984, of the *Chlamydia trachomatis* gene CT827.

SEQ ID NO:72 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0062, of the *Chlamydia trachomatis* gene CT289.

5 SEQ ID NO:73 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0065, of the *Chlamydia trachomatis* gene CT288.

SEQ ID NO:74 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0438, of the *Chlamydia trachomatis* gene CT287.

10 SEQ ID NO:75 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0963, of the *Chlamydia trachomatis* gene CT812.

SEQ ID NO:76 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0778, of the *Chlamydia trachomatis* gene CT603.

SEQ ID NO:77 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn1016, of the *Chlamydia trachomatis* gene CT858.

15 SEQ ID NO:78 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0728, of the *Chlamydia trachomatis* gene CT622.

SEQ ID NO:79 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0557, of the *Chlamydia trachomatis* gene CT460.

20 SEQ ID NO:80 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT872.

SEQ ID NO:81 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT828.

SEQ ID NO:82 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT827.

25 SEQ ID NO:83 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT812.

SEQ ID NO:84 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT798.

30 SEQ ID NO:85 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT681 (MompF).

SEQ ID NO:86 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT603.

SEQ ID NO:87 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT460.

SEQ ID NO:88 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT322.

5 SEQ ID NO:89 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT321.

SEQ ID NO:90 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT289.

10 SEQ ID NO:91 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT288.

SEQ ID NO:92 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT287.

SEQ ID NO:93 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT133.

15 SEQ ID NO:94 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT113.

SEQ ID NO:95 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT872.

20 SEQ ID NO:96 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT828.

SEQ ID NO:97 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT827.

SEQ ID NO:98 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT812.

25 SEQ ID NO:99 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT798.

SEQ ID NO:100 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT681.

30 SEQ ID NO:101 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT603.

SEQ ID NO:102 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT460.

SEQ ID NO:103 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT322.

SEQ ID NO:104 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT321.

5 SEQ ID NO:105 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT289.

SEQ ID NO:106 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT288.

10 SEQ ID NO:107 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT287.

SEQ ID NO:108 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT133.

SEQ ID NO:109 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT113.

15 SEQ ID NO:110 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0695, of the *Chlamydia trachomatis* gene CT681.

SEQ ID NO:111 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0144, of the *Chlamydia trachomatis* gene CT113.

20 SEQ ID NO:112 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0695, of the *Chlamydia trachomatis* gene CT681.

SEQ ID NO:113 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0144, of the *Chlamydia trachomatis* gene CT113.

SEQ ID NO:114 sets forth the DNA sequence of the clone E7-B12-65 (CHH037 positive) that contains a 1179 bp insert containing complete ORF for 376.

25 SEQ ID NO:115 sets forth the DNA sequence of the clone E4-H9-83 (CHH037 positive) that contains the partial ORF for the heat shock protein GroEL (CT110).

SEQ ID NO:116 sets forth the DNA sequence of the clone E9-B10-52 (CHH037 positive) that contains the partial ORF for the gene *yscC* (CT674).

30 SEQ ID NO:117 sets forth the DNA sequence of the clone E7-A7-79 (CHH037 positive) that contains the complete ORF for the histone like development

gene *hctA* (CT743) and a partial ORF for the rRNA methyltransferase gene *ygca* (CT742).

SEQ ID NO:118 sets forth the DNA sequence of the clone E2-D11-18 (CHH037 positive) that contains the partial ORF for *hctA* (CT743).

5 SEQ ID NO:119 sets forth the DNA sequence for the *Chlamydia trachomatis* serovar E hypothetical protein CT694.

SEQ ID NO:120 sets forth the DNA sequence for the *Chlamydia trachomatis* serovar E hypothetical protein CT695.

10 SEQ ID NO:121 sets forth the DNA sequence for the *Chlamydia trachomatis* serovar E L1 ribosomal protein.

SEQ ID NO:122 sets forth the amino acid sequence for the *Chlamydia trachomatis* serovar E hypothetical protein CT694.

SEQ ID NO:123 sets forth the amino acid sequence for the *Chlamydia trachomatis* serovar E hypothetical protein CT695.

15 SEQ ID NO:124 sets forth the amino acid sequence for the *Chlamydia trachomatis* serovar E L1 ribosomal protein.

SEQ ID NO:125 sets forth the DNA sequence of the clone E9-H6-15 (CT3 positive) that contains the partial ORF for the *pmpB* gene (CT413).

20 SEQ ID NO:126 sets forth the DNA sequence of the clone E3-D10-87 (CT1 positive) that contains the partial ORFs for the hypothetical genes CT388 and CT389.

SEQ ID NO:127 sets forth the DNA sequence of the clone E9-D6-43 (CT3 positive) that contains the partial ORF for the CT858.

25 SEQ ID NO:128 sets forth the DNA sequence of the clone E3-D10-4 (CT1 positive) that contains the partial ORF for *pGP3-D*, an ORF encoded on the plasmid *pCHL1*.

SEQ ID NO:129 sets forth the DNA sequence of the clone E3-G8-7 (CT1 positive) that contains the partial ORFs for the CT557 (*LpdA*) and CT558 (*LipA*).

30 SEQ ID NO:130 sets forth the DNA sequence of the clone E3-F11-32 (CT1 positive) that contains the partial ORF for *pmpD* (CT812).

SEQ ID NO:131 sets forth the DNA sequence of the clone E2-F8-5 (CT12 positive) that contains the complete ORF for the 15 kDa ORF (CT442) and a partial ORF for the 60kDa ORF (CT443).

SEQ ID NO:132 sets forth the DNA sequence of the clone E2-G4-39 (CT12 positive) that contains the partial ORF for the 60kDa ORF (CT443).

SEQ ID NO:133 sets forth the DNA sequence of the clone E9-D1-16 (CT10 positive) that contains the partial ORF for pmpH (CT872).

SEQ ID NO:134 sets forth the DNA sequence of the clone E3-F3-6 (CT1 positive) that contains the partial ORFs for the genes accB (CT123), L1 ribosomal (CT125) and S9 ribosomal (CT126).

SEQ ID NO:135 sets forth the DNA sequence of the clone E2-D4-70 (CT12 positive) that contains the partial ORF for the pmpC gene (CT414).

SEQ ID NO:136 sets forth the DNA sequence of the clone E5-A1-79 (CT1 positive) that contains the partial ORF for ydhO (CT127), a complete ORF for S9 ribosomal gene (CT126), a complete ORF for the L1 ribosomal gene (CT125) and a partial ORF for accC (CT124).

SEQ ID NO:137 sets forth the DNA sequence of the clone E1-F7-16 (CT12, CT3, and CT11 positive) that contains the partial ORF for the ftsH gene (CT841) and the entire ORF for the pnp gene (CT842).

SEQ ID NO:138 sets forth the DNA sequence of the clone E1-D8-62 (CT12 positive) that contains the partial ORFs for the ftsH gene (CT841) and for the pnp gene (CT842).

SEQ ID NO:139 sets forth the amino acid sequence for the serovar E protein CT875.

SEQ ID NO:140 sets forth the amino acid sequence for the serovar E protein CT622.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

As noted above, the present invention is generally directed to compositions and methods for the diagnosis and treatment of Chlamydial infection. In one aspect, the compositions of the subject invention include polypeptides that comprise at least one immunogenic portion of a *Chlamydia* antigen, or a variant thereof.

In specific embodiments, the subject invention discloses polypeptides comprising an immunogenic portion of a *Chlamydia* antigen, wherein the *Chlamydia* antigen comprises an amino acid sequence encoded by a polynucleotide molecule including a sequence selected from the group consisting of (a) nucleotide sequences recited in SEQ ID NO:1-48, 114-121, and 125-138 (b) the complements of said nucleotide sequences, and (c) variants of such sequences.

POLYNUCLEOTIDE COMPOSITIONS

As used herein, the terms "DNA segment" and "polynucleotide" refer to a DNA molecule that has been isolated free of total genomic DNA of a particular species. Therefore, a DNA segment encoding a polypeptide refers to a DNA segment that contains one or more coding sequences yet is substantially isolated away from, or purified free from, total genomic DNA of the species from which the DNA segment is obtained. Included within the terms "DNA segment" and "polynucleotide" are DNA segments and smaller fragments of such segments, and also recombinant vectors, including, for example, plasmids, cosmids, phagemids, phage, viruses, and the like.

As will be understood by those skilled in the art, the DNA segments of this invention can include genomic sequences, extra-genomic and plasmid-encoded sequences and smaller engineered gene segments that express, or may be adapted to express, proteins, polypeptides, peptides and the like. Such segments may be naturally isolated, or modified synthetically by the hand of man.

"Isolated," as used herein, means that a polynucleotide is substantially away from other coding sequences, and that the DNA segment does not contain large portions of unrelated coding DNA, such as large chromosomal fragments or other functional genes or polypeptide coding regions. Of course, this refers to the DNA segment as originally isolated, and does not exclude genes or coding regions later added to the segment by the hand of man.

As will be recognized by the skilled artisan, polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. RNA molecules include HnRNA molecules, which contain introns and correspond to a DNA molecule in a one-to-one manner, and mRNA molecules, which do not contain introns. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide of the present

invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials.

Polynucleotides may comprise a native Chlamydia sequence or may comprise a variant, or a biological or antigenic functional equivalent of such a sequence.

- 5 Polynucleotide variants may contain one or more substitutions, additions, deletions and/or insertions, as further described below, preferably such that the immunogenicity of the encoded polypeptide is not diminished, relative to a native Chlamydia protein. The effect on the immunogenicity of the encoded polypeptide may generally be assessed as described herein. The term "variants" also encompasses homologous genes of
- 10 xenogenic origin.

- When comparing polynucleotide or polypeptide sequences, two sequences are said to be "identical" if the sequence of nucleotides or amino acids in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the
- 15 sequences over a comparison window to identify and compare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

- 20 Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M.O. (1978) A model of evolutionary change in proteins – Matrices for detecting distant relationships.
- 25 In Dayhoff, M.O. (ed.) Atlas of Protein Sequence and Structure, National Biomedical Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) Unified Approach to Alignment and Phylogenies pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M. (1989) *CABIOS* 5:151-153; Myers, E.W. and Muller W. (1988) *CABIOS* 4:11-17; Robinson,
- 30 E.D. (1971) *Comb. Theor* 11:105; Santou, N. Nes, M. (1987) *Mol. Biol. Evol.* 4:406-425; Sneath, P.H.A. and Sokal, R.R. (1973) *Numerical Taxonomy – the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, CA; Wilbur, W.J. and

Lipman, D.J. (1983) *Proc. Natl. Acad. Sci. USA* 80:726-730.

Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) *Add. APL. Math* 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity methods of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

One preferred example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1977) *Nucl. Acids Res.* 25:3389-3402 and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410, respectively. BLAST and BLAST 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides and polypeptides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. In one illustrative example, cumulative scores can be calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix can be used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915) alignments, (B) of 50, expectation (E) of 10, M=5, N=-4 and a comparison of both strands.

Preferably, the "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide or polypeptide sequence in the comparison

window may comprise additions or deletions (*i.e.*, gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid bases or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (*i.e.*, the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

Therefore, the present invention encompasses polynucleotide and polypeptide sequences having substantial identity to the sequences disclosed herein, for example those comprising at least 50% sequence identity, preferably at least 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher, sequence identity compared to a polynucleotide or polypeptide sequence of this invention using the methods described herein, (e.g., BLAST analysis using standard parameters, as described below). One skilled in this art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like.

In additional embodiments, the present invention provides isolated polynucleotides and polypeptides comprising various lengths of contiguous stretches of sequence identical to or complementary to one or more of the sequences disclosed herein. For example, polynucleotides are provided by this invention that comprise at least about 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 400, 500 or 1000 or more contiguous nucleotides of one or more of the sequences disclosed herein as well as all intermediate lengths there between. It will be readily understood that "intermediate lengths", in this context, means any length between the quoted values, such as 16, 17, 18, 19, *etc.*; 21, 22, 23, *etc.*; 30, 31, 32, *etc.*; 50, 51, 52, 53, *etc.*; 100, 101, 102, 103, *etc.*; 150, 151, 152, 153, *etc.*; including all integers through 200-500; 500-1,000, and the like.

The polynucleotides of the present invention, or fragments thereof, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, polyadenylation signals, additional restriction enzyme

sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. For
5 example, illustrative DNA segments with total lengths of about 10,000, about 5000, about 3000, about 2,000, about 1,000, about 500, about 200, about 100, about 50 base pairs in length, and the like, (including all intermediate lengths) are contemplated to be useful in many implementations of this invention.

In other embodiments, the present invention is directed to
10 polynucleotides that are capable of hybridizing under moderately stringent conditions to a polynucleotide sequence provided herein, or a fragment thereof, or a complementary sequence thereof. Hybridization techniques are well known in the art of molecular biology. For purposes of illustration, suitable moderately stringent conditions for testing the hybridization of a polynucleotide of this invention with other polynucleotides
15 include prewashing in a solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5 X SSC, overnight; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS.

Moreover, it will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences
20 that encode a polypeptide as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention. Further, alleles of the genes comprising the polynucleotide sequences provided herein are within the scope of the present invention.
25 Alleles are endogenous genes that are altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides. The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard techniques (such as hybridization, amplification and/or database sequence comparison).

PROBES AND PRIMERS

In other embodiments of the present invention, the polynucleotide sequences provided herein can be advantageously used as probes or primers for nucleic acid hybridization. As such, it is contemplated that nucleic acid segments that comprise
5 a sequence region of at least about 15 nucleotide long contiguous sequence that has the same sequence as, or is complementary to, a 15 nucleotide long contiguous sequence disclosed herein will find particular utility. Longer contiguous identical or complementary sequences, *e.g.*, those of about 20, 30, 40, 50, 100, 200, 500, 1000 (including all intermediate lengths) and even up to full length sequences will also be of
10 use in certain embodiments.

The ability of such nucleic acid probes to specifically hybridize to a sequence of interest will enable them to be of use in detecting the presence of complementary sequences in a given sample. However, other uses are also envisioned, such as the use of the sequence information for the preparation of mutant species
15 primers, or primers for use in preparing other genetic constructions.

Polynucleotide molecules having sequence regions consisting of contiguous nucleotide stretches of 10-14, 15-20, 30, 50, or even of 100-200 nucleotides or so (including intermediate lengths as well), identical or complementary to a polynucleotide sequence disclosed herein, are particularly contemplated as hybridization
20 probes for use in, *e.g.*, Southern and Northern blotting. This would allow a gene product, or fragment thereof, to be analyzed, both in diverse cell types and also in various bacterial cells. The total size of fragment, as well as the size of the complementary stretch(es), will ultimately depend on the intended use or application of the particular nucleic acid segment. Smaller fragments will generally find use in
25 hybridization embodiments, wherein the length of the contiguous complementary region may be varied, such as between about 15 and about 100 nucleotides, but larger contiguous complementarity stretches may be used, according to the length complementary sequences one wishes to detect.

The use of a hybridization probe of about 15-25 nucleotides in length
30 allows the formation of a duplex molecule that is both stable and selective. Molecules having contiguous complementary sequences over stretches greater than 15 bases in length are generally preferred, though, in order to increase stability and selectivity of the

hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained. One will generally prefer to design nucleic acid molecules having gene-complementary stretches of 15 to 25 contiguous nucleotides, or even longer where desired.

5 Hybridization probes may be selected from any portion of any of the sequences disclosed herein. All that is required is to review the sequence set forth in SEQ ID NO:1-48, 114-121, and 125-138, or to any continuous portion of the sequence, from about 15-25 nucleotides in length up to and including the full length sequence, that one wishes to utilize as a probe or primer. The choice of probe and primer sequences
10 may be governed by various factors. For example, one may wish to employ primers from towards the termini of the total sequence.

Small polynucleotide segments or fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer. Also, fragments may be
15 obtained by application of nucleic acid reproduction technology, such as the PCRTM technology of U. S. Patent 4,683,202 (incorporated herein by reference), by introducing selected sequences into recombinant vectors for recombinant production, and by other recombinant DNA techniques generally known to those of skill in the art of molecular biology.

20 The nucleotide sequences of the invention may be used for their ability to selectively form duplex molecules with complementary stretches of the entire gene or gene fragments of interest. Depending on the application envisioned, one will typically desire to employ varying conditions of hybridization to achieve varying degrees of selectivity of probe towards target sequence. For applications requiring high selectivity,
25 one will typically desire to employ relatively stringent conditions to form the hybrids, *e.g.*, one will select relatively low salt and/or high temperature conditions, such as provided by a salt concentration of from about 0.02 M to about 0.15 M salt at temperatures of from about 50°C to about 70°C. Such selective conditions tolerate little, if any, mismatch between the probe and the template or target strand, and would
30 be particularly suitable for isolating related sequences.

Of course, for some applications, for example, where one desires to prepare mutants employing a mutant primer strand hybridized to an underlying

template, less stringent (reduced stringency) hybridization conditions will typically be needed in order to allow formation of the heteroduplex. In these circumstances, one may desire to employ salt conditions such as those of from about 0.15 M to about 0.9 M salt, at temperatures ranging from about 20°C to about 55°C. Cross-hybridizing species can thereby be readily identified as positively hybridizing signals with respect to control hybridizations. In any case, it is generally appreciated that conditions can be rendered more stringent by the addition of increasing amounts of formamide, which serves to destabilize the hybrid duplex in the same manner as increased temperature. Thus, hybridization conditions can be readily manipulated, and thus will generally be a method of choice depending on the desired results.

POLYNUCLEOTIDE IDENTIFICATION AND CHARACTERIZATION

Polynucleotides may be identified, prepared and/or manipulated using any of a variety of well established techniques. For example, a polynucleotide may be identified, by screening a microarray of cDNAs for Chlamydia expression. Such screens may be performed, for example, using a Synteni microarray (Palo Alto, CA) according to the manufacturer's instructions (and essentially as described by Schena et al., *Proc. Natl. Acad. Sci. USA* 93:10614-10619, 1996 and Heller et al., *Proc. Natl. Acad. Sci. USA* 94:2150-2155, 1997). Alternatively, polynucleotides may be amplified from cDNA prepared from cells expressing the proteins described herein. Such polynucleotides may be amplified via polymerase chain reaction (PCR). For this approach, sequence-specific primers may be designed based on the sequences provided herein, and may be purchased or synthesized.

An amplified portion of a polynucleotide of the present invention may be used to isolate a full length gene from a suitable library (e.g., Chlamydia eDNA library) using well known techniques. Within such techniques, a library (cDNA or genomic) is screened using one or more polynucleotide probes or primers suitable for amplification. Preferably, a library is size-selected to include larger molecules. Random primed libraries may also be preferred for identifying 5' and upstream regions of genes. Genomic libraries are preferred for obtaining introns and extending 5' sequences.

For hybridization techniques, a partial sequence may be labeled (e.g., by nick-translation or end-labeling with ³²P) using well known techniques. A bacterial or

bacteriophage library is then generally screened by hybridizing filters containing denatured bacterial colonies (or lawns containing phage plaques) with the labeled probe (see Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989). Hybridizing colonies or plaques are
5 selected and expanded, and the DNA is isolated for further analysis. cDNA clones may be analyzed to determine the amount of additional sequence by, for example, PCR using a primer from the partial sequence and a primer from the vector. Restriction maps and partial sequences may be generated to identify one or more overlapping clones. The complete sequence may then be determined using standard techniques, which may
10 involve generating a series of deletion clones. The resulting overlapping sequences can then be assembled into a single contiguous sequence. A full length cDNA molecule can be generated by ligating suitable fragments, using well known techniques.

Alternatively, there are numerous amplification techniques for obtaining a full length coding sequence from a partial cDNA sequence. Within such techniques,
15 amplification is generally performed via PCR. Any of a variety of commercially available kits may be used to perform the amplification step. Primers may be designed using, for example, software well known in the art. Primers are preferably 22-30 nucleotides in length, have a GC content of at least 50% and anneal to the target sequence at temperatures of about 68°C to 72°C. The amplified region may be
20 sequenced as described above, and overlapping sequences assembled into a contiguous sequence.

One such amplification technique is inverse PCR (see Triglia et al., *Nucl. Acids Res.* 16:8186, 1988), which uses restriction enzymes to generate a fragment in the known region of the gene. The fragment is then circularized by intramolecular ligation
25 and used as a template for PCR with divergent primers derived from the known region. Within an alternative approach, sequences adjacent to a partial sequence may be retrieved by amplification with a primer to a linker sequence and a primer specific to a known region. The amplified sequences are typically subjected to a second round of amplification with the same linker primer and a second primer specific to the known
30 region. A variation on this procedure, which employs two primers that initiate extension in opposite directions from the known sequence, is described in WO 96/38591. Another such technique is known as "rapid amplification of cDNA ends" or

RACE. This technique involves the use of an internal primer and an external primer, which hybridizes to a polyA region or vector sequence, to identify sequences that are 5' and 3' of a known sequence. Additional techniques include capture PCR (Lagerstrom et al., *PCR Methods Applic. 1*:111-19, 1991) and walking PCR (Parker et al., *Nucl. Acids. Res. 19*:3055-60, 1991). Other methods employing amplification may also be employed to obtain a full length cDNA sequence.

In certain instances, it is possible to obtain a full length cDNA sequence by analysis of sequences provided in an expressed sequence tag (EST) database, such as that available from GenBank. Searches for overlapping ESTs may generally be performed using well known programs (e.g., NCBI BLAST searches), and such ESTs may be used to generate a contiguous full length sequence. Full length DNA sequences may also be obtained by analysis of genomic fragments.

POLYNUCLEOTIDE EXPRESSION IN HOST CELLS

In other embodiments of the invention, polynucleotide sequences or fragments thereof which encode polypeptides of the invention, or fusion proteins or functional equivalents thereof, may be used in recombinant DNA molecules to direct expression of a polypeptide in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences that encode substantially the same or a functionally equivalent amino acid sequence may be produced and these sequences may be used to clone and express a given polypeptide.

As will be understood by those of skill in the art, it may be advantageous in some instances to produce polypeptide-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce a recombinant RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

Moreover, the polynucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter polypeptide encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. For example, DNA shuffling by random fragmentation and PCR reassembly of gene

fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. In addition, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, or introduce mutations, and so forth.

5 In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences may be ligated to a heterologous sequence to encode a fusion protein. For example, to screen peptide libraries for inhibitors of polypeptide activity, it may be useful to encode a chimeric protein that can be recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site
10 located between the polypeptide-encoding sequence and the heterologous protein sequence, so that the polypeptide may be cleaved and purified away from the heterologous moiety.

Sequences encoding a desired polypeptide may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers, M. H. et al.
15 (1980) *Nucl. Acids Res. Symp. Ser.* 215-223, Horn, T. et al. (1980) *Nucl. Acids Res. Symp. Ser.* 225-232). Alternatively, the protein itself may be produced using chemical methods to synthesize the amino acid sequence of a polypeptide, or a portion thereof. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge, J. Y. et al. (1995) *Science* 269:202-204) and automated synthesis may be
20 achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer, Palo Alto, CA).

A newly synthesized peptide may be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, T. (1983) *Proteins, Structures and Molecular Principles*, WH Freeman and Co., New York, N.Y.) or other comparable
25 techniques available in the art. The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure). Additionally, the amino acid sequence of a polypeptide, or any part thereof, may be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

30 In order to express a desired polypeptide, the nucleotide sequences encoding the polypeptide, or functional equivalents, may be inserted into appropriate expression

vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding a polypeptide of interest and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described in Sambrook, J. et al. (1989) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Plainview, N.Y., and Ausubel, F. M. et al. (1989) *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, N.Y.

A variety of expression vector/host systems may be utilized to contain and express polynucleotide sequences. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems.

The "control elements" or "regulatory sequences" present in an expression vector are those non-translated regions of the vector--enhancers, promoters, 5' and 3' untranslated regions--which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the PBLUESCRIPT phagemid (Stratagene, La Jolla, Calif.) or PSPORT1 plasmid (Gibco BRL, Gaithersburg, MD) and the like may be used. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are generally preferred. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding a polypeptide, vectors based on SV40 or EBV may be advantageously used with an appropriate selectable marker.

In bacterial systems, a number of expression vectors may be selected

depending upon the use intended for the expressed polypeptide. For example, when large quantities are needed, for example for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and
5 expression vectors such as BLUESCRIPT (Stratagene), in which the sequence encoding the polypeptide of interest may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke, G. and S. M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509); and the like. pGEX Vectors (Promega, Madison, Wis.)
10 may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide
15 of interest can be released from the GST moiety at will.

In the yeast, *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al. (1987) *Methods Enzymol.* 153:516-544.

20 In cases where plant expression vectors are used, the expression of sequences encoding polypeptides may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311. Alternatively, plant promoters such as the small subunit of
25 RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, for example, Hobbs, S. or
30 Murry, L. E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, N.Y.; pp. 191-196).

An insect system may also be used to express a polypeptide of interest. For example, in one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The sequences encoding the polypeptide may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the polypeptide-encoding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, *S. frugiperda* cells or *Trichoplusia* larvae in which the polypeptide of interest may be expressed (Engelhard, E. K. et al. (1994) *Proc. Natl. Acad. Sci.* 91 :3224-3227).

In mammalian host cells, a number of viral-based expression systems are generally available. For example, in cases where an adenovirus is used as an expression vector, sequences encoding a polypeptide of interest may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing the polypeptide in infected host cells (Logan, J. and Shenk, T. (1984) *Proc. Natl. Acad. Sci.* 81:3655-3659). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

Specific initiation signals may also be used to achieve more efficient translation of sequences encoding a polypeptide of interest. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding the polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation.

- 5 Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as CHO, HeLa, MDCK, HEK293, and WI38, which have specific cellular machinery and characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

- 10 For long-term, high-yield production of recombinant proteins, stable expression is generally preferred. For example, cell lines which stably express a polynucleotide of interest may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be
15 allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

- 20 Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. et al. (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. et al. (1990) *Cell* 22:817-23) genes which can be employed in tk.sup.- or aprt.sup.- cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can
25 be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler, M. et al. (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); npt, which confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin, F. et al (1981) *J. Mol. Biol.* 150:1-14); and als or pat, which confer resistance to chloresulfuron and phosphinotricin acetyltransferase, respectively (Murry, *supra*).
30 Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman, S. C. and R. C. Mulligan (1988) *Proc. Natl. Acad. Sci.*

85:8047-51). Recently, the use of visible markers has gained popularity with such markers as anthocyanins, beta-glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific
5 vector system (Rhodes, C. A. et al. (1995) *Methods Mol. Biol.* 55:121-131).

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the sequence encoding a polypeptide is inserted within a marker gene sequence, recombinant cells containing sequences can be identified by the
10 absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a polypeptide-encoding sequence under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

Alternatively, host cells which contain and express a desired polynucleotide
15 sequence may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein.

A variety of protocols for detecting and measuring the expression of polynucleotide-encoded products, using either polyclonal or monoclonal antibodies specific for the product are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal
20 antibodies reactive to two non-interfering epitopes on a given polypeptide may be preferred for some applications, but a competitive binding assay may also be employed. These and other assays are described, among other places, in Hampton, R. et al. (1990; *Serological Methods, a Laboratory Manual*, APS Press, St Paul, Minn.) and Maddox, D. E. et al. (1983; *J. Exp. Med.* 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled
30 in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to

polynucleotides include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences, or any portions thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits. Suitable reporter molecules or labels, which may be used include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with a polynucleotide sequence of interest may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides of the invention may be designed to contain signal sequences which direct secretion of the encoded polypeptide through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding a polypeptide of interest to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, Calif.) between the purification domain and the encoded polypeptide may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing a polypeptide of interest and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography) as described in Porath, J. et al. (1992, *Prot. Exp. Purif.* 3:263-281) while the enterokinase cleavage site provides a means for purifying the desired polypeptide from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D. J.

et al. (1993; *DNA Cell Biol.* 12:441-453).

In addition to recombinant production methods, polypeptides of the invention, and fragments thereof, may be produced by direct peptide synthesis using solid-phase techniques (Merrifield J. (1963) *J. Am. Chem. Soc.* 85:2149-2154). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Alternatively, various fragments may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

SITE-SPECIFIC MUTAGENESIS

Site-specific mutagenesis is a technique useful in the preparation of individual peptides, or biologically functional equivalent polypeptides, through specific mutagenesis of the underlying polynucleotides that encode them. The technique, well-known to those of skill in the art, further provides a ready ability to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the DNA. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Mutations may be employed in a selected polynucleotide sequence to improve, alter, decrease, modify, or otherwise change the properties of the polynucleotide itself, and/or alter the properties, activity, composition, stability, or primary sequence of the encoded polypeptide.

In certain embodiments of the present invention, the inventors contemplate the mutagenesis of the disclosed polynucleotide sequences to alter one or more properties of the encoded polypeptide, such as the antigenicity of a polypeptide vaccine. The techniques of site-specific mutagenesis are well-known in the art, and are widely used to create variants of both polypeptides and polynucleotides. For example, site-specific mutagenesis is often used to alter a specific portion of a DNA molecule. In such embodiments, a primer comprising typically about 14 to about 25 nucleotides or so

in length is employed, with about 5 to about 10 residues on both sides of the junction of the sequence being altered.

As will be appreciated by those of skill in the art, site-specific mutagenesis techniques have often employed a phage vector that exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage are readily commercially-available and their use is generally well-known to those skilled in the art. Double-stranded plasmids are also routinely employed in site directed mutagenesis that eliminates the step of transferring the gene of interest from a plasmid to a phage.

In general, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector or melting apart of two strands of a double-stranded vector that includes within its sequence a DNA sequence that encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement.

The preparation of sequence variants of the selected peptide-encoding DNA segments using site-directed mutagenesis provides a means of producing potentially useful species and is not meant to be limiting as there are other ways in which sequence variants of peptides and the DNA sequences encoding them may be obtained. For example, recombinant vectors encoding the desired peptide sequence may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants. Specific details regarding these methods and protocols are found in the teachings of Maloy *et al.*, 1994; Segal, 1976; Prokop and Bajpai, 1991; Kuby, 1994; and Maniatis *et al.*, 1982, each incorporated herein by reference, for that purpose.

As used herein, the term "oligonucleotide directed mutagenesis procedure" refers to template-dependent processes and vector-mediated propagation which result in an increase in the concentration of a specific nucleic acid molecule

relative to its initial concentration, or in an increase in the concentration of a detectable signal, such as amplification. As used herein, the term "oligonucleotide directed mutagenesis procedure" is intended to refer to a process that involves the template-dependent extension of a primer molecule. The term template dependent process refers to nucleic acid synthesis of an RNA or a DNA molecule wherein the sequence of the newly synthesized strand of nucleic acid is dictated by the well-known rules of complementary base pairing (see, for example, Watson, 1987). Typically, vector mediated methodologies involve the introduction of the nucleic acid fragment into a DNA or RNA vector, the clonal amplification of the vector, and the recovery of the amplified nucleic acid fragment. Examples of such methodologies are provided by U. S. Patent No. 4,237,224, specifically incorporated herein by reference in its entirety.

POLYNUCLEOTIDE AMPLIFICATION TECHNIQUES

A number of template dependent processes are available to amplify the target sequences of interest present in a sample. One of the best known amplification methods is the polymerase chain reaction (PCRTM) which is described in detail in U.S. Patent Nos. 4,683,195, 4,683,202 and 4,800,159, each of which is incorporated herein by reference in its entirety. Briefly, in PCRTM, two primer sequences are prepared which are complementary to regions on opposite complementary strands of the target sequence. An excess of deoxynucleoside triphosphates is added to a reaction mixture along with a DNA polymerase (e.g., *Taq* polymerase). If the target sequence is present in a sample, the primers will bind to the target and the polymerase will cause the primers to be extended along the target sequence by adding on nucleotides. By raising and lowering the temperature of the reaction mixture, the extended primers will dissociate from the target to form reaction products, excess primers will bind to the target and to the reaction product and the process is repeated. Preferably reverse transcription and PCRTM amplification procedure may be performed in order to quantify the amount of mRNA amplified. Polymerase chain reaction methodologies are well known in the art.

Another method for amplification is the ligase chain reaction (referred to as LCR), disclosed in Eur. Pat. Appl. Publ. No. 320,308 (specifically incorporated herein by reference in its entirety). In LCR, two complementary probe pairs are

prepared, and in the presence of the target sequence, each pair will bind to opposite complementary strands of the target such that they abut. In the presence of a ligase, the two probe pairs will link to form a single unit. By temperature cycling, as in PCR™, bound ligated units dissociate from the target and then serve as "target sequences" for
5 ligation of excess probe pairs. U.S. Patent No. 4,883,750, incorporated herein by reference in its entirety, describes an alternative method of amplification similar to LCR for binding probe pairs to a target sequence.

Qbeta Replicase, described in PCT Intl. Pat. Appl. Publ. No. PCT/US87/00880, incorporated herein by reference in its entirety, may also be used as
10 still another amplification method in the present invention. In this method, a replicative sequence of RNA that has a region complementary to that of a target is added to a sample in the presence of an RNA polymerase. The polymerase will copy the replicative sequence that can then be detected.

An isothermal amplification method, in which restriction endonucleases
15 and ligases are used to achieve the amplification of target molecules that contain nucleotide 5'-[α -thio]triphosphates in one strand of a restriction site (Walker *et al.*, 1992, incorporated herein by reference in its entirety), may also be useful in the amplification of nucleic acids in the present invention.

Strand Displacement Amplification (SDA) is another method of carrying
20 out isothermal amplification of nucleic acids which involves multiple rounds of strand displacement and synthesis, *i.e.* nick translation. A similar method, called Repair Chain Reaction (RCR) is another method of amplification which may be useful in the present invention and is involves annealing several probes throughout a region targeted for amplification, followed by a repair reaction in which only two of the four bases are
25 present. The other two bases can be added as biotinylated derivatives for easy detection. A similar approach is used in SDA.

Sequences can also be detected using a cyclic probe reaction (CPR). In CPR, a probe having a 3' and 5' sequences of non-target DNA and an internal or "middle" sequence of the target protein specific RNA is hybridized to DNA which is
30 present in a sample. Upon hybridization, the reaction is treated with RNaseH, and the products of the probe are identified as distinctive products by generating a signal that is released after digestion. The original template is annealed to another cycling probe and

the reaction is repeated. Thus, CPR involves amplifying a signal generated by hybridization of a probe to a target gene specific expressed nucleic acid.

Still other amplification methods described in Great Britain Pat. Appl. No. 2 202 328, and in PCT Intl. Pat. Appl. Publ. No. PCT/US89/01025, each of which is incorporated herein by reference in its entirety, may be used in accordance with the present invention. In the former application, "modified" primers are used in a PCR-like, template and enzyme dependent synthesis. The primers may be modified by labeling with a capture moiety (e.g., biotin) and/or a detector moiety (e.g., enzyme). In the latter application, an excess of labeled probes is added to a sample. In the presence of the target sequence, the probe binds and is cleaved catalytically. After cleavage, the target sequence is released intact to be bound by excess probe. Cleavage of the labeled probe signals the presence of the target sequence.

Other nucleic acid amplification procedures include transcription-based amplification systems (TAS) (Kwoh *et al.*, 1989; PCT Intl. Pat. Appl. Publ. No. WO 88/10315, incorporated herein by reference in its entirety), including nucleic acid sequence based amplification (NASBA) and 3SR. In NASBA, the nucleic acids can be prepared for amplification by standard phenol/chloroform extraction, heat denaturation of a sample, treatment with lysis buffer and minispin columns for isolation of DNA and RNA or guanidinium chloride extraction of RNA. These amplification techniques involve annealing a primer that has sequences specific to the target sequence. Following polymerization, DNA/RNA hybrids are digested with RNase H while double stranded DNA molecules are heat-denatured again. In either case the single stranded DNA is made fully double stranded by addition of second target-specific primer, followed by polymerization. The double stranded DNA molecules are then multiply transcribed by a polymerase such as T7 or SP6. In an isothermal cyclic reaction, the RNAs are reverse transcribed into DNA, and transcribed once again with a polymerase such as T7 or SP6. The resulting products, whether truncated or complete, indicate target-specific sequences.

Eur. Pat. Appl. Publ. No. 329,822, incorporated herein by reference in its entirety, disclose a nucleic acid amplification process involving cyclically synthesizing single-stranded RNA ("ssRNA"), ssDNA, and double-stranded DNA (dsDNA), which may be used in accordance with the present invention. The ssRNA is a first template

for a first primer oligonucleotide, which is elongated by reverse transcriptase (RNA-dependent DNA polymerase). The RNA is then removed from resulting DNA:RNA duplex by the action of ribonuclease H (RNase H, an RNase specific for RNA in a duplex with either DNA or RNA). The resultant ssDNA is a second template
5 for a second primer, which also includes the sequences of an RNA polymerase promoter (exemplified by T7 RNA polymerase) 5' to its homology to its template. This primer is then extended by DNA polymerase (exemplified by the large "Klenow" fragment of *E. coli* DNA polymerase I), resulting as a double-stranded DNA ("dsDNA") molecule, having a sequence identical to that of the original RNA between the primers and having
10 additionally, at one end, a promoter sequence. This promoter sequence can be used by the appropriate RNA polymerase to make many RNA copies of the DNA. These copies can then re-enter the cycle leading to very swift amplification. With proper choice of enzymes, this amplification can be done isothermally without addition of enzymes at each cycle. Because of the cyclical nature of this process, the starting sequence can be
15 chosen to be in the form of either DNA or RNA.

PCT Intl. Pat. Appl. Publ. No. WO 89/06700, incorporated herein by reference in its entirety, disclose a nucleic acid sequence amplification scheme based on the hybridization of a promoter/primer sequence to a target single-stranded DNA ("ssDNA") followed by transcription of many RNA copies of the sequence. This
20 scheme is not cyclic; i.e. new templates are not produced from the resultant RNA transcripts. Other amplification methods include "RACE" (Frohman, 1990), and "one-sided PCR" (Ohara, 1989) which are well-known to those of skill in the art.

Methods based on ligation of two (or more) oligonucleotides in the presence of nucleic acid having the sequence of the resulting "di-oligonucleotide",
25 thereby amplifying the di-oligonucleotide (Wu and Dean, 1996, incorporated herein by reference in its entirety), may also be used in the amplification of DNA sequences of the present invention.

BIOLOGICAL FUNCTIONAL EQUIVALENTS

30 Modification and changes may be made in the structure of the polynucleotides and polypeptides of the present invention and still obtain a functional molecule that encodes a polypeptide with desirable characteristics. As mentioned

above, it is often desirable to introduce one or more mutations into a specific polynucleotide sequence. In certain circumstances, the resulting encoded polypeptide sequence is altered by this mutation, or in other cases, the sequence of the polypeptide is unchanged by one or more mutations in the encoding polynucleotide.

5 When it is desirable to alter the amino acid sequence of a polypeptide to create an equivalent, or even an improved, second-generation molecule, the amino acid changes may be achieved by changing one or more of the codons of the encoding DNA sequence, according to Table 1.

10 For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's biological functional activity, certain amino acid sequence substitutions can be made in a protein sequence, and, of course, its underlying DNA
15 coding sequence, and nevertheless obtain a protein with like properties. It is thus contemplated by the inventors that various changes may be made in the peptide sequences of the disclosed compositions, or corresponding DNA sequences which encode said peptides without appreciable loss of their biological utility or activity.

TABLE 1

Amino Acids			Codons						
Alanine	Ala	A	GCA	GCC	GCG	GCU			
Cysteine	Cys	C	UGC	UGU					
Aspartic acid	Asp	D	GAC	GAU					
Glutamic acid	Glu	E	GAA	GAG					
Phenylalanine	Phe	F	UUC	UUU					
Glycine	Gly	G	GGA	GGC	GGG	GGU			
Histidine	His	H	CAC	CAU					
Isoleucine	Ile	I	AUA	AUC	AUU				
Lysine	Lys	K	AAA	AAG					
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG	CUU	
Methionine	Met	M	AUG						
Asparagine	Asn	N	AAC	AAU					
Proline	Pro	P	CCA	CCC	CCG	CCU			
Glutamine	Gln	Q	CAA	CAG					
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGU	
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG	UCU	
Threonine	Thr	T	ACA	ACC	ACG	ACU			
Valine	Val	V	GUA	GUC	GUG	GUU			
Tryptophan	Trp	W	UGG						
Tyrosine	Tyr	Y	UAC	UAU					

In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte and Doolittle, 1982, incorporated herein by reference). It is accepted that the relative hydropathic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics (Kyte and Doolittle, 1982). These values are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (−0.4); threonine (−0.7); serine (−0.8);

tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydrophatic index or score and still result in a protein with similar biological activity, *i.e.* still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydrophatic indices are within ± 2 is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred. It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U. S. Patent 4,554,101 (specifically incorporated herein by reference in its entirety), states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein.

As detailed in U. S. Patent 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 \pm 1); glutamate (+3.0 \pm 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5 \pm 1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4). It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent, and in particular, an immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within ± 2 is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

As outlined above, amino acid substitutions are generally therefore based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions that take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

In addition, any polynucleotide may be further modified to increase stability *in vivo*. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends; the use of phosphorothioate or 2' O-methyl

rather than phosphodiesterase linkages in the backbone; and/or the inclusion of nontraditional bases such as inosine, queosine and wybutosine, as well as acetyl-methyl-, thio- and other modified forms of adenine, cytidine, guanine, thymine and uridine.

5

IN VIVO POLYNUCLEOTIDE DELIVERY TECHNIQUES

In additional embodiments, genetic constructs comprising one or more of the polynucleotides of the invention are introduced into cells *in vivo*. This may be achieved using any of a variety of well known approaches, several of which are outlined
10 below for the purpose of illustration.

I. ADENOVIRUS

One of the preferred methods for *in vivo* delivery of one or more nucleic acid sequences involves the use of an adenovirus expression vector. "Adenovirus expression vector" is meant to include those constructs containing adenovirus sequences
15 sufficient to (a) support packaging of the construct and (b) to express a polynucleotide that has been cloned therein in a sense or antisense orientation. Of course, in the context of an antisense construct, expression does not require that the gene product be synthesized.

The expression vector comprises a genetically engineered form of an
20 adenovirus. Knowledge of the genetic organization of adenovirus, a 36 kb, linear, double-stranded DNA virus, allows substitution of large pieces of adenoviral DNA with foreign sequences up to 7 kb (Grunhaus and Horwitz, 1992). In contrast to retrovirus, the adenoviral infection of host cells does not result in chromosomal integration because adenoviral DNA can replicate in an episomal manner without potential genotoxicity.
25 Also, adenoviruses are structurally stable, and no genome rearrangement has been detected after extensive amplification. Adenovirus can infect virtually all epithelial cells regardless of their cell cycle stage. So far, adenoviral infection appears to be linked only to mild disease such as acute respiratory disease in humans.

Adenovirus is particularly suitable for use as a gene transfer vector
30 because of its mid-sized genome, ease of manipulation, high titer, wide target-cell range and high infectivity. Both ends of the viral genome contain 100-200 base pair inverted repeats (ITRs), which are *cis* elements necessary for viral DNA replication and

packaging. The early (E) and late (L) regions of the genome contain different transcription units that are divided by the onset of viral DNA replication. The E1 region (E1A and E1B) encodes proteins responsible for the regulation of transcription of the viral genome and a few cellular genes. The expression of the E2 region (E2A and E2B) results in the synthesis of the proteins for viral DNA replication. These proteins are involved in DNA replication, late gene expression and host cell shut-off (Renan, 1990). The products of the late genes, including the majority of the viral capsid proteins, are expressed only after significant processing of a single primary transcript issued by the major late promoter (MLP). The MLP, (located at 16.8 m.u.) is particularly efficient during the late phase of infection, and all the mRNA's issued from this promoter possess a 5'-tripartite leader (TPL) sequence which makes them preferred mRNA's for translation.

In a current system, recombinant adenovirus is generated from homologous recombination between shuttle vector and provirus vector. Due to the possible recombination between two proviral vectors, wild-type adenovirus may be generated from this process. Therefore, it is critical to isolate a single clone of virus from an individual plaque and examine its genomic structure.

Generation and propagation of the current adenovirus vectors, which are replication deficient, depend on a unique helper cell line, designated 293, which was transformed from human embryonic kidney cells by Ad5 DNA fragments and constitutively expresses E1 proteins (Graham *et al.*, 1977). Since the E3 region is dispensable from the adenovirus genome (Jones and Shenk, 1978), the current adenovirus vectors, with the help of 293 cells, carry foreign DNA in either the E1, the D3 or both regions (Graham and Prevec, 1991). In nature, adenovirus can package approximately 105% of the wild-type genome (Ghosh-Choudhury *et al.*, 1987), providing capacity for about 2 extra kB of DNA. Combined with the approximately 5.5 kB of DNA that is replaceable in the E1 and E3 regions, the maximum capacity of the current adenovirus vector is under 7.5 kB, or about 15% of the total length of the vector. More than 80% of the adenovirus viral genome remains in the vector backbone and is the source of vector-borne cytotoxicity. Also, the replication deficiency of the E1-deleted virus is incomplete. For example, leakage of viral gene expression has been

observed with the currently available vectors at high multiplicities of infection (MOI) (Mulligan, 1993).

Helper cell lines may be derived from human cells such as human embryonic kidney cells, muscle cells, hematopoietic cells or other human embryonic mesenchymal or epithelial cells. Alternatively, the helper cells may be derived from the cells of other mammalian species that are permissive for human adenovirus. Such cells include, e.g., Vero cells or other monkey embryonic mesenchymal or epithelial cells. As stated above, the currently preferred helper cell line is 293.

Recently, Racher *et al.* (1995) disclosed improved methods for culturing 293 cells and propagating adenovirus. In one format, natural cell aggregates are grown by inoculating individual cells into 1 liter siliconized spinner flasks (Techne, Cambridge, UK) containing 100-200 ml of medium. Following stirring at 40 rpm, the cell viability is estimated with trypan blue. In another format, Fibra-Cel microcarriers (Bibby Sterlin, Stone, UK) (5 g/l) is employed as follows. A cell inoculum, resuspended in 5 ml of medium, is added to the carrier (50 ml) in a 250 ml Erlenmeyer flask and left stationary, with occasional agitation, for 1 to 4 h. The medium is then replaced with 50 ml of fresh medium and shaking initiated. For virus production, cells are allowed to grow to about 80% confluence, after which time the medium is replaced (to 25% of the final volume) and adenovirus added at an MOI of 0.05. Cultures are left stationary overnight, following which the volume is increased to 100% and shaking commenced for another 72 h.

Other than the requirement that the adenovirus vector be replication defective, or at least conditionally defective, the nature of the adenovirus vector is not believed to be crucial to the successful practice of the invention. The adenovirus may be of any of the 42 different known serotypes or subgroups A-F. Adenovirus type 5 of subgroup C is the preferred starting material in order to obtain a conditional replication-defective adenovirus vector for use in the present invention, since Adenovirus type 5 is a human adenovirus about which a great deal of biochemical and genetic information is known, and it has historically been used for most constructions employing adenovirus as a vector.

As stated above, the typical vector according to the present invention is replication defective and will not have an adenovirus E1 region. Thus, it will be most

convenient to introduce the polynucleotide encoding the gene of interest at the position from which the E1-coding sequences have been removed. However, the position of insertion of the construct within the adenovirus sequences is not critical to the invention. The polynucleotide encoding the gene of interest may also be inserted in lieu
5 of the deleted E3 region in E3 replacement vectors as described by Karlsson *et al.* (1986) or in the E4 region where a helper cell line or helper virus complements the E4 defect.

Adenovirus is easy to grow and manipulate and exhibits broad host range *in vitro* and *in vivo*. This group of viruses can be obtained in high titers, *e.g.*, 10^9 - 10^{11}
10 plaque-forming units per ml, and they are highly infective. The life cycle of adenovirus does not require integration into the host cell genome. The foreign genes delivered by adenovirus vectors are episomal and, therefore, have low genotoxicity to host cells. No side effects have been reported in studies of vaccination with wild-type adenovirus (Couch *et al.*, 1963; Top *et al.*, 1971), demonstrating their safety and therapeutic
15 potential as *in vivo* gene transfer vectors.

Adenovirus vectors have been used in eukaryotic gene expression (Leverro *et al.*, 1991; Gomez-Foix *et al.*, 1992) and vaccine development (Grunhaus and Horwitz, 1992; Graham and Prevec, 1992). Recently, animal studies suggested that recombinant adenovirus could be used for gene therapy (Stratford-Perricaudet and
20 Perricaudet, 1991; Stratford-Perricaudet *et al.*, 1990; Rich *et al.*, 1993). Studies in administering recombinant adenovirus to different tissues include trachea instillation (Rosenfeld *et al.*, 1991; Rosenfeld *et al.*, 1992), muscle injection (Ragot *et al.*, 1993), peripheral intravenous injections (Herz and Gerard, 1993) and stereotactic inoculation into the brain (Le Gal La Salle *et al.*, 1993).

25 2. RETROVIRUSES

The retroviruses are a group of single-stranded RNA viruses characterized by an ability to convert their RNA to double-stranded DNA in infected cells by a process of reverse-transcription (Coffin, 1990). The resulting DNA then stably integrates into cellular chromosomes as a provirus and directs synthesis of viral
30 proteins. The integration results in the retention of the viral gene sequences in the recipient cell and its descendants. The retroviral genome contains three genes, gag, pol, and env that code for capsid proteins, polymerase enzyme, and envelope components,

respectively. A sequence found upstream from the gag gene contains a signal for packaging of the genome into virions. Two long terminal repeat (LTR) sequences are present at the 5' and 3' ends of the viral genome. These contain strong promoter and enhancer sequences and are also required for integration in the host cell genome
5 (Coffin, 1990).

In order to construct a retroviral vector, a nucleic acid encoding one or more oligonucleotide or polynucleotide sequences of interest is inserted into the viral genome in the place of certain viral sequences to produce a virus that is replication-defective. In order to produce virions, a packaging cell line containing the gag, pol, and
10 env genes but without the LTR and packaging components is constructed (Mann *et al.*, 1983). When a recombinant plasmid containing a cDNA, together with the retroviral LTR and packaging sequences is introduced into this cell line (by calcium phosphate precipitation for example), the packaging sequence allows the RNA transcript of the recombinant plasmid to be packaged into viral particles, which are then secreted into the
15 culture media (Nicolas and Rubenstein, 1988; Temin, 1986; Mann *et al.*, 1983). The media containing the recombinant retroviruses is then collected, optionally concentrated, and used for gene transfer. Retroviral vectors are able to infect a broad variety of cell types. However, integration and stable expression require the division of host cells (Paskind *et al.*, 1975).

20 A novel approach designed to allow specific targeting of retrovirus vectors was recently developed based on the chemical modification of a retrovirus by the chemical addition of lactose residues to the viral envelope. This modification could permit the specific infection of hepatocytes *via* sialoglycoprotein receptors.

A different approach to targeting of recombinant retroviruses was
25 designed in which biotinylated antibodies against a retroviral envelope protein and against a specific cell receptor were used. The antibodies were coupled *via* the biotin components by using streptavidin (Roux *et al.*, 1989). Using antibodies against major histocompatibility complex class I and class II antigens, they demonstrated the infection of a variety of human cells that bore those surface antigens with an ecotropic virus *in*
30 *vitro* (Roux *et al.*, 1989).

3. ADENO-ASSOCIATED VIRUSES

AAV (Ridgeway, 1988; Hermonat and Muzyczka, 1984) is a parovirus, discovered as a contamination of adenoviral stocks. It is a ubiquitous virus (antibodies are present in 85% of the US human population) that has not been linked to any disease. It is also classified as a dependovirus, because its replications is dependent on the presence of a helper virus, such as adenovirus. Five serotypes have been isolated, of which AAV-2 is the best characterized. AAV has a single-stranded linear DNA that is encapsidated into capsid proteins VP1, VP2 and VP3 to form an icosahedral virion of 20 to 24 nm in diameter (Muzyczka and McLaughlin, 1988).

The AAV DNA is approximately 4.7 kilobases long. It contains two open reading frames and is flanked by two ITRs (FIG. 2). There are two major genes in the AAV genome: *rep* and *cap*. The *rep* gene codes for proteins responsible for viral replications, whereas *cap* codes for capsid protein VP1-3. Each ITR forms a T-shaped hairpin structure. These terminal repeats are the only essential *cis* components of the AAV for chromosomal integration. Therefore, the AAV can be used as a vector with all viral coding sequences removed and replaced by the cassette of genes for delivery. Three viral promoters have been identified and named p5, p19, and p40, according to their map position. Transcription from p5 and p19 results in production of rep proteins, and transcription from p40 produces the capsid proteins (Hermonat and Muzyczka, 1984).

There are several factors that prompted researchers to study the possibility of using rAAV as an expression vector. One is that the requirements for delivering a gene to integrate into the host chromosome are surprisingly few. It is necessary to have the 145-bp ITRs, which are only 6% of the AAV genome. This leaves room in the vector to assemble a 4.5-kb DNA insertion. While this carrying capacity may prevent the AAV from delivering large genes, it is amply suited for delivering the antisense constructs of the present invention.

AAV is also a good choice of delivery vehicles due to its safety. There is a relatively complicated rescue mechanism: not only wild type adenovirus but also AAV genes are required to mobilize rAAV. Likewise, AAV is not pathogenic and not associated with any disease. The removal of viral coding sequences minimizes immune

reactions to viral gene expression, and therefore, rAAV does not evoke an inflammatory response.

4. OTHER VIRAL VECTORS AS EXPRESSION CONSTRUCTS

Other viral vectors may be employed as expression constructs in the present invention for the delivery of oligonucleotide or polynucleotide sequences to a host cell. Vectors derived from viruses such as vaccinia virus (Ridgeway, 1988; Coupar *et al.*, 1988), lentiviruses, polio viruses and herpes viruses may be employed. They offer several attractive features for various mammalian cells (Friedmann, 1989; Ridgeway, 1988; Coupar *et al.*, 1988; Horwich *et al.*, 1990).

With the recent recognition of defective hepatitis B viruses, new insight was gained into the structure-function relationship of different viral sequences. *In vitro* studies showed that the virus could retain the ability for helper-dependent packaging and reverse transcription despite the deletion of up to 80% of its genome (Horwich *et al.*, 1990). This suggested that large portions of the genome could be replaced with foreign genetic material. The hepatotropism and persistence (integration) were particularly attractive properties for liver-directed gene transfer. Chang *et al.* (1991) introduced the chloramphenicol acetyltransferase (CAT) gene into duck hepatitis B virus genome in the place of the polymerase, surface, and pre-surface coding sequences. It was cotransfected with wild-type virus into an avian hepatoma cell line. Culture media containing high titers of the recombinant virus were used to infect primary duckling hepatocytes. Stable CAT gene expression was detected for at least 24 days after transfection (Chang *et al.*, 1991).

5. NON-VIRAL VECTORS

In order to effect expression of the oligonucleotide or polynucleotide sequences of the present invention, the expression construct must be delivered into a cell. This delivery may be accomplished *in vitro*, as in laboratory procedures for transforming cells lines, or *in vivo* or *ex vivo*, as in the treatment of certain disease states. As described above, one preferred mechanism for delivery is *via* viral infection where the expression construct is encapsulated in an infectious viral particle.

Once the expression construct has been delivered into the cell the nucleic acid encoding the desired oligonucleotide or polynucleotide sequences may be positioned and expressed at different sites. In certain embodiments, the nucleic acid

encoding the construct may be stably integrated into the genome of the cell. This integration may be in the specific location and orientation *via* homologous recombination (gene replacement) or it may be integrated in a random, non-specific location (gene augmentation). In yet further embodiments, the nucleic acid may be stably maintained in the cell as a separate, episomal segment of DNA. Such nucleic acid segments or "episomes" encode sequences sufficient to permit maintenance and replication independent of or in synchronization with the host cell cycle. How the expression construct is delivered to a cell and where in the cell the nucleic acid remains is dependent on the type of expression construct employed.

In certain embodiments of the invention, the expression construct comprising one or more oligonucleotide or polynucleotide sequences may simply consist of naked recombinant DNA or plasmids. Transfer of the construct may be performed by any of the methods mentioned above which physically or chemically permeabilize the cell membrane. This is particularly applicable for transfer *in vitro* but it may be applied to *in vivo* use as well. Dubensky *et al.* (1984) successfully injected polyomavirus DNA in the form of calcium phosphate precipitates into liver and spleen of adult and newborn mice demonstrating active viral replication and acute infection. Benvenisty and Reshef (1986) also demonstrated that direct intraperitoneal injection of calcium phosphate-precipitated plasmids results in expression of the transfected genes. It is envisioned that DNA encoding a gene of interest may also be transferred in a similar manner *in vivo* and express the gene product.

Another embodiment of the invention for transferring a naked DNA expression construct into cells may involve particle bombardment. This method depends on the ability to accelerate DNA-coated microprojectiles to a high velocity allowing them to pierce cell membranes and enter cells without killing them (Klein *et al.*, 1987). Several devices for accelerating small particles have been developed. One such device relies on a high voltage discharge to generate an electrical current, which in turn provides the motive force (Yang *et al.*, 1990). The microprojectiles used have consisted of biologically inert substances such as tungsten or gold beads.

Selected organs including the liver, skin, and muscle tissue of rats and mice have been bombarded *in vivo* (Yang *et al.*, 1990; Zelenin *et al.*, 1991). This may require surgical exposure of the tissue or cells, to eliminate any intervening tissue

between the gun and the target organ, *i.e. ex vivo* treatment. Again, DNA encoding a particular gene may be delivered *via* this method and still be incorporated by the present invention.

5 ANTISENSE OLIGONUCLEOTIDES

The end result of the flow of genetic information is the synthesis of protein. DNA is transcribed by polymerases into messenger RNA and translated on the ribosome to yield a folded, functional protein. Thus there are several steps along the route where protein synthesis can be inhibited. The native DNA segment coding for a polypeptide described herein, as all such mammalian DNA strands, has two strands: a sense strand and an antisense strand held together by hydrogen bonding. The messenger RNA coding for polypeptide has the same nucleotide sequence as the sense DNA strand except that the DNA thymidine is replaced by uridine. Thus, synthetic antisense nucleotide sequences will bind to a mRNA and inhibit expression of the protein encoded by that mRNA.

The targeting of antisense oligonucleotides to mRNA is thus one mechanism to shut down protein synthesis, and, consequently, represents a powerful and targeted therapeutic approach. For example, the synthesis of polygalacturonase and the muscarine type 2 acetylcholine receptor are inhibited by antisense oligonucleotides directed to their respective mRNA sequences (U. S. Patent 5,739,119 and U. S. Patent 5,759,829, each specifically incorporated herein by reference in its entirety). Further, examples of antisense inhibition have been demonstrated with the nuclear protein cyclin, the multiple drug resistance gene (MDG1), ICAM-1, E-selectin, STK-1, striatal GABA_A receptor and human EGF (Jaskulski *et al.*, 1988; Vasanthakumar and Ahmed, 1989; Peris *et al.*, 1998; U. S. Patent 5,801,154; U. S. Patent 5,789,573; U. S. Patent 5,718,709 and U. S. Patent 5,610,288, each specifically incorporated herein by reference in its entirety). Antisense constructs have also been described that inhibit and can be used to treat a variety of abnormal cellular proliferations, *e.g.* cancer (U. S. Patent 5,747,470; U. S. Patent 5,591,317 and U. S. Patent 5,783,683, each specifically incorporated herein by reference in its entirety).

Therefore, in exemplary embodiments, the invention provides oligonucleotide sequences that comprise all, or a portion of, any sequence that is

capable of specifically binding to polynucleotide sequence described herein, or a complement thereof. In one embodiment, the antisense oligonucleotides comprise DNA or derivatives thereof. In another embodiment, the oligonucleotides comprise RNA or derivatives thereof. In a third embodiment, the oligonucleotides are modified DNAs
5 comprising a phosphorothioated modified backbone. In a fourth embodiment, the oligonucleotide sequences comprise peptide nucleic acids or derivatives thereof. In each case, preferred compositions comprise a sequence region that is complementary, and more preferably substantially-complementary, and even more preferably, completely complementary to one or more portions of polynucleotides disclosed herein.

10 Selection of antisense compositions specific for a given gene sequence is based upon analysis of the chosen target sequence (*i.e.* in these illustrative examples the rat and human sequences) and determination of secondary structure, T_m , binding energy, relative stability, and antisense compositions were selected based upon their relative inability to form dimers, hairpins, or other secondary structures that would reduce or
15 prohibit specific binding to the target mRNA in a host cell.

Highly preferred target regions of the mRNA, are those which are at or near the AUG translation initiation codon, and those sequences which were substantially complementary to 5' regions of the mRNA. These secondary structure analyses and target site selection considerations were performed using v.4 of the
20 OLIGO primer analysis software (Rychlik, 1997) and the BLASTN 2.0.5 algorithm software (Altschul *et al.*, 1997).

The use of an antisense delivery method employing a short peptide vector, termed MPG (27 residues), is also contemplated. The MPG peptide contains a hydrophobic domain derived from the fusion sequence of HIV gp41 and a hydrophilic
25 domain from the nuclear localization sequence of SV40 T-antigen (Morris *et al.*, 1997). It has been demonstrated that several molecules of the MPG peptide coat the antisense oligonucleotides and can be delivered into cultured mammalian cells in less than 1 hour with relatively high efficiency (90%). Further, the interaction with MPG strongly increases both the stability of the oligonucleotide to nuclease and the ability to cross the
30 plasma membrane (Morris *et al.*, 1997).

RIBOZYMES

Although proteins traditionally have been used for catalysis of nucleic acids, another class of macromolecules has emerged as useful in this endeavor. Ribozymes are RNA-protein complexes that cleave nucleic acids in a site-specific fashion. Ribozymes have specific catalytic domains that possess endonuclease activity (Kim and Cech, 1987; Gerlach *et al.*, 1987; Forster and Symons, 1987). For example, a large number of ribozymes accelerate phosphoester transfer reactions with a high degree of specificity, often cleaving only one of several phosphoesters in an oligonucleotide substrate (Cech *et al.*, 1981; Michel and Westhof, 1990; Reinhold-Hurek and Shub, 1992). This specificity has been attributed to the requirement that the substrate bind via specific base-pairing interactions to the internal guide sequence ("IGS") of the ribozyme prior to chemical reaction.

Ribozyme catalysis has primarily been observed as part of sequence-specific cleavage/ligation reactions involving nucleic acids (Joyce, 1989; Cech *et al.*, 1981). For example, U. S. Patent No. 5,354,855 (specifically incorporated herein by reference) reports that certain ribozymes can act as endonucleases with a sequence specificity greater than that of known ribonucleases and approaching that of the DNA restriction enzymes. Thus, sequence-specific ribozyme-mediated inhibition of gene expression may be particularly suited to therapeutic applications (Scanlon *et al.*, 1991; Sarver *et al.*, 1990). Recently, it was reported that ribozymes elicited genetic changes in some cells lines to which they were applied; the altered genes included the oncogenes *H-ras*, *c-fos* and genes of HIV. Most of this work involved the modification of a target mRNA, based on a specific mutant codon that is cleaved by a specific ribozyme.

Six basic varieties of naturally-occurring enzymatic RNAs are known presently. Each can catalyze the hydrolysis of RNA phosphodiester bonds *in trans* (and thus can cleave other RNA molecules) under physiological conditions. In general, enzymatic nucleic acids act by first binding to a target RNA. Such binding occurs through the target binding portion of a enzymatic nucleic acid which is held in close proximity to an enzymatic portion of the molecule that acts to cleave the target RNA. Thus, the enzymatic nucleic acid first recognizes and then binds a target RNA through complementary base-pairing, and once bound to the correct site, acts enzymatically to cut the target RNA. Strategic cleavage of such a target RNA will destroy its ability to

direct synthesis of an encoded protein. After an enzymatic nucleic acid has bound and cleaved its RNA target, it is released from that RNA to search for another target and can repeatedly bind and cleave new targets.

The enzymatic nature of a ribozyme is advantageous over many technologies, such as antisense technology (where a nucleic acid molecule simply binds to a nucleic acid target to block its translation) since the concentration of ribozyme necessary to affect a therapeutic treatment is lower than that of an antisense oligonucleotide. This advantage reflects the ability of the ribozyme to act enzymatically. Thus, a single ribozyme molecule is able to cleave many molecules of target RNA. In addition, the ribozyme is a highly specific inhibitor, with the specificity of inhibition depending not only on the base pairing mechanism of binding to the target RNA, but also on the mechanism of target RNA cleavage. Single mismatches, or base-substitutions, near the site of cleavage can completely eliminate catalytic activity of a ribozyme. Similar mismatches in antisense molecules do not prevent their action (Woolf *et al.*, 1992). Thus, the specificity of action of a ribozyme is greater than that of an antisense oligonucleotide binding the same RNA site.

The enzymatic nucleic acid molecule may be formed in a hammerhead, hairpin, a hepatitis δ virus, group I intron or RNaseP RNA (in association with an RNA guide sequence) or Neurospora VS RNA motif. Examples of hammerhead motifs are described by Rossi *et al.* (1992). Examples of hairpin motifs are described by Hampel *et al.* (Eur. Pat. Appl. Publ. No. EP 0360257), Hampel and Tritz (1989), Hampel *et al.* (1990) and U. S. Patent 5,631,359 (specifically incorporated herein by reference). An example of the hepatitis δ virus motif is described by Perrotta and Been (1992); an example of the RNaseP motif is described by Guerrier-Takada *et al.* (1983); Neurospora VS RNA ribozyme motif is described by Collins (Saville and Collins, 1990; Saville and Collins, 1991; Collins and Olive, 1993); and an example of the Group I intron is described in (U. S. Patent 4,987,071, specifically incorporated herein by reference). All that is important in an enzymatic nucleic acid molecule of this invention is that it has a specific substrate binding site which is complementary to one or more of the target gene RNA regions, and that it have nucleotide sequences within or surrounding that substrate binding site which impart an RNA cleaving activity to the molecule. Thus the ribozyme constructs need not be limited to specific motifs mentioned herein.

In certain embodiments, it may be important to produce enzymatic cleaving agents which exhibit a high degree of specificity for the RNA of a desired target, such as one of the sequences disclosed herein. The enzymatic nucleic acid molecule is preferably targeted to a highly conserved sequence region of a target mRNA. Such enzymatic nucleic acid molecules can be delivered exogenously to specific cells as required. Alternatively, the ribozymes can be expressed from DNA or RNA vectors that are delivered to specific cells.

Small enzymatic nucleic acid motifs (e.g., of the hammerhead or the hairpin structure) may also be used for exogenous delivery. The simple structure of these molecules increases the ability of the enzymatic nucleic acid to invade targeted regions of the mRNA structure. Alternatively, catalytic RNA molecules can be expressed within cells from eukaryotic promoters (e.g., Scanlon *et al.*, 1991; Kashani-Sabet *et al.*, 1992; Dropulic *et al.*, 1992; Weerasinghe *et al.*, 1991; Ojwang *et al.*, 1992; Chen *et al.*, 1992; Sarver *et al.*, 1990). Those skilled in the art realize that any ribozyme can be expressed in eukaryotic cells from the appropriate DNA vector. The activity of such ribozymes can be augmented by their release from the primary transcript by a second ribozyme (Int. Pat. Appl. Publ. No. WO 93/23569, and Int. Pat. Appl. Publ. No. WO 94/02595, both hereby incorporated by reference; Ohkawa *et al.*, 1992; Taira *et al.*, 1991; and Ventura *et al.*, 1993).

Ribozymes may be added directly, or can be complexed with cationic lipids, lipid complexes, packaged within liposomes, or otherwise delivered to target cells. The RNA or RNA complexes can be locally administered to relevant tissues *ex vivo*, or *in vivo* through injection, aerosol inhalation, infusion pump or stent, with or without their incorporation in biopolymers.

Ribozymes may be designed as described in Int. Pat. Appl. Publ. No. WO 93/23569 and Int. Pat. Appl. Publ. No. WO 94/02595, each specifically incorporated herein by reference) and synthesized to be tested *in vitro* and *in vivo*, as described. Such ribozymes can also be optimized for delivery. While specific examples are provided, those in the art will recognize that equivalent RNA targets in other species can be utilized when necessary.

Hammerhead or hairpin ribozymes may be individually analyzed by computer folding (Jaeger *et al.*, 1989) to assess whether the ribozyme sequences fold

into the appropriate secondary structure. Those ribozymes with unfavorable intramolecular interactions between the binding arms and the catalytic core are eliminated from consideration. Varying binding arm lengths can be chosen to optimize activity. Generally, at least 5 or so bases on each arm are able to bind to, or otherwise
5 interact with, the target RNA.

Ribozymes of the hammerhead or hairpin motif may be designed to anneal to various sites in the mRNA message, and can be chemically synthesized. The method of synthesis used follows the procedure for normal RNA synthesis as described in Usman *et al.* (1987) and in Scaringe *et al.* (1990) and makes use of common nucleic
10 acid protecting and coupling groups, such as dimethoxytrityl at the 5'-end, and phosphoramidites at the 3'-end. Average stepwise coupling yields are typically >98%. Hairpin ribozymes may be synthesized in two parts and annealed to reconstruct an active ribozyme (Chowrira and Burke, 1992). Ribozymes may be modified extensively to enhance stability by modification with nuclease resistant groups, for example, 2'-
15 amino, 2'-C-allyl, 2'-fluoro, 2'-O-methyl, 2'-H (for a review see *e.g.*, Usman and Cedergren, 1992). Ribozymes may be purified by gel electrophoresis using general methods or by high pressure liquid chromatography and resuspended in water.

Ribozyme activity can be optimized by altering the length of the ribozyme binding arms, or chemically synthesizing ribozymes with modifications that
20 prevent their degradation by serum ribonucleases (see *e.g.*, Int. Pat. Appl. Publ. No. WO 92/07065; Perrault *et al.*, 1990; Pieken *et al.*, 1991; Usman and Cedergren, 1992; Int. Pat. Appl. Publ. No. WO 93/15187; Int. Pat. Appl. Publ. No. WO 91/03162; Eur. Pat. Appl. Publ. No. 92110298.4; U. S. Patent 5,334,711; and Int. Pat. Appl. Publ. No. WO 94/13688, which describe various chemical modifications that can be made to the sugar
25 moieties of enzymatic RNA molecules), modifications which enhance their efficacy in cells, and removal of stem II bases to shorten RNA synthesis times and reduce chemical requirements.

Sullivan *et al.* (Int. Pat. Appl. Publ. No. WO 94/02595) describes the general methods for delivery of enzymatic RNA molecules. Ribozymes may be
30 administered to cells by a variety of methods known to those familiar to the art, including, but not restricted to, encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins, biodegradable

nanocapsules, and bioadhesive microspheres. For some indications, ribozymes may be directly delivered *ex vivo* to cells or tissues with or without the aforementioned vehicles. Alternatively, the RNA/vehicle combination may be locally delivered by direct inhalation, by direct injection or by use of a catheter, infusion pump or stent. Other routes of delivery include, but are not limited to, intravascular, intramuscular, subcutaneous or joint injection, aerosol inhalation, oral (tablet or pill form), topical, systemic, ocular, intraperitoneal and/or intrathecal delivery. More detailed descriptions of ribozyme delivery and administration are provided in Int. Pat. Appl. Publ. No. WO 94/02595 and Int. Pat. Appl. Publ. No. WO 93/23569, each specifically incorporated herein by reference.

Another means of accumulating high concentrations of a ribozyme(s) within cells is to incorporate the ribozyme-encoding sequences into a DNA expression vector. Transcription of the ribozyme sequences are driven from a promoter for eukaryotic RNA polymerase I (pol I), RNA polymerase II (pol II), or RNA polymerase III (pol III). Transcripts from pol II or pol III promoters will be expressed at high levels in all cells; the levels of a given pol II promoter in a given cell type will depend on the nature of the gene regulatory sequences (enhancers, silencers, *etc.*) present nearby. Prokaryotic RNA polymerase promoters may also be used, providing that the prokaryotic RNA polymerase enzyme is expressed in the appropriate cells (Elroy-Stein and Moss, 1990; Gao and Huang, 1993; Lieber *et al.*, 1993; Zhou *et al.*, 1990). Ribozymes expressed from such promoters can function in mammalian cells (e.g. Kashani-Saber *et al.*, 1992; Ojwang *et al.*, 1992; Chen *et al.*, 1992; Yu *et al.*, 1993; L'Huillier *et al.*, 1992; Lisiewicz *et al.*, 1993). Such transcription units can be incorporated into a variety of vectors for introduction into mammalian cells, including but not restricted to, plasmid DNA vectors, viral DNA vectors (such as adenovirus or adeno-associated vectors), or viral RNA vectors (such as retroviral, semliki forest virus, sindbis virus vectors).

Ribozymes may be used as diagnostic tools to examine genetic drift and mutations within diseased cells. They can also be used to assess levels of the target RNA molecule. The close relationship between ribozyme activity and the structure of the target RNA allows the detection of mutations in any region of the molecule which alters the base-pairing and three-dimensional structure of the target RNA. By using

multiple ribozymes, one may map nucleotide changes which are important to RNA structure and function *in vitro*, as well as in cells and tissues. Cleavage of target RNAs with ribozymes may be used to inhibit gene expression and define the role (essentially) of specified gene products in the progression of disease. In this manner, other genetic targets may be defined as important mediators of the disease. These studies will lead to better treatment of the disease progression by affording the possibility of combinational therapies (e.g., multiple ribozymes targeted to different genes, ribozymes coupled with known small molecule inhibitors, or intermittent treatment with combinations of ribozymes and/or other chemical or biological molecules). Other *in vitro* uses of ribozymes are well known in the art, and include detection of the presence of mRNA associated with an IL-5 related condition. Such RNA is detected by determining the presence of a cleavage product after treatment with a ribozyme using standard methodology.

15 PEPTIDE NUCLEIC ACIDS

In certain embodiments, the inventors contemplate the use of peptide nucleic acids (PNAs) in the practice of the methods of the invention. PNA is a DNA mimic in which the nucleobases are attached to a pseudopeptide backbone (Good and Nielsen, 1997). PNA is able to be utilized in a number of methods that traditionally have used RNA or DNA. Often PNA sequences perform better in techniques than the corresponding RNA or DNA sequences and have utilities that are not inherent to RNA or DNA. A review of PNA including methods of making, characteristics of, and methods of using, is provided by Corey (1997) and is incorporated herein by reference. As such, in certain embodiments, one may prepare PNA sequences that are complementary to one or more portions of the ACE mRNA sequence, and such PNA compositions may be used to regulate, alter, decrease, or reduce the translation of ACE-specific mRNA, and thereby alter the level of ACE activity in a host cell to which such PNA compositions have been administered.

PNAs have 2-aminoethyl-glycine linkages replacing the normal phosphodiester backbone of DNA (Nielsen *et al.*, 1991; Hanvey *et al.*, 1992; Hyrup and Nielsen, 1996; Nielsen, 1996). This chemistry has three important consequences: firstly, in contrast to DNA or phosphorothioate oligonucleotides, PNAs are neutral

molecules; secondly, PNAs are achiral, which avoids the need to develop a stereoselective synthesis; and thirdly, PNA synthesis uses standard Boc (Dueholm *et al.*, 1994) or Fmoc (Thomson *et al.*, 1995) protocols for solid-phase peptide synthesis, although other methods, including a modified Merrifield method, have been used
5 (Christensen *et al.*, 1995).

PNA monomers or ready-made oligomers are commercially available from PerSeptive Biosystems (Framingham, MA). PNA syntheses by either Boc or Fmoc protocols are straightforward using manual or automated protocols (Norton *et al.*, 1995). The manual protocol lends itself to the production of chemically modified PNAs
10 or the simultaneous synthesis of families of closely related PNAs.

As with peptide synthesis, the success of a particular PNA synthesis will depend on the properties of the chosen sequence. For example, while in theory PNAs can incorporate any combination of nucleotide bases, the presence of adjacent purines can lead to deletions of one or more residues in the product. In expectation of this
15 difficulty, it is suggested that, in producing PNAs with adjacent purines, one should repeat the coupling of residues likely to be added inefficiently. This should be followed by the purification of PNAs by reverse-phase high-pressure liquid chromatography (Norton *et al.*, 1995) providing yields and purity of product similar to those observed during the synthesis of peptides.

20 Modifications of PNAs for a given application may be accomplished by coupling amino acids during solid-phase synthesis or by attaching compounds that contain a carboxylic acid group to the exposed N-terminal amine. Alternatively, PNAs can be modified after synthesis by coupling to an introduced lysine or cysteine. The ease with which PNAs can be modified facilitates optimization for better solubility or
25 for specific functional requirements. Once synthesized, the identity of PNAs and their derivatives can be confirmed by mass spectrometry. Several studies have made and utilized modifications of PNAs (Norton *et al.*, 1995; Haaima *et al.*, 1996; Stetsenko *et al.*, 1996; Petersen *et al.*, 1995; Ulmann *et al.*, 1996; Koch *et al.*, 1995; Orum *et al.*, 1995; Footer *et al.*, 1996; Griffith *et al.*, 1995; Kremsky *et al.*, 1996; Pardridge *et al.*,
30 1995; Boffa *et al.*, 1995; Landsdorp *et al.*, 1996; Gambacorti-Passerini *et al.*, 1996; Armitage *et al.*, 1997; Seeger *et al.*, 1997; Ruskowski *et al.*, 1997). U.S. Patent No. 5,700,922 discusses PNA-DNA-PNA chimeric molecules and their uses in diagnostics,

modulating protein in organisms, and treatment of conditions susceptible to therapeutics.

In contrast to DNA and RNA, which contain negatively charged linkages, the PNA backbone is neutral. In spite of this dramatic alteration, PNAs recognize complementary DNA and RNA by Watson-Crick pairing (Egholm *et al.*, 1993), validating the initial modeling by Nielsen *et al.* (1991). PNAs lack 3' to 5' polarity and can bind in either parallel or antiparallel fashion, with the antiparallel mode being preferred (Egholm *et al.*, 1993).

Hybridization of DNA oligonucleotides to DNA and RNA is destabilized by electrostatic repulsion between the negatively charged phosphate backbones of the complementary strands. By contrast, the absence of charge repulsion in PNA-DNA or PNA-RNA duplexes increases the melting temperature (T_m) and reduces the dependence of T_m on the concentration of mono- or divalent cations (Nielsen *et al.*, 1991). The enhanced rate and affinity of hybridization are significant because they are responsible for the surprising ability of PNAs to perform strand invasion of complementary sequences within relaxed double-stranded DNA. In addition, the efficient hybridization at inverted repeats suggests that PNAs can recognize secondary structure effectively within double-stranded DNA. Enhanced recognition also occurs with PNAs immobilized on surfaces, and Wang *et al.* have shown that support-bound PNAs can be used to detect hybridization events (Wang *et al.*, 1996).

One might expect that tight binding of PNAs to complementary sequences would also increase binding to similar (but not identical) sequences, reducing the sequence specificity of PNA recognition. As with DNA hybridization, however, selective recognition can be achieved by balancing oligomer length and incubation temperature. Moreover, selective hybridization of PNAs is encouraged by PNA-DNA hybridization being less tolerant of base mismatches than DNA-DNA hybridization. For example, a single mismatch within a 16 bp PNA-DNA duplex can reduce the T_m by up to 15°C (Egholm *et al.*, 1993). This high level of discrimination has allowed the development of several PNA-based strategies for the analysis of point mutations (Wang *et al.*, 1996; Carlsson *et al.*, 1996; Thiede *et al.*, 1996; Webb and Hurskainen, 1996; Perry-O'Keefe *et al.*, 1996).

High-affinity binding provides clear advantages for molecular recognition and the development of new applications for PNAs. For example, 11-13 nucleotide PNAs inhibit the activity of telomerase, a ribonucleo-protein that extends telomere ends using an essential RNA template, while the analogous DNA oligomers do not (Norton *et al.*, 1996).

Neutral PNAs are more hydrophobic than analogous DNA oligomers, and this can lead to difficulty solubilizing them at neutral pH, especially if the PNAs have a high purine content or if they have the potential to form secondary structures. Their solubility can be enhanced by attaching one or more positive charges to the PNA termini (Nielsen *et al.*, 1991).

Findings by Allfrey and colleagues suggest that strand invasion will occur spontaneously at sequences within chromosomal DNA (Boffa *et al.*, 1995; Boffa *et al.*, 1996). These studies targeted PNAs to triplet repeats of the nucleotides CAG and used this recognition to purify transcriptionally active DNA (Boffa *et al.*, 1995) and to inhibit transcription (Boffa *et al.*, 1996). This result suggests that if PNAs can be delivered within cells then they will have the potential to be general sequence-specific regulators of gene expression. Studies and reviews concerning the use of PNAs as antisense and anti-gene agents include Nielsen *et al.* (1993b), Hanvey *et al.* (1992), and Good and Nielsen (1997). Koppelhus *et al.* (1997) have used PNAs to inhibit HIV-1 inverse transcription, showing that PNAs may be used for antiviral therapies.

Methods of characterizing the antisense binding properties of PNAs are discussed in Rose (1993) and Jensen *et al.* (1997). Rose uses capillary gel electrophoresis to determine binding of PNAs to their complementary oligonucleotide, measuring the relative binding kinetics and stoichiometry. Similar types of measurements were made by Jensen *et al.* using BIAcore™ technology.

Other applications of PNAs include use in DNA strand invasion (Nielsen *et al.*, 1991), antisense inhibition (Hanvey *et al.*, 1992), mutational analysis (Orum *et al.*, 1993), enhancers of transcription (Mollegaard *et al.*, 1994), nucleic acid purification (Orum *et al.*, 1995), isolation of transcriptionally active genes (Boffa *et al.*, 1995), blocking of transcription factor binding (Vickers *et al.*, 1995), genome cleavage (Veselkov *et al.*, 1996), biosensors (Wang *et al.*, 1996), *in situ* hybridization (Thisted *et al.*, 1996), and in a alternative to Southern blotting (Perry-O'Keefe, 1996).

POLYPEPTIDE COMPOSITIONS AND USES

The present invention, in other aspects, provides polypeptide compositions. Generally, a polypeptide of the invention will be an isolated polypeptide (or an epitope, variant, or active fragment thereof) derived from a mammalian species. Preferably, the polypeptide is encoded by a polynucleotide sequence disclosed herein or a sequence which hybridizes under moderately stringent conditions to a polynucleotide sequence disclosed herein. Alternatively, the polypeptide may be defined as a polypeptide which comprises a contiguous amino acid sequence from an amino acid sequence disclosed herein, or which polypeptide comprises an entire amino acid sequence disclosed herein.

Likewise, a polypeptide composition of the present invention is understood to comprise one or more polypeptides that are capable of eliciting antibodies that are immunologically reactive with one or more polypeptides encoded by one or more contiguous nucleic acid sequences contained in SEQ ID NO:1-48, 114-121, and 125-138, or to active fragments, or to variants thereof, or to one or more nucleic acid sequences which hybridize to one or more of these sequences under conditions of moderate to high stringency.

As used herein, an active fragment of a polypeptide includes a whole or a portion of a polypeptide which is modified by conventional techniques, *e.g.*, mutagenesis, or by addition, deletion, or substitution, but which active fragment exhibits substantially the same structure function, antigenicity, etc., as a polypeptide as described herein.

In certain illustrative embodiments, the polypeptides of the invention will comprise at least an immunogenic portion of a Chlamydia protein or a variant thereof, as described herein. Proteins that are Chlamydia proteins generally also react detectably within an immunoassay (such as an ELISA) with antisera from a patient with a Chlamydial infection. Polypeptides as described herein may be of any length. Additional sequences derived from the native protein and/or heterologous sequences may be present, and such sequences may (but need not) possess further immunogenic or antigenic properties.

An "immunogenic portion," as used herein is a portion of a protein that

is recognized (*i.e.*, specifically bound) by a B-cell and/or T-cell surface antigen receptor. Such immunogenic portions generally comprise at least 5 amino acid residues, more preferably at least 10, and still more preferably at least 20 amino acid residues of a Chlamydia protein or a variant thereof. Certain preferred immunogenic portions include
5 peptides in which an N-terminal leader sequence and/or transmembrane domain have been deleted. Other preferred immunogenic portions may contain a small N- and/or C-terminal deletion (*e.g.*, 1-30 amino acids, preferably 5-15 amino acids), relative to the mature protein.

Immunogenic portions may generally be identified using well known
10 techniques, such as those summarized in Paul, *Fundamental Immunology*, 3rd ed., 243-247 (Raven Press, 1993) and references cited therein. Such techniques include screening polypeptides for the ability to react with antigen-specific antibodies, antisera and/or T-cell lines or clones. As used herein, antisera and antibodies are "antigen-specific" if they specifically bind to an antigen (*i.e.*, they react with the protein in an
15 ELISA or other immunoassay, and do not react detectably with unrelated proteins). Such antisera and antibodies may be prepared as described herein, and using well known techniques. An immunogenic portion of a native Chlamydia protein is a portion that reacts with such antisera and/or T-cells at a level that is not substantially less than the reactivity of the full length polypeptide (*e.g.*, in an ELISA and/or T-cell reactivity
20 assay). Such immunogenic portions may react within such assays at a level that is similar to or greater than the reactivity of the full length polypeptide. Such screens may generally be performed using methods well known to those of ordinary skill in the art, such as those described in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. For example, a polypeptide may be immobilized on a
25 solid support and contacted with patient sera to allow binding of antibodies within the sera to the immobilized polypeptide. Unbound sera may then be removed and bound antibodies detected using, for example, ¹²⁵I-labeled Protein A.

As noted above, a composition may comprise a variant of a native Chlamydia protein. A polypeptide "variant," as used herein, is a polypeptide that differs
30 from a native Chlamydia protein in one or more substitutions, deletions, additions and/or insertions, such that the immunogenicity of the polypeptide is not substantially diminished. In other words, the ability of a variant to react with antigen-specific

antisera may be enhanced or unchanged, relative to the native protein, or may be diminished by less than 50%, and preferably less than 20%, relative to the native protein. Such variants may generally be identified by modifying one of the above polypeptide sequences and evaluating the reactivity of the modified polypeptide with
5 antigen-specific antibodies or antisera as described herein. Preferred variants include those in which one or more portions, such as an N-terminal leader sequence or transmembrane domain, have been removed. Other preferred variants include variants in which a small portion (*e.g.*, 1-30 amino acids, preferably 5-15 amino acids) has been removed from the N- and/or C-terminal of the mature protein.

10 Polypeptide variants encompassed by the present invention include those exhibiting at least about 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more identity (determined as described above) to the polypeptides disclosed herein.

Preferably, a variant contains conservative substitutions. A
15 "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydropathic nature of the polypeptide to be substantially unchanged. Amino acid substitutions may generally be made on the basis of similarity in polarity, charge, solubility, hydrophobicity,
20 hydrophilicity and/or the amphipathic nature of the residues. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine and valine; glycine and alanine; asparagine and glutamine; and serine, threonine, phenylalanine and tyrosine.
25 Other groups of amino acids that may represent conservative changes include: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. A variant may also, or alternatively, contain nonconservative changes. In a preferred embodiment, variant polypeptides differ from a native sequence by substitution, deletion or addition of five amino acids or fewer.
30 Variants may also (or alternatively) be modified by, for example, the deletion or addition of amino acids that have minimal influence on the immunogenicity, secondary structure and hydropathic nature of the polypeptide.

As noted above, polypeptides may comprise a signal (or leader) sequence at the N-terminal end of the protein, which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide
5 (e.g., poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

Polypeptides may be prepared using any of a variety of well known techniques. Recombinant polypeptides encoded by DNA sequences as described above may be readily prepared from the DNA sequences using any of a variety of expression
10 vectors known to those of ordinary skill in the art. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast, and higher eukaryotic cells, such as mammalian cells and plant cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell
15 line such as COS or CHO. Supernatants from suitable host/vector systems which secrete recombinant protein or polypeptide into culture media may be first concentrated using a commercially available filter. Following concentration, the concentrate may be applied to a suitable purification matrix such as an affinity matrix or an ion exchange resin. Finally, one or more reverse phase HPLC steps can be employed to further purify
20 a recombinant polypeptide.

Portions and other variants having less than about 100 amino acids, and generally less than about 50 amino acids, may also be generated by synthetic means, using techniques well known to those of ordinary skill in the art. For example, such polypeptides may be synthesized using any of the commercially available solid-phase
25 techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. See Merrifield, *J. Am. Chem. Soc.* 85:2149-2146, 1963. Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Perkin Elmer/Applied BioSystems Division (Foster City, CA), and may be operated according to the manufacturer's
30 instructions.

Within certain specific embodiments, a polypeptide may be a fusion protein that comprises multiple polypeptides as described herein, or that comprises at

least one polypeptide as described herein and an unrelated sequence, such as a known Chlamydia protein. A fusion partner may, for example, assist in providing T helper epitopes (an immunological fusion partner), preferably T helper epitopes recognized by humans, or may assist in expressing the protein (an expression enhancer) at higher yields than the native recombinant protein. Certain preferred fusion partners are both immunological and expression enhancing fusion partners. Other fusion partners may be selected so as to increase the solubility of the protein or to enable the protein to be targeted to desired intracellular compartments. Still further fusion partners include affinity tags, which facilitate purification of the protein.

Fusion proteins may generally be prepared using standard techniques, including chemical conjugation. Preferably, a fusion protein is expressed as a recombinant protein, allowing the production of increased levels, relative to a non-fused protein, in an expression system. Briefly, DNA sequences encoding the polypeptide components may be assembled separately, and ligated into an appropriate expression vector. The 3' end of the DNA sequence encoding one polypeptide component is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide component so that the reading frames of the sequences are in phase. This permits translation into a single fusion protein that retains the biological activity of both component polypeptides.

A peptide linker sequence may be employed to separate the first and second polypeptide components by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S.

Patent No. 4,751,180. The linker sequence may generally be from 1 to about 50 amino acids in length. Linker sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

5 The ligated DNA sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of DNA are located only 5' to the DNA sequence encoding the first polypeptides. Similarly, stop codons required to end translation and transcription termination signals are only present 3' to the DNA sequence encoding the
10 second polypeptide.

 Fusion proteins are also provided. Such proteins comprise a polypeptide as described herein together with an unrelated immunogenic protein. Preferably the immunogenic protein is capable of eliciting a recall response. Examples of such proteins include tetanus, tuberculosis and hepatitis proteins (*see*, for example, Stoute
15 et al. *New Engl. J. Med.*, 336:86-91, 1997).

 Within preferred embodiments, an immunological fusion partner is derived from protein D, a surface protein of the gram-negative bacterium *Haemophilus influenza B* (WO 91/18926). Preferably, a protein D derivative comprises approximately the first third of the protein (*e.g.*, the first N-terminal 100-110 amino
20 acids), and a protein D derivative may be lipidated. Within certain preferred embodiments, the first 109 residues of a Lipoprotein D fusion partner is included on the N-terminus to provide the polypeptide with additional exogenous T-cell epitopes and to increase the expression level in *E. coli* (thus functioning as an expression enhancer). The lipid tail ensures optimal presentation of the antigen to antigen presenting cells.
25 Other fusion partners include the non-structural protein from influenzae virus, NS1 (hemagglutinin). Typically, the N-terminal 81 amino acids are used, although different fragments that include T-helper epitopes may be used.

 In another embodiment, the immunological fusion partner is the protein known as LYTA, or a portion thereof (preferably a C-terminal portion). LYTA is
30 derived from *Streptococcus pneumoniae*, which synthesizes an N-acetyl-L-alanine amidase known as amidase LYTA (encoded by the *LytA* gene; *Gene* 43:265-292, 1986). LYTA is an autolysin that specifically degrades certain bonds in the peptidoglycan

backbone. The C-terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E. coli* C-LYTA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LYTA
5 fragment at the amino terminus has been described (see *Biotechnology* 10:795-798, 1992). Within a preferred embodiment, a repeat portion of LYTA may be incorporated into a fusion protein. A repeat portion is found in the C-terminal region starting at residue 178. A particularly preferred repeat portion incorporates residues 188-305.

In general, polypeptides (including fusion proteins) and polynucleotides
10 as described herein are isolated. An "isolated" polypeptide or polynucleotide is one that is removed from its original environment. For example, a naturally-occurring protein is isolated if it is separated from some or all of the coexisting materials in the natural system. Preferably, such polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. A polynucleotide is
15 considered to be isolated if, for example, it is cloned into a vector that is not a part of the natural environment.

ILLUSTRATIVE THERAPEUTIC COMPOSITIONS AND USES

In another aspect, the present invention provides methods for using one
20 or more of the above polypeptides or fusion proteins (or polynucleotides encoding such polypeptides or fusion proteins) to induce protective immunity against Chlamydial infection in a patient. As used herein, a "patient" refers to any warm-blooded animal, preferably a human. A patient may be afflicted with a disease, or may be free of detectable disease and/or infection. In other words, protective immunity may be
25 induced to prevent or treat Chlamydial infection.

In this aspect, the polypeptide, fusion protein or polynucleotide molecule is generally present within a pharmaceutical composition or a vaccine. Pharmaceutical compositions may comprise one or more polypeptides, each of which may contain one or more of the above sequences (or variants thereof), and a physiologically acceptable
30 carrier. Vaccines may comprise one or more of the above polypeptides and an immunostimulant, such as an adjuvant or a liposome (into which the polypeptide is incorporated). Such pharmaceutical compositions and vaccines may also contain other

Chlamydia antigens, either incorporated into a combination polypeptide or present within a separate polypeptide.

Alternatively, a vaccine may contain polynucleotides encoding one or more polypeptides or fusion proteins as described above, such that the polypeptide is generated *in situ*. In such vaccines, the polynucleotides may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacterial and viral expression systems. Appropriate nucleic acid expression systems contain the necessary polynucleotide sequences for expression in the patient (such as a suitable promoter and terminating signal). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an immunogenic portion of the polypeptide on its cell surface. In a preferred embodiment, the polynucleotides may be introduced using a viral expression system (e.g., vaccinia or other pox virus, retrovirus, or adenovirus), which may involve the use of a non-pathogenic (defective) virus. Techniques for incorporating polynucleotides into such expression systems are well known to those of ordinary skill in the art. The polynucleotides may also be administered as "naked" plasmid vectors as described, for example, in Ulmer et al., *Science* 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. Techniques for incorporating DNA into such vectors are well known to those of ordinary skill in the art. A retroviral vector may additionally transfer or incorporate a gene for a selectable marker (to aid in the identification or selection of transduced cells) and/or a targeting moiety, such as a gene that encodes a ligand for a receptor on a specific target cell, to render the vector target specific. Targeting may also be accomplished using an antibody, by methods known to those of ordinary skill in the art.

Other formulations for therapeutic purposes include colloidal dispersion systems, such as macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. A preferred colloidal system for use as a delivery vehicle *in vitro* and *in vivo* is a liposome (i.e., an artificial membrane vesicle). The uptake of naked polynucleotides may be increased by incorporating the polynucleotides into and/or onto biodegradable beads, which are efficiently transported into the cells. The preparation and use of such systems is well known in the art.

In a related aspect, a polynucleotide vaccine as described above may be administered simultaneously with or sequentially to either a polypeptide of the present invention or a known *Chlamydia* antigen. For example, administration of polynucleotides encoding a polypeptide of the present invention, either "naked" or in a delivery system as described above, may be followed by administration of an antigen in order to enhance the protective immune effect of the vaccine.

Polypeptides and polynucleotides disclosed herein may also be employed in adoptive immunotherapy for the treatment of *Chlamydial* infection. Adoptive immunotherapy may be broadly classified into either active or passive immunotherapy. In active immunotherapy, treatment relies on the *in vivo* stimulation of the endogenous host immune system with the administration of immune response-modifying agents (for example, vaccines, bacterial adjuvants, and/or cytokines).

In passive immunotherapy, treatment involves the delivery of biologic reagents with established immune reactivity (such as effector cells or antibodies) that can directly or indirectly mediate anti-*Chlamydia* effects and does not necessarily depend on an intact host immune system. Examples of effector cells include T lymphocytes (for example, CD8+ cytotoxic T-lymphocyte, CD4+ T-helper), killer cells (such as Natural Killer cells, lymphokine-activated killer cells), B cells, or antigen presenting cells (such as dendritic cells and macrophages) expressing the disclosed antigens. The polypeptides disclosed herein may also be used to generate antibodies or anti-idiotypic antibodies (as in U.S. Patent No. 4,918,164), for passive immunotherapy.

The predominant method of procuring adequate numbers of T-cells for adoptive immunotherapy is to grow immune T-cells *in vitro*. Culture conditions for expanding single antigen-specific T-cells to several billion in number with retention of antigen recognition *in vivo* are well known in the art. These *in vitro* culture conditions typically utilize intermittent stimulation with antigen, often in the presence of cytokines, such as IL-2, and non-dividing feeder cells. As noted above, the immunoreactive polypeptides described herein may be used to rapidly expand antigen-specific T cell cultures in order to generate sufficient number of cells for immunotherapy. In particular, antigen-presenting cells, such as dendritic, macrophage, monocyte, fibroblast, or B-cells, may be pulsed with immunoreactive polypeptides, or polynucleotide sequence(s) may be introduced into antigen presenting cells, using a

variety of standard techniques well known in the art. For example, antigen presenting cells may be transfected or transduced with a polynucleotide sequence, wherein said sequence contains a promoter region appropriate for increasing expression, and can be expressed as part of a recombinant virus or other expression system. Several viral
5 vectors may be used to transduce an antigen presenting cell, including pox virus, vaccinia virus, and adenovirus; also, antigen presenting cells may be transfected with polynucleotide sequences disclosed herein by a variety of means, including gene-gun technology, lipid-mediated delivery, electroporation, osmotic shock, and particulate delivery mechanisms, resulting in efficient and acceptable expression levels as
10 determined by one of ordinary skill in the art. For cultured T-cells to be effective in therapy, the cultured T-cells must be able to grow and distribute widely and to survive long term *in vivo*. Studies have demonstrated that cultured T-cells can be induced to grow *in vivo* and to survive long term in substantial numbers by repeated stimulation with antigen supplemented with IL-2 (see, for example, Cheever, M., *et al*, "Therapy
15 With Cultured T Cells: Principles Revisited," *Immunological Reviews*, 157:177, 1997).

The polypeptides disclosed herein may also be employed to generate and/or isolate chlamydial-reactive T-cells, which can then be administered to the patient. In one technique, antigen-specific T-cell lines may be generated by *in vivo* immunization with short peptides corresponding to immunogenic portions of the
20 disclosed polypeptides. The resulting antigen specific CD8+ or CD4+ T-cell clones may be isolated from the patient, expanded using standard tissue culture techniques, and returned to the patient.

Alternatively, peptides corresponding to immunogenic portions of the polypeptides may be employed to generate *Chlamydia* reactive T cell subsets by
25 selective *in vitro* stimulation and expansion of autologous T cells to provide antigen-specific T cells which may be subsequently transferred to the patient as described, for example, by Chang *et al*, (*Crit. Rev. Oncol. Hematol.*, 22(3), 213, 1996). Cells of the immune system, such as T cells, may be isolated from the peripheral blood of a patient, using a commercially available cell separation system, such as Isolex™ System,
30 available from Nexell Therapeutics, Inc. Irvine, CA. The separated cells are stimulated with one or more of the immunoreactive polypeptides contained within a delivery vehicle, such as a microsphere, to provide antigen-specific T cells. The population of

antigen-specific T cells is then expanded using standard techniques and the cells are administered back to the patient.

In other embodiments, T-cell and/or antibody receptors specific for the polypeptides disclosed herein can be cloned, expanded, and transferred into other vectors or effector cells for use in adoptive immunotherapy. In particular, T cells may be transfected with the appropriate genes to express the variable domains from chlamydia specific monoclonal antibodies as the extracellular recognition elements and joined to the T cell receptor signaling chains, resulting in T cell activation, specific lysis, and cytokine release. This enables the T cell to redirect its specificity in an MHC-independent manner. See for example, Eshhar, Z., *Cancer Immunol Immunother*, 45(3-4):131-6, 1997 and Hwu, P., et al, *Cancer Res*, 55(15):3369-73, 1995. Another embodiment may include the transfection of chlamydia antigen specific alpha and beta T cell receptor chains into alternate T cells, as in Cole, DJ, et al, *Cancer Res*, 55(4):748-52, 1995.

In a further embodiment, syngeneic or autologous dendritic cells may be pulsed with peptides corresponding to at least an immunogenic portion of a polypeptide disclosed herein. The resulting antigen-specific dendritic cells may either be transferred into a patient, or employed to stimulate T cells to provide antigen-specific T cells which may, in turn, be administered to a patient. The use of peptide-pulsed dendritic cells to generate antigen-specific T cells and the subsequent use of such antigen-specific T cells to eradicate disease in a murine model has been demonstrated by Cheever et al, *Immunological Reviews*, 157:177, 1997). Additionally, vectors expressing the disclosed polynucleotides may be introduced into stem cells taken from the patient and clonally propagated *in vitro* for autologous transplant back into the same patient.

Within certain aspects, polypeptides, polynucleotides, T cells and/or binding agents disclosed herein may be incorporated into pharmaceutical compositions or immunogenic compositions (*i.e.*, vaccines). Alternatively, a pharmaceutical composition may comprise an antigen-presenting cell (*e.g.* a dendritic cell) transfected with a *Chlamydial* polynucleotide such that the antigen presenting cell expresses a *Chlamydial* polypeptide. Pharmaceutical compositions comprise one or more such compounds and a physiologically acceptable carrier. Vaccines may comprise one or more such compounds and an immunostimulant. An immunostimulant may be any

substance that enhances or potentiates an immune response to an exogenous antigen. Examples of immunostimulants include adjuvants, biodegradable microspheres (*e.g.*, polylactic galactide) and liposomes (into which the compound is incorporated; *see e.g.*, Fullerton, U.S. Patent No. 4,235,877). Vaccine preparation is generally described in, 5 for example, M.F. Powell and M.J. Newman, eds., "Vaccine Design (the subunit and adjuvant approach)," Plenum Press (NY, 1995). Pharmaceutical compositions and vaccines within the scope of the present invention may also contain other compounds, which may be biologically active or inactive. For example, one or more immunogenic portions of other *Chlamydial* antigens may be present, either incorporated into a fusion 10 polypeptide or as a separate compound, within the composition or vaccine.

A pharmaceutical composition or vaccine may contain DNA encoding one or more of the polypeptides as described above, such that the polypeptide is generated *in situ*. As noted above, the DNA may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid 15 expression systems, bacteria and viral expression systems. Numerous gene delivery techniques are well known in the art, such as those described by Rolland, *Crit. Rev. Therap. Drug Carrier Systems* 15:143-198, 1998, and references cited therein. Appropriate nucleic acid expression systems contain the necessary DNA sequences for expression in the patient (such as a suitable promoter and terminating signal). Bacterial 20 delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an immunogenic portion of the polypeptide on its cell surface or secretes such an epitope.

In a preferred embodiment, the DNA may be introduced using a viral expression system (*e.g.*, vaccinia or other pox virus, retrovirus, adenovirus, baculovirus, 25 togavirus, bacteriophage, and the like), which often involves the use of a non-pathogenic (defective), replication competent virus.

For example, many viral expression vectors are derived from viruses of the retroviridae family. This family includes the murine leukemia viruses, the mouse mammary tumor viruses, the human foamy viruses, Rous sarcoma virus, and the 30 immunodeficiency viruses, including human, simian, and feline. Considerations when designing retroviral expression vectors are discussed in Comstock *et al.* (1997).

Excellent murine leukemia virus (MLV)-based viral expression vectors have been developed by Kim *et al.* (1998). In creating the MLV vectors, Kim *et al.* found that the entire *gag* sequence, together with the immediate upstream region, could be deleted without significantly affecting viral packaging or gene expression. Further, it was found that nearly the entire U3 region could be replaced with the immediately-early promoter of human cytomegalovirus without deleterious effects. Additionally, MCR and internal ribosome entry sites (IRES) could be added without adverse effects. Based on their observations, Kim *et al.* have designed a series of MLV-based expression vectors comprising one or more of the features described above.

As more has been learned about human foamy virus (HFV), characteristics of HFV that are favorable for its use as an expression vector have been discovered. These characteristics include the expression of pol by splicing and start of translation at a defined initiation codon. Other aspects of HFV viral expression vectors are reviewed in Bodem *et al.* (1997).

Murakami *et al.* (1997) describe a Rous sarcoma virus (RSV)-based replication-competent avian retrovirus vectors, IR1 and IR2 to express a heterologous gene at a high level. In these vectors, the IRES derived from encephalomyocarditis virus (EMCV) was inserted between the *env* gene and the heterologous gene. The IR1 vector retains the splice-acceptor site that is present downstream of the *env* gene while the IR2 vector lacks it. Murakami *et al.* have shown high level expression of several different heterologous genes by these vectors.

Recently, a number of lentivirus-based retroviral expression vectors have been developed. Kafri *et al.* (1997) have shown sustained expression of genes delivered directly into liver and muscle by a human immunodeficiency virus (HIV)-based expression vector. One benefit of the system is the inherent ability of HIV to transduce non-dividing cells. Because the viruses of Kafri *et al.* are pseudotyped with vesicular stomatitis virus G glycoprotein (VSVG), they can transduce a broad range of tissues and cell types.

A large number of adenovirus-based expression vectors have been developed, primarily due to the advantages offered by these vectors in gene therapy applications. Adenovirus expression vectors and methods of using such vectors are the subject of a number of United States patents, including United States Patent No.

5,698,202, United States Patent No. 5,616,326, United States Patent No. 5,585,362, and United States Patent No. 5,518,913, all incorporated herein by reference.

Additional adenoviral constructs are described in Khatri *et al.* (1997) and Tomanin *et al.* (1997). Khatri *et al.* describe novel ovine adenovirus expression vectors and their ability to infect bovine nasal turbinate and rabbit kidney cells as well as a range of human cell type, including lung and foreskin fibroblasts as well as liver, prostate, breast, colon and retinal lines. Tomanin *et al.* describe adenoviral expression vectors containing the T7 RNA polymerase gene. When introduced into cells containing a heterologous gene operably linked to a T7 promoter, the vectors were able to drive gene expression from the T7 promoter. The authors suggest that this system may be useful for the cloning and expression of genes encoding cytotoxic proteins.

Poxviruses are widely used for the expression of heterologous genes in mammalian cells. Over the years, the vectors have been improved to allow high expression of the heterologous gene and simplify the integration of multiple heterologous genes into a single molecule. In an effort to diminish cytopathic effects and to increase safety, vaccinia virus mutant and other poxviruses that undergo abortive infection in mammalian cells are receiving special attention (Oertli *et al.*, 1997). The use of poxviruses as expression vectors is reviewed in Carroll and Moss (1997).

Togaviral expression vectors, which includes alphaviral expression vectors have been used to study the structure and function of proteins and for protein production purposes. Attractive features of togaviral expression vectors are rapid and efficient gene expression, wide host range, and RNA genomes (Huang, 1996). Also, recombinant vaccines based on alphaviral expression vectors have been shown to induce a strong humoral and cellular immune response with good immunological memory and protective effects (Tubulekas *et al.*, 1997). Alphaviral expression vectors and their use are discussed, for example, in Lundstrom (1997).

In one study, Li and Garoff (1996) used Semliki Forest virus (SFV) expression vectors to express retroviral genes and to produce retroviral particles in BHK-21 cells. The particles produced by this method had protease and reverse transcriptase activity and were infectious. Furthermore, no helper virus could be detected in the virus stocks. Therefore, this system has features that are attractive for its use in gene therapy protocols.

Baculoviral expression vectors have traditionally been used to express heterologous proteins in insect cells. Examples of proteins include mammalian chemokine receptors (Wang *et al.*, 1997), reporter proteins such as green fluorescent protein (Wu *et al.*, 1997), and FLAG fusion proteins (Wu *et al.*, 1997; Koh *et al.*, 1997).

5 Recent advances in baculoviral expression vector technology, including their use in virion display vectors and expression in mammalian cells is reviewed by Possee (1997). Other reviews on baculoviral expression vectors include Jones and Morikawa (1996) and O'Reilly (1997).

Other suitable viral expression systems are disclosed, for example, in

10 Fisher-Hoeh et al., *Proc. Natl. Acad. Sci. USA* 86:317-321, 1989; Flexner et al., *Ann. N.Y. Acad. Sci.* 569:86-103, 1989; Flexner et al., *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfeld et al., *Science* 252:431-434, 1991; Kolls et al., *Proc. Natl. Acad. Sci. USA*

15 91:215-219, 1994; Kass-Eisler et al., *Proc. Natl. Acad. Sci. USA* 90:11498-11502, 1993; Guzman et al., *Circulation* 88:2838-2848, 1993; and Guzman et al., *Cir. Res.* 73:1202-1207, 1993. Techniques for incorporating DNA into such expression systems are well known to those of ordinary skill in the art. In other systems, the DNA may be introduced as "naked" DNA, as described, for example, in Ulmer et al., *Science*

20 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

It will be apparent that a vaccine may comprise a polynucleotide and/or a polypeptide component, as desired. It will also be apparent that a vaccine may contain

25 pharmaceutically acceptable salts of the polynucleotides and/or polypeptides provided herein. Such salts may be prepared from pharmaceutically acceptable non-toxic bases, including organic bases (*e.g.*, salts of primary, secondary and tertiary amines and basic amino acids) and inorganic bases (*e.g.*, sodium, potassium, lithium, ammonium, calcium and magnesium salts). While any suitable carrier known to those of ordinary

30 skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. Compositions of the present invention may be formulated for any appropriate manner of administration,

including for example, topical, oral, nasal, intravenous, intracranial, intraperitoneal, subcutaneous or intramuscular administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, 5 such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (e.g., polylactate polyglycolate) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109.

10 Such compositions may also comprise buffers (e.g., neutral buffered saline or phosphate buffered saline), carbohydrates (e.g., glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, bacteriostats, chelating agents such as EDTA or glutathione, adjuvants (e.g., aluminum hydroxide), solutes that render the formulation isotonic, hypotonic or weakly hypertonic 15 with the blood of a recipient, suspending agents, thickening agents and/or preservatives. Alternatively, compositions of the present invention may be formulated as a lyophilizate. Compounds may also be encapsulated within liposomes using well known technology.

Any of a variety of immunostimulants may be employed in the vaccines 20 of this invention. For example, an adjuvant may be included. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a stimulator of immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis* derived proteins. Suitable adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant 25 and Complete Adjuvant (Difco Laboratories, Detroit, MI); Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ); AS-2 (SmithKline Beecham, Philadelphia, PA); aluminum salts such as aluminum hydroxide gel (alum) or aluminum phosphate; salts of calcium, iron or zinc; an insoluble suspension of acylated tyrosine; acylated sugars; cationically or anionically derivatized polysaccharides; polyphosphazenes; 30 biodegradable microspheres; monophosphoryl lipid A and quil A. Cytokines, such as GM-CSF or interleukin-2, -7, or -12, may also be used as adjuvants.

Within the vaccines provided herein, under select circumstances, the

adjuvant composition may be designed to induce an immune response predominantly of the Th1 type or Th2 type. High levels of Th1-type cytokines (e.g., IFN- γ , TNF α , IL-2 and IL-12) tend to favor the induction of cell mediated immune responses to an administered antigen. In contrast, high levels of Th2-type cytokines (e.g., IL-4, IL-5, IL-6 and IL-10) tend to favor the induction of humoral immune responses. Following application of a vaccine as provided herein, a patient will support an immune response that includes Th1- and Th2-type responses. Within a preferred embodiment, in which a response is predominantly Th1-type, the level of Th1-type cytokines will increase to a greater extent than the level of Th2-type cytokines. The levels of these cytokines may be readily assessed using standard assays. For a review of the families of cytokines, see Mosmann and Coffman, *Ann. Rev. Immunol.* 7:145-173, 1989.

Preferred adjuvants for use in eliciting a predominantly Th1-type response include, for example, a combination of monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A (3D-MPL), together with an aluminum salt. MPL adjuvants are available from Corixa Corporation (Seattle, WA; see US Patent Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094). CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) also induce a predominantly Th1 response. Such oligonucleotides are well known and are described, for example, in WO 96/02555 and WO 99/33488. Immunostimulatory DNA sequences are also described, for example, by Sato et al., *Science* 273:352, 1996. Another preferred adjuvant is a saponin, preferably QS21 (Aquila Biopharmaceuticals Inc., Framingham, MA), which may be used alone or in combination with other adjuvants. For example, an enhanced system involves the combination of a monophosphoryl lipid A and saponin derivative, such as the combination of QS21 and 3D-MPL as described in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol, as described in WO 96/33739. Other preferred formulations comprise an oil-in-water emulsion and tocopherol. A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil-in-water emulsion is described in WO 95/17210.

Other preferred adjuvants include Montanide ISA 720 (Seppic, France), SAF (Chiron, California, United States), ISCOMS (CSL), MF-59 (Chiron), the SBAS series of adjuvants (e.g., SBAS-2 or SBAS-4, available from SmithKline Beecham, Rixensart, Belgium), Detox (Corixa Corporation; Seattle, WA), RC-529 (Corixa

Corporation; Seattle, WA) and other aminoalkyl glucosaminide 4-phosphates (AGPs), such as those described in pending U.S. Patent Application Serial Nos. 08/853,826 and 09/074,720, the disclosures of which are incorporated herein by reference in their entireties.

5 Any vaccine provided herein may be prepared using well known methods that result in a combination of antigen, immunostimulant and a suitable carrier or excipient. The compositions described herein may be administered as part of a sustained release formulation (*i.e.*, a formulation such as a capsule, sponge or gel (composed of polysaccharides, for example) that effects a slow release of compound
10 following administration). Such formulations may generally be prepared using well known technology (*see, e.g.*, Coombes et al., *Vaccine* 14:1429-1438, 1996) and administered by, for example, oral, rectal or subcutaneous implantation, or by implantation at the desired target site. Sustained-release formulations may contain a polypeptide, polynucleotide or antibody dispersed in a carrier matrix and/or contained
15 within a reservoir surrounded by a rate controlling membrane.

Carriers for use within such formulations are biocompatible, and may also be biodegradable; preferably the formulation provides a relatively constant level of active component release. Such carriers include microparticles of poly(lactide-co-glycolide), as well as polyacrylate, latex, starch, cellulose and dextran. Other delayed-
20 release carriers include supramolecular biovectors, which comprise a non-liquid hydrophilic core (*e.g.*, a cross-linked polysaccharide or oligosaccharide) and, optionally, an external layer comprising an amphiphilic compound, such as a phospholipid (*see e.g.*, U.S. Patent No. 5,151,254 and PCT applications WO 94/20078, WO/94/23701 and WO 96/06638). The amount of active compound contained within a sustained release
25 formulation depends upon the site of implantation, the rate and expected duration of release and the nature of the condition to be treated or prevented.

Any of a variety of delivery vehicles may be employed within pharmaceutical compositions and vaccines to facilitate production of an antigen-specific immune response that targets *Chlamydia*-infected cells. Delivery vehicles include
30 antigen presenting cells (APCs), such as dendritic cells, macrophages, B cells, monocytes and other cells that may be engineered to be efficient APCs. Such cells may, but need not, be genetically modified to increase the capacity for presenting the antigen,

to improve activation and/or maintenance of the T cell response, to have anti-*Chlamydia* effects *per se* and/or to be immunologically compatible with the receiver (i.e., matched HLA haplotype). APCs may generally be isolated from any of a variety of biological fluids and organs, and may be autologous, allogeneic, syngeneic or
5 xenogeneic cells.

Certain preferred embodiments of the present invention use dendritic cells or progenitors thereof as antigen-presenting cells. Dendritic cells are highly potent APCs (Banchereau and Steinman, *Nature* 392:245-251, 1998) and have been shown to be effective as a physiological adjuvant for eliciting prophylactic or therapeutic
10 immunity (see Timmerman and Levy, *Ann. Rev. Med.* 50:507-529, 1999). In general, dendritic cells may be identified based on their typical shape (stellate *in situ*, with marked cytoplasmic processes (dendrites) visible *in vitro*), their ability to take up, process and present antigens with high efficiency, and their ability to activate naïve T cell responses. Dendritic cells may, of course, be engineered to express specific cell-
15 surface receptors or ligands that are not commonly found on dendritic cells *in vivo* or *ex vivo*, and such modified dendritic cells are contemplated by the present invention. As an alternative to dendritic cells, secreted vesicles antigen-loaded dendritic cells (called exosomes) may be used within a vaccine (see Zitvogel et al., *Nature Med.* 4:594-600, 1998).

20 Dendritic cells and progenitors may be obtained from peripheral blood, bone marrow, lymph nodes, spleen, skin, umbilical cord blood or any other suitable tissue or fluid. For example, dendritic cells may be differentiated *ex vivo* by adding a combination of cytokines such as GM-CSF, IL-4, IL-13 and/or TNF α to cultures of monocytes harvested from peripheral blood. Alternatively, CD34 positive cells
25 harvested from peripheral blood, umbilical cord blood or bone marrow may be differentiated into dendritic cells by adding to the culture medium combinations of GM-CSF, IL-3, TNF α , CD40 ligand, LPS, flt3 ligand and/or other compound(s) that induce differentiation, maturation and proliferation of dendritic cells.

Dendritic cells are conveniently categorized as "immature" and "mature"
30 cells, which allows a simple way to discriminate between two well characterized phenotypes. However, this nomenclature should not be construed to exclude all possible intermediate stages of differentiation. Immature dendritic cells are

characterized as APC with a high capacity for antigen uptake and processing, which correlates with the high expression of Fc γ receptor and mannose receptor. The mature phenotype is typically characterized by a lower expression of these markers, but a high expression of cell surface molecules responsible for T cell activation such as class I and
5 class II MHC, adhesion molecules (*e.g.*, CD54 and CD11) and costimulatory molecules (*e.g.*, CD40, CD80, CD86 and 4-1BB).

APCs may generally be transfected with a polynucleotide encoding a *Chlamydial* protein (or portion or other variant thereof) such that the *Chlamydial* polypeptide, or an immunogenic portion thereof, is expressed on the cell surface. Such
10 transfection may take place *ex vivo*, and a composition or vaccine comprising such transfected cells may then be used for therapeutic purposes, as described herein. Alternatively, a gene delivery vehicle that targets a dendritic or other antigen presenting cell may be administered to a patient, resulting in transfection that occurs *in vivo*. *In vivo* and *ex vivo* transfection of dendritic cells, for example, may generally be performed
15 using any methods known in the art, such as those described in WO 97/24447, or the gene gun approach described by Mahvi et al., *Immunology and cell Biology* 75:456-460, 1997. Antigen loading of dendritic cells may be achieved by incubating dendritic cells or progenitor cells with the *Chlamydial* polypeptide, DNA (naked or within a plasmid vector) or RNA; or with antigen-expressing recombinant bacterium or viruses (*e.g.*,
20 vaccinia, fowlpox, adenovirus or lentivirus vectors). Prior to loading, the polypeptide may be covalently conjugated to an immunological partner that provides T cell help (*e.g.*, a carrier molecule). Alternatively, a dendritic cell may be pulsed with a non-conjugated immunological partner, separately or in the presence of the polypeptide.

Routes and frequency of administration of pharmaceutical compositions
25 and vaccines, as well as dosage, will vary from individual to individual. In general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g.*, intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (*e.g.*, by aspiration) or orally. Between 1 and 3 doses may be administered for a 1-36 week period. Preferably, 3 doses are administered, at intervals of 3-4 months, and booster
30 vaccinations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of polypeptide or DNA that, when administered as described above, is capable of raising an immune

response in an immunized patient sufficient to protect the patient from *Chlamydial* infection for at least 1-2 years. In general, the amount of polypeptide present in a dose (or produced *in situ* by the DNA in a dose) ranges from about 1 pg to about 100 mg per kg of host, typically from about 10 pg to about 1 mg, and preferably from about 100 pg to about 1 µg. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.1 mL to about 5 mL.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (e.g., polylactic galactide) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109.

In general, an appropriate dosage and treatment regimen provides the active compound(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit. Such a response can be monitored by establishing an improved clinical outcome in treated patients as compared to non-treated patients. Increases in preexisting immune responses to a *Chlamydial* protein generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using standard proliferation, cytotoxicity or cytokine assays, which may be performed using samples obtained from a patient before and after treatment.

DETECTION AND DIAGNOSIS

In another aspect, the present invention provides methods for using the polypeptides described above to diagnose *Chlamydial* infection. In this aspect, methods are provided for detecting *Chlamydial* infection in a biological sample, using one or more of the above polypeptides, either alone or in combination. For clarity, the term "polypeptide" will be used when describing specific embodiments of the inventive

diagnostic methods. However, it will be clear to one of skill in the art that the fusion proteins of the present invention may also be employed in such methods.

As used herein, a "biological sample" is any antibody-containing sample obtained from a patient. Preferably, the sample is whole blood, sputum, serum, plasma, saliva, cerebrospinal fluid or urine. More preferably, the sample is a blood, serum or plasma sample obtained from a patient. The polypeptides are used in an assay, as described below, to determine the presence or absence of antibodies to the polypeptide(s) in the sample, relative to a predetermined cut-off value. The presence of such antibodies indicates previous sensitization to *Chlamydia* antigens which may be indicative of *Chlamydia*-infection.

In embodiments in which more than one polypeptide is employed, the polypeptides used are preferably complementary (*i.e.*, one component polypeptide will tend to detect infection in samples where the infection would not be detected by another component polypeptide). Complementary polypeptides may generally be identified by using each polypeptide individually to evaluate serum samples obtained from a series of patients known to be infected with *Chlamydia*. After determining which samples test positive (as described below) with each polypeptide, combinations of two or more polypeptides may be formulated that are capable of detecting infection in most, or all, of the samples tested.

A variety of assay formats are known to those of ordinary skill in the art for using one or more polypeptides to detect antibodies in a sample. *See, e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988, which is incorporated herein by reference. In a preferred embodiment, the assay involves the use of polypeptide immobilized on a solid support to bind to and remove the antibody from the sample. The bound antibody may then be detected using a detection reagent that contains a reporter group. Suitable detection reagents include antibodies that bind to the antibody/polypeptide complex and free polypeptide labeled with a reporter group (*e.g.*, in a semi-competitive assay). Alternatively, a competitive assay may be utilized, in which an antibody that binds to the polypeptide is labeled with a reporter group and allowed to bind to the immobilized antigen after incubation of the antigen with the sample. The extent to which components of the sample inhibit the

binding of the labeled antibody to the polypeptide is indicative of the reactivity of the sample with the immobilized polypeptide.

The solid support may be any solid material known to those of ordinary skill in the art to which the antigen may be attached. For example, the solid support may be a test well in a microtiter plate, or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681.

The polypeptides may be bound to the solid support using a variety of techniques known to those of ordinary skill in the art. In the context of the present invention, the term "bound" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the antigen and functional groups on the support or may be a linkage by way of a cross-linking agent). Binding by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the polypeptide, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of polypeptide ranging from about 10 ng to about 1 μ g, and preferably about 100 ng, is sufficient to bind an adequate amount of antigen.

Covalent attachment of polypeptide to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the polypeptide. For example, the polypeptide may be bound to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the polypeptide (*see, e.g.,* Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is an enzyme linked immunosorbent assay (ELISA). This assay may be performed by first contacting a polypeptide antigen that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that antibodies to the polypeptide within the sample are allowed

to bind to the immobilized polypeptide. Unbound sample is then removed from the immobilized polypeptide and a detection reagent capable of binding to the immobilized antibody-polypeptide complex is added. The amount of detection reagent that remains bound to the solid support is then determined using a method appropriate for the
5 specific detection reagent.

More specifically, once the polypeptide is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin (BSA) or Tween 20™ (Sigma Chemical Co., St. Louis, MO)
10 may be employed. The immobilized polypeptide is then incubated with the sample, and antibody is allowed to bind to the antigen. The sample may be diluted with a suitable diluent, such as phosphate-buffered saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is that period of time that is sufficient to detect the presence of antibody within an HGE-infected sample. Preferably, the contact
15 time is sufficient to achieve a level of binding that is at least 95% of that achieved at equilibrium between bound and unbound antibody. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. Detection reagent may then be added to the solid support. An appropriate detection reagent is any compound that binds to the immobilized antibody-polypeptide complex and that can be detected by any of a variety of means known to those in the art. Preferably, the
25 detection reagent contains a binding agent (such as, for example, Protein A, Protein G, immunoglobulin, lectin or free antigen) conjugated to a reporter group. Preferred reporter groups include enzymes (such as horseradish peroxidase), substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and biotin. The conjugation of binding agent to reporter group may be achieved using standard methods
30 known to those of ordinary skill in the art. Common binding agents may also be purchased conjugated to a variety of reporter groups from many commercial sources (*e.g.*, Zymed Laboratories, San Francisco, CA, and Pierce, Rockford, IL).

The detection reagent is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound antibody. An appropriate amount of time may generally be determined from the manufacturer's instructions or by assaying the level of binding that occurs over a period of time.

5 Unbound detection reagent is then removed and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin

10 may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

To determine the presence or absence of anti-*Chlamydia* antibodies in

15 the sample, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value is the average mean signal obtained when the immobilized antigen is incubated with samples from an uninfected patient. In general, a sample generating a signal that is three standard deviations above

20 the predetermined cut-off value is considered positive for *Chlamydia*-infection. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, pp. 106-107. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true

25 positive rates (i.e., sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (i.e., the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered

30 positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In

general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for Chlamydial infection.

In a related embodiment, the assay is performed in a rapid flow-through or strip test format, wherein the antigen is immobilized on a membrane, such as nitrocellulose. In the flow-through test, antibodies within the sample bind to the immobilized polypeptide as the sample passes through the membrane. A detection reagent (e.g., protein A-colloidal gold) then binds to the antibody-polypeptide complex as the solution containing the detection reagent flows through the membrane. The detection of bound detection reagent may then be performed as described above. In the strip test format, one end of the membrane to which polypeptide is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing detection reagent and to the area of immobilized polypeptide. Concentration of detection reagent at the polypeptide indicates the presence of anti-*Chlamydia* antibodies in the sample. Typically, the concentration of detection reagent at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of polypeptide immobilized on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of antibodies that would be sufficient to generate a positive signal in an ELISA, as discussed above. Preferably, the amount of polypeptide immobilized on the membrane ranges from about 25 ng to about 1 µg, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount (e.g., one drop) of patient serum or blood.

Of course, numerous other assay protocols exist that are suitable for use with the polypeptides of the present invention. The above descriptions are intended to be exemplary only. One example of an alternative assay protocol which may be usefully employed in such methods is a Western blot, wherein the proteins present in a biological sample are separated on a gel, prior to exposure to a binding agent. Such techniques are well known to those of skill in the art.

30 BINDING AGENTS AND THEIR USES

The present invention further provides agents, such as antibodies and antigen-binding fragments thereof, that specifically bind to a *Chlamydial* protein. As

used herein, an antibody, or antigen-binding fragment thereof, is said to "specifically bind" to a *Chlamydial* protein if it reacts at a detectable level (within, for example, an ELISA) with a *Chlamydial* protein, and does not react detectably with unrelated proteins under similar conditions. As used herein, "binding" refers to a noncovalent association between two separate molecules such that a complex is formed. The ability to bind may be evaluated by, for example, determining a binding constant for the formation of the complex. The binding constant is the value obtained when the concentration of the complex is divided by the product of the component concentrations. In general, two compounds are said to "bind," in the context of the present invention, when the binding constant for complex formation exceeds about 10^3 L/mol. The binding constant may be determined using methods well known in the art.

Binding agents may be further capable of differentiating between patients with and without a *Chlamydial* infection using the representative assays provided herein. In other words, antibodies or other binding agents that bind to a *Chlamydial* protein will generate a signal indicating the presence of a *Chlamydial* infection in at least about 20% of patients with the disease, and will generate a negative signal indicating the absence of the disease in at least about 90% of individuals without infection. To determine whether a binding agent satisfies this requirement, biological samples (e.g., blood, sera, sputum urine and/or tissue biopsies) from patients with and without *Chlamydial* infection (as determined using standard clinical tests) may be assayed as described herein for the presence of polypeptides that bind to the binding agent. It will be apparent that a statistically significant number of samples with and without the disease should be assayed. Each binding agent should satisfy the above criteria; however, those of ordinary skill in the art will recognize that binding agents may be used in combination to improve sensitivity.

Any agent that satisfies the above requirements may be a binding agent. For example, a binding agent may be a ribosome, with or without a peptide component, an RNA molecule or a polypeptide. In a preferred embodiment, a binding agent is an antibody or an antigen-binding fragment thereof. Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, antibodies can be produced by cell culture techniques, including the generation

of monoclonal antibodies as described herein, or via transfection of antibody genes into suitable bacterial or mammalian cell hosts, in order to allow for the production of recombinant antibodies. In one technique, an immunogen comprising the polypeptide is initially injected into any of a wide variety of mammals (*e.g.*, mice, rats, rabbits, sheep or goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

Monoclonal antibodies specific for an antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (*i.e.*, reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and their culture supernatants tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable

vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process
5 in, for example, an affinity chromatography step.

Within certain embodiments, the use of antigen-binding fragments of antibodies may be preferred. Such fragments include Fab fragments, which may be prepared using standard techniques. Briefly, immunoglobulins may be purified from rabbit serum by affinity chromatography on Protein A bead columns (Harlow and Lane,
10 *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988) and digested by papain to yield Fab and Fc fragments. The Fab and Fc fragments may be separated by affinity chromatography on protein A bead columns.

Monoclonal antibodies of the present invention may be coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides,
15 differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include ^{90}Y , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{211}At , and ^{212}Bi . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, *Shigella* toxin, and pokeweed
20 antiviral protein.

A therapeutic agent may be coupled (*e.g.*, covalently bonded) to a suitable monoclonal antibody either directly or indirectly (*e.g.*, via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such
25 as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an
30 antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical

reactivity may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spittler), by irradiation of a photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers which provide multiple sites for attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as albumins (*e.g.*, U.S. Patent No. 4,507,234, to Kato et al.), peptides and polysaccharides such as aminodextran (*e.g.*, U.S. Patent No. 4,699,784, to Shih et al.). A carrier may also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating

compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For
5 example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

A variety of routes of administration for the antibodies and immunoconjugates may be used. Typically, administration will be intravenous, intramuscular, subcutaneous or in site-specific regions by appropriate methods. It will
10 be evident that the precise dose of the antibody/immunoconjugate will vary depending upon the antibody used, the antigen density, and the rate of clearance of the antibody.

Antibodies may be used in diagnostic tests to detect the presence of *Chlamydia* antigens using assays similar to those detailed above and other techniques well known to those of skill in the art, thereby providing a method for detecting
15 Chlamydial infection in a patient.

Diagnostic reagents of the present invention may also comprise DNA sequences encoding one or more of the above polypeptides, or one or more portions thereof. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify *Chlamydia*-specific cDNA
20 derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for a DNA molecule encoding a polypeptide of the present invention. The presence of the amplified cDNA is then detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes specific for a DNA molecule encoding a polypeptide of the present invention may be used in a
25 hybridization assay to detect the presence of an inventive polypeptide in a biological sample.

The following Examples are offered by way of illustration and not by way of limitation.

EXAMPLE 1

CD4 T CELL EXPRESSION CLONING FOR THE IDENTIFICATION
OF T CELL STIMULATING ANTIGENS FROM CHLAMYDIA TRACHOMATIS
SEROVAR E

5

In this example, a CD4+ T cell expression cloning strategy was used to identify Chlamydia trachomatis antigens recognized by patients enrolled in Corixa Corporation's blood donor program. A genomic library of Chlamydia trachomatis serovar E was constructed and screened with Chlamydia specific T cell lines generated by stimulating PBMCs from these donors. Donor CT1 is a 27 yr. old male whose clinical manifestation was non-gonococcal urethritis and his urine was tested positive for Chlamydia by ligase chain reaction. Donor CT3 is a 43 yr. old male who is asymptomatic and infected with serovar J. Donor CT10 is a 24 yr. old female who is asymptomatic and was exposed to Chlamydia through her partner but did not develop the disease. Donor CT11 is a 24 yr. old female with multiple infections (serovar J, F and E).

Chlamydia specific T-cell lines were generated from donors with chlamydial genital tract infection or donors exposed to chlamydia who did not develop the disease. T cell lines from donor CT-1, CT-3 and CT-10 were generated by stimulating PBMC's with reticulate bodies of C. trachomatis serovar E. T-cell lines from donor CT-11 were generated by stimulating PBMC's with either reticulate bodies or elementary bodies of C. trachomatis serovar E. A randomly sheared genomic library of C. trachomatis serovar E was constructed in lambda Zap II vector and an amplified library plated out in 96 well microtiter plates at a density of 25 clones/well. Bacteria were induced to express the recombinant protein in the presence of 2mM IPTG for 2hr, then pelleted and resuspended in 200ul RPMI/10% FBS. 10 ul of the induced bacterial suspension was transferred to 96 well plates containing autologous monocyte-derived dendritic cells. After a 2 hour incubation, dendritic cells were washed to remove E. coli and the T cells were added. Positive E. coli pools were identified by determining IFN gamma production and proliferation of T cells in the pools. The number of pools identified by each T-cell line is as follows: CT1 line : 30/480 pools; CT3 line : 91/960

pools; CT10 line : 40/480 pools; CT11 line : 51/480 pools. The clones identified using this approach are set forth in SEQ ID NO:1-14.

In another example using substantially the same approach described above, we identified 12 additional T-cell reactive clones from *Chlamydia trachomatis* serovar E expression screening. Clone E5-E9-3 (CT1 positive) contains a 636 bp insert that encodes partially the ORF for dnaK like gene. Part of this sequence was also identified in clone E1-A5-53. Clone E4-H3-56 (CT1 positive, 463 bp insert) contains a partial ORF for the TSA gene (CT603) on the complementary strand. The insert for clone E2-G12-52 (1265 bp) was identified with the CT11 line. It contains a partial ORF for clpB, a protease ATPase. Another clone identified with the CT11 line, E1-F9-79 (167 bp), contains a partial ORF for the gene CT133 on the complementary strand. CT133 is a predicted rRNA methylase. Clone E4-D2-79 (CT3 positive) contains a 1181 bp insert that is a partial ORF for nrdA gene. The ORF for this gene was also identified in clone E2-B10-52 (CT10 positive). Clone E6-C8-95 contains a 731 bp insert that was identified using the donor lines CT3, CT1, and CT12. This insert has a carboxy terminal half for the gene for the 60 kDa ORF. Clone E7-H11-61 (CT3 positive-1135 bp) has partial inserts for flhA (CT061), tyrS (CT062), TSA (CT603) and a hypothetical protein (CT602). The insert for clone E5-A11-8 (CT10 positive-1736 bp) contains the complete ORF for groES (CT111) and a majority of the ORF for groEL (CT110). Clone E3-F2-37 (CT10, CT3, CT11, and CT12 positive-1377bp insert) contains a partial ORF for gene tRNA-Trp (CT322) and a complete ORF for the gene secE (CT321). E4-G9-75 is another CT10 clone that contains a partial ORF (723 bp insert) for the amino terminal region of the pmpH gene (CT872). Clone E2-D5-89 (516bp) is also a CT10 positive clone that contains a partial ORF for pmpD gene (12). The insert for clone E5-E2-10 (CT10 positive) is 427 bp and contains a partial ORF for the major outer membrane protein omp1.

Example 2

ADDITIONAL CD4 T CELL EXPRESSION CLONING FOR THE IDENTIFICATION OF T CELL STIMULATING ANTIGENS FROM CHLAMYDIA TRACHOMATIS SEROVAR E

Twenty sequences were isolated from single clones using a Chlamydia trachomatis serovar E (Ct E) library expression screening method. Descriptions of how the clones and lines were generated are provided in Example 1.

Clone E5-A8-85 (identified using the CT1 patient line) was found to
5 contain a 1433 bp insert. This insert contains a large region of the C-terminal half of the CT875, a Chlamydia trachomatis hypothetical specific gene that is disclosed in SEQ ID NO:34. Also present in the clone is a partial open reading frame (ORF) of a hypothetical protein CT001 which is on the complementary strand.

The clone E9-G2-93 (identified using the C10 patient line) was shown to
10 contain a 554 bp insert, the sequence of which is disclosed in SEQ ID NO:33. This sequence encodes a partial ORF for CT178, a hypothetical CT protein.

Clone E7-B1-16 (identified using the patient lines CT10, CT3, CT5, CT11, CT13, and CHH037) has a 2577 bp insert, the sequence of which is disclosed in SEQ ID NO:32. This clone was found to contain three ORFs. The first ORF contains
15 almost the entire ORF for CT694, a Chlamydia trachomatis (CT) specific hypothetical protein. The second ORF is a full length ORF for CT695, another hypothetical CT protein. The third ORF is the N-terminal portion of CT696.

Clone E9-D5-8 (identified using the patient lines CT10, CT1, CT4, and CT11) contains a 393 bp insert, which is disclosed in SEQ ID NO:31. It was found to
20 encode a partial ORF for CT680, the S2 ribosomal protein.

Clone E9-E10-51 (identified using the patient line CT10) contains an 883 bp insert, the sequence of which is disclosed in SEQ ID NO:30. This clone contains two partial ORF. The first of these is for the C-terminal half of CT680, which may show some overlap with the insert present in clone E9-D5-8. The second ORF is
25 the N-terminal partial ORF for CT679, which is the elongation factor TS.

Clone E3-B4-18 (identified using the CT1 patient line) contains a 1224 bp insert, the sequence of which is disclosed in SEQ ID NO:29. This clone contains 4 ORFs. At the N-terminal end of the clone is the complete ORF for CT772, coding for inorganic pyrophosphatase. The second ORF is a small portion of the C-terminal end of
30 CT771, on the complementary frame. The third is a partial ORF of the hypothetical protein, CT191 and the fourth is a partial ORF for CT190, DNA gyrase-B.

Clone E10-B2-57 (identified using the CT10 patient line) contains an 822 bp insert, the sequence of which is disclosed in SEQ ID NO:42. This clone contains the complete ORF for CT066, a hypothetical protein, on the complementary strand.

5 Clone E3-F3-18 (identified using the CT1 patient line) contains an 1141 bp insert, the sequence of which is disclosed in SEQ ID NO:41. It contains a partial ORF for pmpG (CT871) in frame with the β -gal gene.

 Clone E4-D6-21 (identified using the CT3 patient line) contains a 1297 bp insert, the sequence of which is disclosed in SEQ ID NO:40. This clone contains a
10 very small portion of xseA (CT329), the entire ORF for tpiS (CT328) on the complementary strand, and a partial amino terminal ORF for trpC (CT327) on the top frame.

 Clone E1-G9-23 (identified using the CT3 patient line) contains an 1180 bp insert, the sequence of which is disclosed in SEQ ID NO:39. This clone contains
15 almost the entire ORF for glycogen synthase (CT798).

 Clone E3-A3-31 (identified using the CT1 patient line) contains an 1834 bp insert, the sequence of which is disclosed in SEQ ID NO:38. This clone contains a large region of the hypothetical gene CT622.

 Clone E2-F7-11 (identified using both the CT3 and CT10 patient lines)
20 contains a 2093 bp insert, the sequence of which is disclosed in SEQ ID NO:37. This clone contains a large region of the rpoN gene (CT609) in frame with β -gal and the complete ORF for the hypothetical gene CT610 on the complementary strand. In addition, it also contains the carboxy-terminal end of CT611, another hypothetical gene.

 Clone E7-H11-10 (identified using the CT3 patient line) contains a 1990
25 bp insert, the sequence of which is disclosed in SEQ ID NO:36. This clone contains the amino terminal partial ORF for CT610, a complete ORF for CT611, another complete ORF for CT612, and a carboxy-terminal portion of CT613. All of these genes are hypothetical and all are present on the complementary strand.

 Clone E10-C6-45 (identified using the CT3 patient line) contains a 196
30 bp insert, the sequence of which is disclosed in SEQ ID NO:35. This clone contains a partial ORF for nrdA (CT827) in frame with β -gal. This clone contains a relatively

small insert and has particular utility in determining the epitope of this gene that contributes to the immunogenicity of Serovar E.

Clone E3-H6-10 (identified using the CT12 patient line) contains a 3734 bp insert, the sequence of which is disclosed in SEQ ID NO:48. This clone contains
5 ORFs for a series of hypothetical proteins. It contains the partial ORFs for CT223 and CT229 and the complete ORFs for CT224, CT225, CT226, CT227, and CT228.

Clone E4-C3-40 (identified using the CT10 patient line) contains a 2044 bp insert, the sequence of which is disclosed in SEQ ID NO:47. This clone contains a partial ORF for *nrdA* (CT827) and the complete ORF for *nrdB* (CT828).

10 Clone E2-D8-19 (identified using the CT1 patient line) contains a 2010 bp insert, the sequence of which is disclosed in SEQ ID NO:46. This clone contains ORF from the *Chlamydia trachomatis* plasmid as well as containing partial ORFs for ORF3 and ORF6, and complete ORFs for ORF4 and ORF5.

15 Clone E3-D10-46 (identified using the patient lines CT1, CT3, CT4, CT11, and CT12) contains a 1666 bp insert, the sequence of which is identified in SEQ ID NO: 45. This clone contains a partial ORF for CT770 (*fab F*), a complete ORF for CT771 (hydrolase/phosphatase homologue), a complete ORF for CT772 (*ppa*, inorganic phosphatase), and a partial ORF for CT773 (*ldh*, Leucine dehydrogenase).

20 Clone E10-H8-1 (identified using both the CT3 and CT10 patient lines) contains an 1862 bp insert, the sequence of which is disclosed in SEQ ID NO:44. It contains the partial ORFs for CT871 (*pmpG*) as well as CT872 (*pmpH*).

Clone E3-F3-7 (identified using the CT1 patient line) contains a 1643 bp insert, the sequence of which is identified in SEQ ID NO:43. It contains the partial ORFs for both CT869 (*pmpE*) and CT870 (*pmpF*).

25

EXAMPLE 3

ADDITIONAL CD4 T CELL EXPRESSION CLONING FOR THE IDENTIFICATION OF T CELL STIMULATING ANTIGENS FROM CHLAMYDIA TRACHOMATIS SEROVAR E

30

The T cell line CHH037 was generated from a 22 year-old healthy female sero-negative for *Chlamydia*. This line was used to screen the *Chlamydia*

trachomatis serovar E library. Nineteen clones were identified from this screen, as described below.

Clone E7-B12-65, contains an 1179 bp insert, the sequence of which is disclosed in SEQ ID NO:114. It contains the complete ORF of the gene for Malate
5 dehydrogenase (CT376) on the complementary strand.

Clone E4-H9-83 contains a 772 bp insert, the sequence of which is identified in SEQ ID NO:115. It contains the partial ORF for the heat shock protein GroEL (CT110).

Clone E9-B10-52 contains a 487 bp insert, the sequence of which is
10 identified in SEQ ID NO:116. It contains a partial ORF for the gene yscC (CT674), a general secretion pathway protein.

Clone E7-A7-79 contains a 1014 bp insert, the sequence of which is disclosed in SEQ ID NO:117. It contains the complete ORF for the histone like development gene, hetA (CT743) and a partial ORF for the rRNA methyltransferase
15 gene ygcA (CT742).

Clone E2-D11-18 contains a 287 bp insert, the sequence of which is disclosed in SEQ ID NO:118. It contains the partial ORF for hetA (CT743).

Clone E9-H6-15, identified using the CT3 line, contains a 713 bp insert
20 the sequence of which is disclosed in SEQ ID NO:125. It contains the partial ORF of the pmpB gene (CT413).

Clone E3-D10-87, identified using the CT1 line, contains a 780 bp insert, the sequence of which is disclosed in SEQ ID NO:126. It contains the partial ORF for CT388, a hypothetical gene, on the complementary strand, and a partial ORF
25 for CT389, another hypothetical protein.

Clone E9-D6-43, identified using the CT3 line, contains a 433 bp insert, the sequence of which is disclosed in SEQ ID NO:127. It contains a partial ORF for CT858.

Clone E3-D10-4, identified using the CT1 line, contains an 803 bp
30 insert, the sequence of which is disclosed in SEQ ID NO:128. It contains a partial ORF for pGP3-D, an ORF encoded on the plasmid pCHL1.

Clone E3-G8-7, identified using the CT1 line, contains an 842 bp insert, the sequence of which is disclosed in SEQ ID NO:129. It contains partial ORFs for CT557 (Lpda) and CT558 (LipA).

5 Clone E3-F11-32, identified using the CT1 line, contains an 813 bp insert, the sequence of which is disclosed in SEQ ID NO:130. It contains a partial ORF for pmpD (CT812).

Clone E2-F8-5, identified using the CT12 line, contains a 1947 bp insert, the sequence of which is disclosed in SEQ ID NO:131. It contains a complete ORF for the 15 kDa ORF (CT442) and a partial ORF for the 60 kDa ORF (CT443).

10 Clone E2-G4-39, identified using the CT12 line, contains a 1278 bp insert, the sequence of which is disclosed in SEQ ID NO:132. It contains the partial ORF of the 60kDa ORF (CT443).

Clone E9-D1-16, identified using the CT10 line, contains a 916 bp insert, the sequence of which is disclosed in SEQ ID NO:133. It contains the partial
15 ORF for the pmpH (CT872).

Clone E3-F3-6, identified using the CT1 line, contains a 751 bp insert, the sequence of which is disclosed in SEQ ID NO:134. It contains the partial ORFs, all on the complementary strand, for genes accB (CT123), L13 ribosomal (CT125), and S9 ribosomal (CT126).

20 Clone E2-D4-70, identified using the CT12 line, contains a 410 bp insert, the sequence of which is disclosed in SEQ ID NO:135. It contains the partial ORF for the pmpC gene (CT414).

Clone E5-A1-79, identified using the CT1 line, contains a 2719 bp insert, the sequence of which is disclosed in SEQ ID NO:136. It contains a partial ORF
25 for ydhO (CT127), a complete ORF for S9 ribosomal gene (CT126 on the complementary strand), a complete ORF for the L13 ribosomal gene (CT125 on the complementary strand) and a partial ORF for accC (CT124 on the complementary strand).

Clone E1-F7-16, identified using the lines CT12, CT3, and CT11,
30 contains a 2354 bp insert, the sequence of which is disclosed in SEQ ID NO:137. It contains a partial ORF of the ftsH gene (CT841) and the entire ORF for the pnp gene (CT842) on the complementary strand.

Clone E1-D8-62, identified using the CT12 line, contains an 898 bp insert, the sequence of which is disclosed in SEQ ID NO:138. It contains partial ORFs for the *ftsH* gene (CT841) and for the *pnp* gene (CT842).

5

EXAMPLE 4

EXPRESSION OF CHLAMYDIA TRACOMATIS RECOMBINANT PROTEINS

Several *Chlamydia trachomatis* serovar E specific genes were cloned into pET17b. This plasmid incorporates a 6X histidine tag at the N-terminal to allow for
10 expression and purification of recombinant protein.

Two full-length recombinant proteins, CT622 and CT875, were expressed in *E. coli*. Both of these genes were identified using CtlGVII expression screening, but the serovar E homologues were expressed. The primers used to amplify these genes were based on serovar D sequences. The genes were amplified using
15 serovar E genomic DNA as the template. Once amplified, the fragments were cloned in pET-17b with a N-terminal 6X-His Tag. After transforming the recombinant plasmid in XL-I blue cells, the DNA was prepared and the clones fully sequenced. The DNA was then transformed into the expression host BL21-pLysS cells (Novagen) for production of the recombinant proteins. The proteins were induced with IPTG and purified on Ni-
20 NTA agarose using standard methods. The DNA sequences for CTE622 and CTE875 are disclosed in SEQ ID NO:28 and 27 respectively, and their amino acid sequences are disclosed in SEQ ID NO: 140 and 139, respectively

Five additional *Chlamydia trachomatis* genes were cloned. The *Chlamydia trachomatis* specific protein CT694, the protein CT695, and the L1
25 ribosomal protein, the DNA sequences of which are disclosed in SEQ ID NO:119, 120 and 121 respectively. The protein sequences of these 6X-histidine recombinant proteins are disclosed in SEQ ID NO: 122 (CT694), 123 (CT695), and 124 (L1 ribosomal protein). The genes CT875 and CT622, from serovar E were also cloned using pET17b as 6X-His fusion proteins. These recombinant proteins were expressed and purified and
30 their amino acid sequences disclosed in SEQ ID NO:139 and 140, respectively.

EXAMPLE 5

RECOMBINANT CHLAMYDIAL ANTIGENS RECOGNIZED BY T CELL LINES

- Patient T cell lines were generated from the following donors: CT1, CT2, CT3,
 5 CT4, CT5, CT6, CT7, CT8, CT9, CT10, CT11, CT12, CT13, CT14, CT15, and CT16.

A summary of their details is included in Table II.

Table II: <i>C. trachomatis</i> patients						
Patients	Gender	Age	Clinical Manifestation	Serovar	IgG titer	Multiple Infections
CT1	M	27	NGU	LCR	Negative	No
CT2	M	24	NGU	D	Negative	E
CT3	M	43	Asymptomatic Shed Eb Dx was HPV	J	Ct 1:512 Cp 1:1024 Cps 1:256	No
CT4	F	25	Asymptomatic Shed Eb	J	Ct 1:1024	Y
CT5	F	27	BV	LCR	Ct 1:256 Cp 1:256	F/F
CT6	M	26	Perinial rash Discharge, dysuria	G	Cp 1:1024	N
CT7	F	29	BV Genital ulcer	E	Ct 1:512 Cp 1:1024	N
CT8	F	24	Not Known	LCR	Not tested	NA
CT9	M	24	asymptomatic	LCR	Ct 1:128 Cp 1:128	N

CT10	F	20	Mild itch vulvar	negative	negative	12/1/98
CT11	F	21	BV Abnormal pap smear	J	Ct 1:512	F/F/J/E/E PID 6/96
CT12	M	20	asymptomatic	LCR	Cp 1:512	N
CT13	F	18	BV, gonorrhea, Ct vaginal discharge, dysuria	G	Ct 1:1024	N
CT14	M	24	NGU	LCR	Ct 1:256 Cp 1:256	N
CT15	F	21	Muco-purulent cervicitis Vaginal discharge	culture	Ct 1:256 Ct IgM 1:320 Cp 1:64	N
CT16	M	26	Asymptomatic/ contact	LCR	NA	N
CL8	M	38	No clinical history of disease	negative	negative	No

NGU=Non-Gonococcal Urethritis; BV=Bacterial Vaginosis; CT=Chlamydia trachomatis; Cp=Chlamydia pneumoniae; Eb=Chlamydia elementary bodies; HPV=human papilloma virus; Dx=diagnosis; PID=pelvic inflammatory disease;

5 LCR=Ligase change reaction.

PBMC were collected from a second series of donors and T cell lines have been generated from a sub-set of these. A summary of the details for three such T cell lines is listed in the table below.

Table III: Normal Donors

Donor	Gender	Age	CT IgG Titer	CP IgG Titer
CHH011	F	49	1:64	1:16
CHH037	F	22	0	0
CHH042	F	25	0	1:16

Donor CHH011 is a healthy 49 year old female donor sero-negative for *C. trachomatis*. PBMC produced higher quantities of IFN-gamma in response to *C. trachomatis* elementary bodies as compared to *C. pneumoniae* elementary bodies, indicating a *C. trachomatis*-specific response. Donor CHH037 is a 22 year old healthy female donor sero-negative for *C. trachomatis*. PBMC produced higher quantities of IFN-gamma in response to *C. trachomatis* elementary bodies as compared to *C. pneumoniae* elementary bodies, indicating a *C. trachomatis*-specific response. CHH042 is a 25 year old healthy female donor with an IgG titer of 1:16 to *C. pneumoniae*. PBMC produced higher quantities of IFN-gamma in response to *C. trachomatis* elementary bodies as compared to *C. pneumoniae* elementary bodies, indicating a *C. trachomatis*-specific response.

Recombinant proteins for several *Chlamydia trachomatis* genes were generated as described above. Sequences for MOMP was derived from serovar F. The genes CT875, CT622, pmp-B-2, pmpA, and CT529 were derived from serovar E and sequences for the genes gro-EL, Swib, pmpD, pmpG, TSA, CT610, pmpC, pmpE, S13, lpdA, pmpI, and pmpH-C were derived from LII.

Several of the patient and donor lines described above were tested against the recombinant Chlamydia proteins. Table IV summarizes the results of the T cell responses to the recombinant Chlamydia proteins.

Table IV: Recombinant Chlamydia Antigens Recognized By T Cell Lines

Antigen	Serovar	#of hits	CL 8 L2	CT 10 E	CT 1 E	CT 3 E	CT 4 L2	CT 5 E	CT 11 E	CT 12 E	CT 13 E	CHH- 011 E	CHH- 037 E
gro-EL (CT110)	L2	10	-	+	+	+	+	+	+	+	+	+	+
MompF (CT681)	F	10	-	+	+	+	+	+	+	+	+	+	+
CT875	E	8	-	+	+	-	+	+	+	+	+	-	+
SWIB (CT460)	L2	8	+	+	-	+	-	+	-	+	+	+	+
pmpD (CT812)	L2	5	-	+	+	+	+	-	-	+	+	-	-
pmpG (CT871)	L2	6	-	+	+	-	+	+	nt	-	+	+	-
TSA (CT603)	L2	6	-	-	+	+	+	+	-	-	+	-	+
CT622	E	3	-	-	+	-	+	-	-	-	+	-	-
CT610	L2	3	-	+	-	+	-	-	-	+	-	-	-
pmpB-2 (CT413)	E	3	-	-	+	+	+	-	-	-	-	-	-
pmpC (CT414)	L2	4	-	-	-	+	-	+	-	+	-	-	+
pmpE (CT869)	L2	3	-	+	+	-	-	-	-	-	+	-	-
S13 (CT509)	L2	2	+	-	-	-	+	-	-	-	-	-	-
lpdA (CT557)	L2	3	-	-	+	+	-	-	-	-	-	+	-
pmpI (CT874)	L2	2	-	-	+	-	-	-	-	-	-	+	-
pmpH-C (CT872)	L2	1	-	-	-	-	-	-	-	+	-	-	-

pmpA (CT412)	E	0	-	-	-	-	-	-	-	-	-	-	-
CT529	E	0	-	-	-	-	-	-	-	-	-	-	-

Although the present invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, changes and
5 modifications can be carried out without departing from the scope of the invention which is intended to be limited only by the scope of the appended claims.

Claims

What is Claimed:

1. An isolated polynucleotide comprising a sequence selected from the group consisting of:

- (a) sequences provided in SEQ ID NO:1-48, 114-121, and 125-138;
- (b) complements of the sequences provided in SEQ ID NO: 1-48, 114-121, and 125-138;
- (c) sequences consisting of at least 20 contiguous residues of a sequence provided in SEQ ID NO:1-48, 114-121, 125-138;
- (d) sequences that hybridize to a sequence provided in SEQ ID NO:1-48, 114-121, and 125-138, under highly stringent conditions;
- (e) sequences having at least 95% identity to a sequence of SEQ ID NO:1-48, 114-121, and 125-138;
- (f) sequences having at least 99% identity to a sequence of SEQ ID NO: 1-48, 114-121, and 125-138; and
- (g) degenerate variants of a sequence provided in SEQ ID NO: 1-48, 114-121, and 125-138.

2. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- (a) sequences encoded by a polynucleotide of claim 1;
- (b) sequences having at least 95% identity to a sequence encoded by a polynucleotide of claim 1; and
- (c) sequences having at least 99% identity to a sequence encoded by a polynucleotide of claim 1.

3. An isolated polypeptide comprising at least an immunogenic fragment of a polypeptide sequence selected from the group consisting of:

- (a) a polypeptide sequence set forth in SEQ ID NO: 122-124 and 139-140,

(b) a polypeptide sequence having at least 95% identity with a sequence set forth in SEQ ID NO: 122-124 and 139-140, and

(c) a polypeptide sequence having at least 99% identity with a sequence set forth in SEQ ID NO: 122-124 and 139-140.

4. An expression vector comprising a polynucleotide of claim 1 operably linked to an expression control sequence.

5. A host cell transformed or transfected with an expression vector according to claim 4.

6. An isolated antibody, or antigen-binding fragment thereof, that specifically binds to a polypeptide of claim 2 or claim 3.

7. A method for detecting the presence of Chlamydia in a patient, comprising the steps of:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with a binding agent that binds to a polypeptide of claim 2 or claim 3;
- (c) detecting in the sample an amount of polypeptide that binds to the binding agent; and
- (d) comparing the amount of polypeptide to a predetermined cut-off value and therefrom determining the presence of Chlamydia in the patient.

8. A fusion protein comprising at least one polypeptide according to claim 2 or claim 3.

9. An oligonucleotide that hybridizes to a sequence recited in any one of SEQ ID NO: 1-48, 114-121, and 125-138 under highly stringent conditions.

10. A method for stimulating and/or expanding T cells specific for a Chlamydia protein, comprising contacting T cells with at least one component selected from the group consisting of:

- (a) a polypeptide according to claim 2 or claim 3;
- (b) a polynucleotide according to claim 1; and
- (c) an antigen-presenting cell that expresses a polynucleotide

according to claim 1,

under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells.

11. An isolated T cell population, comprising T cells prepared according to the method of claim 10.

12. A composition comprising a first component selected from the group consisting of physiologically acceptable carriers and immunostimulants, and a second component selected from the group consisting of:

- (a) a polypeptide according to claim 2 or claim 3;
- (b) a polynucleotide according to claim 1;
- (c) an antibody according to claim 6;
- (d) a fusion protein according to claim 8;
- (e) a T cell population according to claim 11; and
- (f) an antigen presenting cell that expresses a polypeptide according

to claim 2 or claim 3.

13. A method for stimulating an immune response in a patient, comprising administering to the patient a composition selected from the group consisting of:

- (a) a composition of claim 12;
- (b) a polynucleotide sequence of any one of SEQ ID NO:80-94; and
- (c) a polypeptide sequence of any one of SEQ ID NO:95-109.

14. A method for the treatment of Chlamydia infection in a patient,

comprising administering to the patient a composition selected from the group consisting of:

- (a) a composition of claim 12;
- (b) a polynucleotide sequence of any one of SEQ ID NO:80-94; and
- (d) a polypeptide sequence of any one of SEQ ID NO:95-109.

15. A method for determining the presence of Chlamydia in a patient, comprising the steps of:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with an oligonucleotide according to claim 9;
- (c) detecting in the sample an amount of a polynucleotide that hybridizes to the oligonucleotide; and
- (d) comparing the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, and therefore determining the presence of the cancer in the patient.

16. A diagnostic kit comprising at least one oligonucleotide according to claim 9.

17. A diagnostic kit comprising at least one antibody according to claim 6 and a detection reagent, wherein the detection reagent comprises a reporter group.

18. A method for the treatment of Chlamydia in a patient, comprising the steps of:

- (a) incubating CD4+ and/or CD8+ T cells isolated from a patient with at least one component selected from the group consisting of:
 - (i) a polypeptide according to any one of claims 2 or 3;
 - (ii) a polypeptide sequence of any one of SEQ ID NO: 95-109;
 - (iii) a polynucleotide according to claim 1;
 - (iv) a polynucleotide sequence of any one of SEQ ID NO:80-94;

- (v) an antigen presenting cell that expresses a polypeptide sequence set forth in any one of claims 2 or 3;
- (vi) an antigen presenting cell that expresses a polypeptide sequence of any one of SEQ ID NO:95-109, such that the T cells proliferate; and
- (b) administering to the patient an effective amount of the proliferated T cells.

SEQUENCE LISTING

<110> Corixa Corporation
Bhatia, Ajay
Probst, Peter
Stromberg, Erika Jean

<120> COMPOUNDS AND METHODS FOR TREATMENT AND DIAGNOSIS
OF CHLAMYDIAL INFECTION

<130> 210121.515PC

<140> PCT

<141> 2001-04-23

<160> 140

<210> 1

<211> 1311

<212> DNA

<213> Chlamydia trachomatis

<400> 1

taattgctgt	ttacctctct	tottgctgaa	gacttggtta	tggtttttat	tttgacgata	60
aacctagtta	aggcataaaa	gagttgogaa	ggaagagccc	taaacctttc	ttatcatctt	120
ctttaactag	gagtcaccca	tgagtcaaaa	taagaactct	gctttcatgc	agcctgtgaa	180
cgatccgctg	gatttagctg	ccatcgttgg	tgacgagcct	atgcctcgca	cagagatcat	240
taagaaaaatg	tggtattaca	ttaagaagaa	tagccttcaa	gatcctacaa	acaaacgtaa	300
tatcaatccc	gatgataaat	tggtctaaat	ttttggaact	gaaaaacota	togataatgt	360
ccaaatgaca	aaaatggttt	ctcaacacat	cattaaataa	aatagaat	gactcaogtg	420
ttctctgctt	taaagatgag	gaactagttc	attctttttg	ttcgtttctg	tggttatatc	480
tgatatctta	acaactatct	tagcagcacc	tggttttgaca	tggttttggtg	ccaatcactt	540
agagcctaac	ctattgagag	taacgcgttt	aaattggaat	ctgcctaaaa	aaattgctca	600
tcttcattgg	cttcgcatta	tacagatttc	ggatttacac	ctaaacacat	cgaagcctga	660
tgccctttcta	aaaaaagtat	ctcgtaaagt	ctctctctct	tctccagata	ttctgtattt	720
tacaggagac	tttgtctgtc	gcgctaaagt	agaaactcct	gaaagattaa	aacatttcct	780
atgttctctg	catgcgccct	taggtctgtt	tgcttgcccta	ggaatcatg	attacgccac	840
ctacgtatcc	cggtgatatt	acgggaaaaat	taataccatc	tcagcaatga	atagccgtcc	900
tttaaaaaaga	gctttttacct	ctgttttaca	aagtctattc	gcctcttctc	gcaatgaatt	960
tgagatact	ctgaatccac	aaattctcaa	tccacaccta	gtcagtatat	tacgcaatc	1020
tccatttcaa	tatttgcata	atcaaaagcgc	gacactttcc	gatacataca	acatcgtggg	1080
attagcgcat	ttttttgcca	acaaattcga	tcccaaaaaa	gcttttactg	actataatcc	1140
caagttacct	ggtattatcc	tttctcataa	tcccgatagc	attcaccatc	tccaagatta	1200
ccaggtgat	gtgttttttt	ccgggcactc	gcattgcccct	caaatctctc	ttccctggcc	1260
taagtgtgcc	aatacagata	ccaataaact	ttcagggtta	gaaaaccagg	a	1311

<210> 2

<211> 1516

<212> DNA

<213> Chlamydia trachomatis

<400> 2

tttgagctcg	tgccgctcgt	gccggtgcgt	gtgaaccgct	tcttcaaaag	cttgtcttaa	60
aagatattgt	ctcgcttcgg	gattagttac	atgtttaaaa	attgctagaa	caatattatt	120
cccaaccaag	ctctctcggt	tgctgaaaaa	acctaatttc	aaaagaatga	ctcgccgcctc	180
atcttcagaa	agaagatccg	acttccataa	ttcgtatgct	ttcccatggg	ggatctctgt	240
agggaagcag	ttattttgct	agccattcaa	ataatgttcc	caagcccatc	tgtaacttaat	300
aggaaacaagt	tggttgacat	cgaactggtt	gcagtctcat	agacgcttgc	tatttagatt	360
aaacgcttgt	tggttttccat	ctaaaaatcc	tgcttgcata	agaacgctta	attttattgt	420
tataatttat	gatttaattac	tgacatgctt	cacacacgtc	ttccaaagaa	cattagagtg	480
ctttctctgc	cttttcaaca	ataattctct	ccgaagcaga	cttattcttc	atccaaacag	540

gctgaattcc	tctcttatta	atatctacaa	aagatttttc	aacggtcggt	gctgatgaag	600
atctcagata	atcgttagtt	ttcaaacctt	ttttccaagc	cggttaaatac	atattcgaca	660
gtttttttcc	gtctgcgtgg	gcaagataaa	gggtgaggga	ttgcccacata	tcaatccatt	720
tttgtctctg	agacgcgcac	tgcataatcc	atctcgtgtc	aatctcaaaa	gctgtcaaga	780
aaatagtgtt	taagtgtatc	ggtatacgtc	cgattttocaa	taaagaccoca	tcaaaattatt	840
tcaggtctac	taacatatca	gcataccaga	taacctaat	cttcaacctc	tcaattaaat	900
acacatttgg	acaactggaat	tctccggaca	aattagactc	cacaaacaaa	gtttgttagc	960
ttggctcaat	agattgagtt	aotcctataa	tggtggagat	cgctcgctgc	ggagctatag	1020
coataagctg	acaatgtcgc	ataccatgct	ctttaaccaa	actacggata	ggttcccaat	1080
ctttttctga	tgcagtatcc	atctggagat	ttgctttccc	tcgatagtgc	gctaacaact	1140
gaactgtatc	ataagggagc	aaacctctat	cccatcttgc	tcctttataa	gagctgaag	1200
tgctcgtctc	tttagcgagc	agacaagaag	cttgaatcgc	atagtataag	atcaactctg	1260
aactgtagtc	agcaaatctc	acagcttctt	gcgaagcata	gcttataatc	agcttataca	1320
agggactctg	gaatcccatc	acccttaatc	caatagcgcg	gtgagcaaa	ttcgcccttt	1380
tagcttctct	tggttgataa	aagttaatat	caatcacggt	atccacacata	caggatcgat	1440
ttagagatgt	ctcagagagt	tttctctcat	caaacccatc	coctacgata	tggtgaacta	1500
agttaatoga	tcttaa					1516

<210> 3

<211> 2397

<212> DNA

<213> Chlamydia trachomatis

<400> 3

agagtgtgct	ggaggagcta	tttttgcaaa	acgggttcgt	attgtagata	accaaggagg	60
cggtgtattc	tgcacaacct	tctctgatat	tctatggcgc	gccatttita	caggtctctc	120
ggagagaag	gataaagttag	atgggcataa	ccctgaagtc	ttgatctcag	gcaatgcagg	180
tgatgttgtt	ttttccggaa	attcctcgaa	gcgtgatgag	catcttctcc	atacaggtgg	240
ggagagcatt	tgtactcaaa	atttgacgat	ttctcagaat	acaagggaatg	ttctgtttta	300
taacacacgtg	gcctgttcgg	gaggagctgt	tcgtatagag	gatcatggtt	atgtttcttt	360
agaagctttt	ggaggagata	ttgtttttaa	aggaattctc	tctttcagag	cacaaggatc	420
cgatgcctac	tattttgcag	gtaaaagaac	gcattattaca	gccctgaatg	ctcagggaag	480
acatgctatt	gttttccacg	acgcattagt	ttttgaaaa	ctagaagaaa	ggaattctgc	540
tgaagtattg	ttaatacaata	gtcgagaaaa	tccaggttca	aaatttctca	agtttgatgc	600
aattgtgcta	ttcgtacctc	ttagtttctc	atgtccacgg	taaagggtac	ggaaagatac	660
gcattttatt	tcatagtctt	tagcttcgat	ccctagtgtc	tcgcacatga	ctcgtctgcc	720
aagacttttg	gttacgaaaa	caacaggctc	tcgttgagaa	atgatttgga	gtagctctag	780
cgtagggtgt	ttttctgtgt	tctcgtgtgt	tgaagatttg	actagaggag	agacttcaat	840
acataactcg	ctgcggtttt	ttaataaaat	ttgaccagag	gagggtcttt	ccgactgtcc	900
tagtaataga	cgaaattatg	ccaatgctct	gtgaaacgtt	ttccctgatt	catctcgaaa	960
ctttgcgcag	gattccaatt	cttcgattac	tgtaaaaggg	ataatgatgc	gaggtgttag	1020
aaaaggagaa	agggccttag	gatcgttaaa	caaaacgctg	gtatcaataa	caaagaggtt	1080
ttctactaca	aattctcaaa	tgaactcaag	gtaaggggga	gatagtactt	tgattgtgta	1140
tcatatccag	aaaaattaaa	acatgtcttt	gttagagaga	agtcggggga	gaggggtttt	1200
agcaatcaac	ctccgcgtgt	gctaactctg	ttgtcaaaaa	tgtaaccttc	aactacaagt	1260
cgaagagaa	cgagctcttc	tggttgaggt	tgattgaaa	gtcaaaatta	tcatcagatt	1320
catgttgatt	tcccaataca	ttctcgtcgc	atggaatcag	atagctttca	tagagctgtg	1380
tgatggaggg	acggagaggg	gtattacttt	gaaactccat	ttaattgatg	gagagacagt	1440
ctctatacc	aatctaggac	aagcagattg	tgatgaggtg	ttccaagagc	acttgctata	1500
tttagagctc	acagctcctc	agaaaaacaa	ggaagaggaa	aaaattagct	ctttgttagg	1560
agctgttcaa	caaatggcta	aagatgcga	agtaacaggt	ttttctcaaa	agggcttggt	1620
ttctgttgta	ctaggaggag	ctggttcgat	taatatgttg	ttgcaacatt	ctccagaaa	1680
taagatcatc	ctgatctctc	ctacccgatt	actggagagg	atagcgcaaa	tgatgcgttc	1740
attatctata	ggaccaactt	ctatttttag	taagccagag	ccctattgca	actgttttgc	1800
ttgtcaaaat	ggacgagcta	cagtggaaag	agaggtatgc	ggagtatcgg	atgagatctc	1860
caacttttgt	tcatgggata	tctctcaaa	tggagaaaag	gtgtacactg	ttcaagatcc	1920
tttgaataca	gaagtatacc	ttttgttttt	tttatacag	ccagcactcc	aaattctcag	1980
tgtagagaata	tatcataaaat	agacgggctc	ctagcgtctc	gaatagaaaa	agctcttctg	2040
atagacactat	caagcctctc	ctttatacgc	tcaagcaata	gaacgggaga	tctcagcaat	2100
ggatttttgt	gtactcat	aaacgagcgg	aaaatgaaat	tactcaaat	ttcttcaggt	2160
ctacacacgc	tcaaatcatc	gaggaaaacc	gtatgagaaa	cggatctact	cggtccgaat	2220

tcggcacgag gtctctaatt ttgcagaagg agcacaaatt ttgtctgtcc aagggttaaa 2280
 tcttgcgtgga gaataaggat actgccctcc ttgccctcca gatgcgaagc atcgctatta 2340
 cttttatgct tatgcgtcgc atgtgtgtgct ttccgatgaa gaaggagtga ccaaga 2397

<210> 4
 <211> 1094
 <212> DNA
 <213> Chlamydia trachomatis

<400> 4
 tgaatgcagaa gacactgita agaagtitaca agaagccggt gctaaggctg ttgtctaaagg 60
 gctgtaattg ttatgggaaa gagaatgctt tgggggttgc ttgcaagctt ctctttttctg 120
 ttatgctgcac agtagctggg cacagagggg ttcccggtac gtcttaacag atttgcctgg 180
 acttaacttt tagtgtttgg catcgcaaac agaataatc ttgtgcaatg gttttttctt 240
 aatggaatca aggtgatagt atttgtcgga tggacaagt gtgacaagt atctgtccta ttccgtgtta 300
 tctgtattgg atagactctg ttttgtccta gtctgcaaac atctgtccta ttccgtgtta 360
 gagatcacag agggactaaa tagggaatg gtatcgccaa aagctcttaa gtctttaggag 420
 agctgcagtg ttaagtgc ccgagcgggt cagcgtcaaa aagaaagaag aattttttaa 480
 ttctctaat ctgttcgaag tcaaatcaa gtctataag cagttttctc aaatcgggaa 540
 gcttgctgaa gagcgagaaa acattgtttt agaagaagtc ttcaagaa ttcttcctca 600
 caagctttat aaagaagcta cgtattttaga gtacctctc ttacaagttag gagggtccaa 660
 taactcccaca gaagagtgtta ttgtcgtggg aatcacctat agtgttaact taagggttgg 720
 ttccgtgtta actgatgaaa cggggattaa agaagaagaa gtctatatgg gaacctccoc 780
 catcatgact caaaaggaaa cctttattat taatggggca gagagagctg ttgtttctca 840
 agtcaacgtg tctccagaaa tcaattttga acagaaaaaa catctcaaa ggaattgttt 900
 attttttttt agaattatc ctatttttga aagttgtgaa gaagctgtct tgcagattaa 960
 tgcacatttc tatatccata ttgataggaa aaaaagtgc agaaagatt tagctattga 1020
 cggttatccg agtttttaga tattcaaacg atgcagatat tattgaagag ttcttttctg 1080
 tagaggagcg ttcc 1094

<210> 5
 <211> 2129
 <212> DNA
 <213> Chlamydia trachomatis

<400> 5
 gcttttttaa gagataagca acaaccgag aatccactcc tccagacata gcaacaatga 60
 tagtttttac cacaatgagc ccagaaaacg ctttctgtta ttgaagtgtg cacattcaaa 120
 agggccatca tggttagcaaa aaaacaggat caaaaaaac tatttctcaa gcgcctgttt 180
 ttaaatctta attacaaaaa taataatcaa ttcaactttt caaaaaaga atttaaacat 240
 taattgttat aaaaacaata ttattataaa aataataacc atagtgtgog ggaatctct 300
 ttcatgtgtt atttttaggc tcatcaacct aggcatacgc ctaaaaact tcttttggaa 360
 gttcaccaat cgtttccaga taagcatcct caaattgcta aagctatgcg gattacgggg 420
 ataaccctcg cagctotatc tctgtcgtcg gtatcgctcg cgtctatgcg cgtctctgcg 480
 ggagagctcg ccattctctt tgcgtcatt ggtggaattg ctgcaatgic tgccctctta 540
 tctcgtgcga ccattatctg ttctgcacaa aaggctctgg ctcaacgaaa caaaaaacaa 600
 ctagaagagt tgcttcggtt agataatgog accgagcatg tgaattacct gacctcagac 660
 acctcttatt ttaatacagt ggaatcctta gatgctctaa ataacagtt ttctcagatt 720
 gacttaacta ttcaagctcc cgaaaaaaaa ctattaaaaa aagtcttgg ttccagatgc 780
 gatttcaacta taactccat cgaagagatc tccgatcgct ttacgaaaa gtctctcttt 840
 cttcgattaa gagaacattt ttgtcggaga gaagagcgtt atgcccccta tttaagccct 900
 cctctactta acaagaatg ttgtgcgaac caatcacat caaatatgat taggatgcta 960
 ccaaaatccg gtgtgttttt ttccctcaaa gccaatcac taagtcatgc cagccgcaca 1020
 ctatatacag tattgaaagt cgttttatcc ttaggagttc tcgctggagt cgtcgtcttt 1080
 atcatcttcc ttcccccctg cctgcctttt atcgctgtta taggagatc ttcccttagca 1140
 ttgggagtg catctttctt tatgattcgg ggcattaaat attgtctcga acattctctc 1200
 ctgaatagaa agcaattagc taaagatttt caaaaaacca ttatcccgga tgcctgtgcc 1260
 tctatgtgttc attaccgca tcaattacta tcaactctac atgaaactct attagatgaa 1320
 gccatcacag ctatagtgag cgagcccttc ttatttgaa acgctaatct taaggcaaaa 1380
 attgaagatt tgacaaaaca atatgatata ttgaacgag ctttaataa actcttttaa 1440
 caagatgagc cgtcctgttc tcaattagag aaacgagctt acttattccc aattcctaatt 1500

aacgacgaaa	atgctaaaac	taaagaatcg	cagcttctag	actcagaaaa	tgattcfaat	1560
tctgaatttc	aggagattat	aaataaagga	ctagaagctg	ccaataaacg	acgagctgac	1620
gctaagtcaa	aattctatag	ggaagacgaa	acotctgaca	aaagattctc	tatatggaaa	1680
cccaacaaga	acttggcatt	agaagatttg	tggagagctg	atgaagcttg	caatgaagag	1740
caacaagctc	tcctcttaga	agatttatag	agttataaaa	cctcagaatg	tcaagctgca	1800
ctccaaaaag	tgagtcaaga	actgaagcgg	gcacaaaaat	cattcgcaat	cctgaaaaag	1860
catgctctag	acagacttta	tgaatccagt	gtagccatga	tggatttagc	tagagcgaat	1920
caagaaacac	accggtctct	gaacatctct	tctgaattac	aaacaactag	acaataacctg	1980
ttagataatc	actaacgggt	ctctataaat	gacaaaaaga	aaaaggagag	ctgtgtctgt	2040
gctctctttt	ttctctaaat	attctgtaaa	gactaacctt	tttatgggtg	cgttgagcct	2100
cctctctctg	ttcccgagga	gcccgcaac				2129

<210> 6

<211> 288

<212> DNA

<213> Chlamydia trachomatis

<400> 6

gagggagcag	cctaactctc	ccctctcttc	ttaaaaaga	ggggagcctt	tttctcttac	60
aaagatacgc	tagcttittc	ctgaagaatc	tcatacaagag	atattttgcat	tttccacagg	120
ataaaggcoat	ccaaagggaag	ccctggaatc	acttcabatt	ctcccggttg	tagcatttga	180
caaggggaac	caaagattaa	atcttccggt	aatccatagg	gattgtggctc	cgaacacact	240
ccggaagaaa	accattctcc	ttcttttggc	tgatatattg	atcgagcagc	ctctgtctaaa	300
gctcgtgctg	cagaagctgc	cgaagacttc	ccctgtgctt	cgattactgc	actacacaga	360
ctctgtacag	aaggcaccat	aattattctc	aaaccaatc	gattccgctat	cgctctcgct	420
ataggaaggt	catlaactcag	agctgtcgta	aaatcaggca	ctgtgttggc	ggagtgtatt	480
ccccaacaaca	caacttctga	tacagccgat	aaaggtlactt	ctgctctatg	cgataacatg	540
ctatgcatac	gattctggctc	caactgtagc	atcgcalgaa	agttctttct	caataactctg	600
ggagcctatg	ctactgtcat	ccagcaattg	gtattcaacg	ggctcccaac	aacaaaaatc	660
tttgcatacc	ctgtgtgctg	tggtgtcaaa	gcttttccct	gctgtagcaa	aatctcccca	720
tttttcttta	gaagatccct	tcctccatct	ctcgggcttc	taggaactga	ccctataagg	780
aatgcgcgat	caatgcacac	aaaagcatca	tgcacatgat	tcgttacctg	caacagctgt	840
aataaaggga	aagcaccatc	atctagcttc	atggcgcac	cagataaagc	cccttctgtt	900
ccaggaaatc	ctagatacgc	cagatcgatg	ccacaatcaa	ggccaaaaac	atctccatga	960
gcogagaaa	atagaagctt	ataggctatt	tgcctgttct	ctctgtttac	tgctaaactc	1020
aactgtttgag	aaacataaag	ccacctcttc	tttactttta	caaaacgcac	ataactctca	1080
caactcgttt	gcaactaaat	aattttggctc	ccaaacatag	tttgatgatg	aaaagaatca	1140
agtaacccat	ttctcttagta	aaagcttttg	gcaaaaaaaa	gctcatctat	ttttcaatg	1200
atgacgcgac	ttttaaactgaa	taagaactta	gaaaaactta	taaaaaatag	gcgcgtgtga	1260
tcctaccatc	atacttgatc	cgcagcgcac	aacttgttgt	cccttttttg	cagccaataa	1320
accgtggaca	tttaaaaaac	caataaaacg	tgcgcgaata	aagaacataa	agccctctaa	1380
aaaaacgattt	taagagagaa	gtaaatagaca	gattgtaaca	tatttaaaat	aaaaactctg	1440
caaaacaaaa	aactttgect	ggcgtctcc	gtagaagcaa	ctttatgtta	aaagctttaa	1500
aaagtcttaac	atacctcgag	cttcgggaaa	ctctacagga	gcattccccg	acatgatccg	1560
tataatttgc	gttgcccaatt	ctttccctaa	tgaaacccct	ctttgatcaa	aagaatttga	1620
ttcccaagcaa	aaccottgaa	atgcacaaatt	atgctcataa	aaagccaata	aactccacga	1680
aatacagagga	gaaagctgtt	gogctaccaa	tatcgaagaa	ggctgtgtcc	ctttaaacct	1740
cttattcggg	ttcgcattat	ctctacccgt	agctaaagct	aaagatttga	caacaaggtt	1800
tgcagaagagc	ttttgagatc	togtgcgc				1828

<210> 7

<211> 861

<212> DNA

<213> Chlamydia trachomatis

<400> 7

ggcgccacta	ctttaaagat	togtctctct	tttggtatac	cgagagaagt	tctgtgtgaa	60
tggcgttatg	ttcctgaag	tgtaggagat	ttggctacca	tagctctctc	tatcagggct	120
ccacagttac	agaaatcgat	gagaagcttt	ttccctaaag	aagatgatgc	gtttctacgg	180
tttagttcgc	tttctactc	tccaatgggt	cgcatttttt	ggcgagagct	tgccaatcat	240
tatgcacaga	gtgggttgaa	aagcgggtac	aatatgggga	gtaccagatg	gtttctccct	300

```

gtcattgggc ctgttatacg ggagtcggag ggttttttc gcgcttatat ttcttcgggtg 360
actgatgggg atggttaagag ccataaagta ggattttctaa gaatttctatc atatagttgg 420
caggacatgg aagatttttga tccttcaggga cgcctcctt gggaagaatt tgctaagatt 480
attcoagtat ttctttctaa tacagaagct ttgattatcg accaaacgaa caaccaggt 540
ggtagtgctc ttlatcttta tgcactgctt tccatgttga cagaccgtcc tttagaactt 600
ctgaacacata gaattgattct gactcaggat gaagtgtgtg atgctttaga ttggttaacc 660
ctgttggaaa acgtagacac aaacgtggag tctgcctctg ctctgggaga caacatgaa 720
ggatatactg tggatctaca ggttcogag tatttaaaaa gctttggacg tcaagtattg 780
aattgttggg gtaaaagggg tatcgagtta tcaacgccta ttctctttt ttggttttag 840
aagattcctc cacatcctcg a

```

<210> 8

<211> 763

<212> DNA

<213> *Chlamydia trachomatis*

<400> 8

```

ataacaaaaa catcttgatt atttttgtta aaagaaatc ttaatgagtt ttatttaatt 60
aacgaaacga aaagcttgct aatgaaaatt attcacacag ctatogaatt tgctccggta 120
atacaaggcg gaggcctggg agacgcgcta tacggactag caaaagcctt agccgctaat 180
cacacaacgg aagtgtgta ccttttatac cctaatttat ttactttgoc caaagaacaa 240
gatctttgct cgatccaaaa attatcttat ttttttgctg gagagcaaga agcaactgct 300
ttctctact ttatgaagg aattaaahta actctattca aactcgacac acgtccagag 360
ttatcggaga attcgggaac aatctacaca agcgatgatg ccttcgcttt ttgctgcttt 420
ttctgctgtg cgcctcctca catccaaaaa gaaggagcca atatcgttca tttaacagt 480
tgggatacag gatagttlge tggactactc aaacaacagc cctgctctca attcaaaaag 540
attgttctta cctacacataa ttttggttat cgaggctata caacacagga aatattagaa 600
gcctcctctt tgaatgaatt ttatatcage cagtaccaac tatllogcga tccacaaact 660
tgtgtgttgc taaaaggagc tttataactg tcagatttgc ttctcctaca 720
tacgcgaag aaattcttga agattattcc gattacgaaa ttc 763

```

<210> 9

<211> 665

<212> DNA

<213> *Chlamydia trachomatis*

<400> 9

```

ttgaaactaa aaacctaat tatttaaacg tcaaaataaa aaagagtttt aaaatgggaa 60
attctggttt ttatttgtat aacactgaaa actgcgtttt tctgataat atcaaatgtg 120
ggcaaatgac agacgcgctc aaggaccagc aaataatcct tgggacacaa toaacactgt 180
tcgcagccaa aatgacagct tctgatggaa tatotttaac agtctccaat aattcatcaa 240
ccaatgcttc tattacaatt ggtttggatg ogaaaaagc ttaccagctt attctagaaa 300
agttgggaga tcaaatcttt gatggaattg ctgatactat tgttgatagt acagtccaag 360
atatttttag caaaatcaaa acagaccctt ctctaggttt gtggaagct tttaacaact 420
ttccaatcac taataaaatt caatgcaacg ggttattcac tccagtaaac attgaaactt 480
tattgagagg aactgaataa ggaataattca cagtacaccc caaaagctct gggagcattg 540
tcttagtctc agcagatatt attgcatcaa gaatgggaag cgcggttgtt ctacttttgg 600
tacgagaagg tgattctaag cctgcgcgca ttagtattgg atactatca ggcattccta 660
attta

```

<210> 10

<211> 843

<212> DNA

<213> *Chlamydia trachomatis*

<400> 10

```

tgggaattgc gaagataacg attacgtttc cgtatctata ggagcgcgtt tgaatacaga 60
aaatatgtgc ttggataaag ctggtgttat ttgtgatgaa cgcggagtaa tccctacaga 120
tgccacattc cgacacaaag tacctaactat ttatgcttat ggagatataca caggaaatgt 180
gcaacttgcc ctatgagctt ctcatcaagg aatcattgca gcacggaata tgcctggcca 240
taaaaggaaa atcgattact ctgcgctccc ttctgtgac tttaccttcc ctgaagtgcg 300

```

ttcagtaggc	ctctcccaa	cagcagctca	acaacaaaa	atccccgtca	aagtaacaaa	360
attccatttt	cgagctattg	gaaaagcggg	cgcaatgggc	gaggccgatg	gatttgcagc	420
cattatcagc	catgagacta	ctcagcagat	cctaggagct	tatgtgattg	gcctcatgc	480
ctcatcctg	atttccgaaa	ttaccctcagc	agttcgtaat	gaactgactc	ttccttgat	540
ttacgaaact	atccacgcac	atccaacctt	agcagaagtt	tggcggtgaa	gtgcgtttgt	600
agotcgtgat	acccattacc	atatgcccc	tgtcaaaaa	tgaccgatcc	agatctctcc	660
actctcaaaa	actctatacc	cgccagattc	cctaagtcgc	taccgcagaa	actcccttta	720
gggcggtgat	ttgtccaaac	tgataatact	atcaaaaaata	aagggtctcc	tacagtctgt	780
gaggaagcct	cttgtccgaa	tgcaccccat	tgttggtcta	gacatacagc	tgacctatct	840
aga						843

<210> 11

<211> 1474

<212> DNA

<213> Chlamydia trachomatis

<400> 11

acagaaggga	cggcagagta	atcgatttcc	totttatggc	cagctatatt	ccgtgctgca	60
atgattccct	gatgagaagc	tacatgggca	agttgccatt	ttcctgtgat	atctccaata	120
tcataaattg	taggtacgtt	tgtgcgcatt	gtggcatcgg	tagggatgac	tcocggctta	180
gcacaataaa	caccagcttt	atccaaagca	atattttctg	tattcaaacg	gctcctata	240
gatacagaaa	cgtaatcgta	ttotcgcaga	ttccaatgta	tagttaacgc	aaacgcatct	300
actatatcct	caatatctga	tacagaggct	ttcagtaaga	aaacggagtc	ttgtcgggtg	360
gcttcgatca	acatggtttt	tgaaatatct	ggattattca	aagcaaggat	ttgagagctt	420
atcaacacaa	cgccaataat	ggagcctaac	gtatggaata	aggaagcgaa	ttgcgaacgc	480
gttagcagta	aaatccgagg	ggagcctatt	tgagggattt	ctttgaggtt	tagcaagcct	540
tcagagccgg	agattctcgc	gaaataaggaa	tcocggggaa	agctcgttgg	600	
atttttactt	tgctcagagt	aatggagtg	gctttgatta	agaagaggtt	ttctccataag	660
ctgcgaatga	gaccattaa	tcacccggg	ctctccagc	agaagacagt	gatcttatgt	720
accatagcgg	gatagttgat	gctgaatcct	ttcatatgaa	tcocaaactg	gtcagatctg	780
cgattttggg	taacgacttc	agctcctgct	aagagggctt	tagaaggaa	acacccctgg	840
tttcaaacgg	ttccgcagc	ctctcgtctt	tcgattagcg	cagtttttag	tcctgcttga	900
gggcaagtag	ttgtcgaac	atagcctcct	ggcccgctct	cgataactac	acagctgaaa	960
gcttcaattca	taacatttcc	tottcaatga	gtgttttaga	ttgcaacgat	ccatattgaa	1020
tgattatctg	aaggaagagg	attctccttc	caagcctttc	taggaaggg	aaagagaggt	1080
ccttccagaca	aatacatttc	ccggatttga	catctgggtg	gataaaatct	caatgagagg	1140
aagtggtagc	aggagagaaa	aaataggaa	gtaagagtg	tatttcgaat	gctcaggagg	1200
agagcggtac	ccacgataag	caagcagaat	cccgactagt	gcattatgat	atgagcgatt	1260
ctttggccag	gagagaacga	gtccagagcc	tgtcgaaaac	aagagaatca	gtgagcagaa	1320
ggtaagggaaa	ccgcaaccca	agaagagagc	tgcagtcggc	caatatttga	ggcagtcaca	1380
ctggaggagg	gcaggctctt	gaacaggctc	ctca			1440

<210> 12

<211> 2017

<212> DNA

<213> Chlamydia trachomatis

<400> 12

ataagcattc	tcattctacc	agaagtagaa	gtcaaaacct	tcataagtat	ctaaaaagac	60
tgcattataa	tcctcgatcc	catccggagg	cgctcctcgc	atccatattc	catggatgat	120
ttctccaaca	gtacacgat	ggccttttaa	ttctgttttg	atggtttcaa	gaacaccttc	180
aatccggagt	gtcttaggtt	ttctctgggc	ttctgtcttc	ttagcttttg	ctgttttagg	240
ctgagcctgc	gatgatcgtc	gaagcttctt	ctgaatggca	tcgacgtatt	ttcctgtgtg	300
aatcaaggaaa	ttctgtccgc	cttcgaattt	ttatctgggc	atagagttgt	aagcaataat	360
gacctttttt	agtttattta	ataggttctt	aacagtagat	ttctgttcag	gagtaattcc	420
tagttttttt	ttatgtttct	tgaggagtag	atcgtatttg	atcatcatca	gattttctat	480
ctttccagaa	gaagctttct	cctcttcttc	ttctatagca	cgcttttttc	togataaaaa	540
agctcgtcta	ggaggaactg	caatagcaga	aatcgttttt	cccccccccc	ctctgaaacgc	600
agtaagtaag	aacgttcaat	ggctgtgtaa	taaaactcgt	ctttctotta	cgagagagag	660
ttttgtcgtt	acttctcgtt	cttttagagat	tgtagtgcac	ttattctctcg	aagtgaaggt	720

ctctgcccgtc	tctgtccgaa	tctggccagca	gaagccatgt	tatcttttgt	tagatacaatg	780
cctcttttgtt	ttttgaattc	atcaagcatc	cagttgatga	tgactccgtc	gaagtgcgtct	840
cctcccaagt	gagtatcccc	gttgggttag	agaacttcaa	aaactccgtc	accgatttcc	900
aagatagaaa	tatcgaaggt	tctctctcct	aagtcgaaga	cggcgatttt	tttatctcct	960
tcttatcaca	taccataagc	aagagcgcc	gctgttggtt	caggaataat	gcgtttaaca	1020
tctaatctctg	cgatacgtcc	agcatctttt	gtagaagctc	tttgagaatc	gttaagatga	1080
gctgttacgg	taatgactgc	tccgttaact	gtttctcgca	gataagctc	cagtttaacc	1140
ttcatcttca	tgaggatctg	agcgcggatt	tctcttgagg	tgtagacgtt	ttgttccaca	1200
tcaaaagacg	catctctctt	cgagttagga	gcaactttgt	aggggactgt	tttaatttca	1260
gatttcgactt	cagagaattt	tctaccgatg	aatcgcttag	tagaagccaa	tgtttttttc	1320
ggatttggtta	tctcctgacg	ttttgcagga	attccacaaa	gagtttcgcc	acctttaaaa	1380
gcaacgatag	aaggagtagt	acgagttcct	tcaagaaggg	caataacttt	aggttggcca	1440
ccttccataa	cagagacgca	agagttggtc	gtccctagg	cgatacaca	aattttgtta	1500
gaactttctt	tttgcgtcat	altgaacacc	taattttctg	gataattatt	ctttttcttc	1560
gttacogtct	gagtttcttt	tagcaggaag	tttgctact	ttcacttlgg	ctacgcgaat	1620
aggaacgatct	cctatcttat	aaactttagt	aaattctctc	aagatagctc	cttctggaa	1680
tggttggtgtt	ttctcgattt	ctacacgttc	atgcaggtac	ggattaaata	gttctctctt	1740
cgagaaatat	tcaaccacac	ctttctcttc	gaagatttgc	ttaaatttgt	gaaggatcat	1800
ttggaatat	atagcccaat	ttttacttcc	tccagaggtt	tgagaagcga	atcccaaacg	1860
ctttttccata	cttttcgatag	aaggaaggaa	atccataaga	gcattttcta	cagcatactg	1920
catcatctct	gtcgtgtctt	tctgtagtcg	ttttcttgag	ttttctgctt	cagcgcagac	1980
cctcagatat	cgatcattct	gttcttgctc	cgtgcgg			2017

<210> 13

<211> 1171

<212> DNA

<213> Chlamydia trachomatis

<400> 13

ggtaaacagag	ttaaaacaag	agcatcacagg	gctaaccgag	tgcgctttag	tgaaaaaagc	60
tgaggagcag	attagtcga	caaaaaaaga	tattcaagag	atcaaaccta	gtgggtcggg	120
tattctctatc	gttggtccga	gtgggtccag	tgcttccgca	ggaagtgcgg	caggagcgtt	180
gaaatctctc	aacaattcag	gaagaatttc	ctgtgtgctt	gatgatgtag	acaatgaat	240
ggcagcgatt	gcactgcga	gtttctgac	tatgatcgaa	caattttaat	taaaccaatc	300
tgcaacagct	aaagagctac	aagctatgga	ggctcagctg	actgcgatgt	cgatcaact	360
ggttggtgcg	gtggcgagc	tccacgcgca	aatacaagca	atcaaatgat	cttcttgcga	420
agctttgcaa	caaccatcag	cagatgtgtt	ggctacagct	atgggacag	tggcttttgc	480
agctgcgaag	gttgagagag	gctccgcagg	aacagctggc	actgtccaga	tgaatgtaaa	540
acagctttac	aagacagcgt	tttcttcgac	ttcttccag	tcttatcgag	cagcatcttc	600
cgatggatatt	tctgtttaca	aaacactgaa	ctttttatat	tcgaaaagca	gaagcggcgt	660
gcagtcagct	attagtcaaa	ctgcacattc	cgcgctttcc	agaagcgttt	ctcgtctcgt	720
catagaaaat	caaggacgca	gtgcagatgc	tagcacaaga	gcagcagaaa	ctatttgcag	780
agatagccaa	acgttaggtg	atgtatatag	cgcgttacag	gttctgattt	cttttgatgc	840
tacgatttgtg	agcaattccg	aagcaaatca	agaagagatt	atgcagaagc	tcacggcatc	900
tattagcaaa	gttccacaat	ttgggtatcc	tgctgttccg	aattctgcgg	atagcttgca	960
gaagtttgtct	gcgcaattgg	aaagagagtt	tggtgatggg	gaacgtagtc	tcgcagaatc	1020
tcaagagaat	gcgtttagaa	aacagccgcg	tttcatccaa	caggtgttgg	taaacattgc	1080
ttctctattc	tctggttacc	tttcttaaac	tggtgatgaa	gtttgtgaat	gagggggagc	1140
caaaaaagaa	tttctttttt	ggctcttttt	t			1171

<210> 14

<211> 877

<212> DNA

<213> Chlamydia trachomatis

<400> 14

cagagaatttc	togacatact	atctaactcg	atatgtaaag	ctgctttaca	tcccttgaa	60
tagaaaataa	atggaataaa	aaagcccaag	acaaagagag	ttgttctcgg	ctgcagaga	120
ctgtcagatc	attttataaa	gattgatgac	aactacgaca	agttctcggg	tccaaaagag	180
aatctaaaaa	gcctacataaa	gattgogtta	ctttcttgca	tgctctaac	cttttaaatc	240
gctcatcttt	gagaagcatt	tcaatgagcg	ctttttcttc	tctagatcgc	cgcacatccg	300

8

```

ctttctcatg tttctgtaa tatgcatagt cttcaggatt ggaaatcca aagtactcag 360
tcaatccacg aattttctct ctacgcgatac gtggaaattg actctcataa gaatacaaaag 420
cagccactcc tgcagctaaa gaattctctg tacaccaccc cagcaaaagta gctactttcg 480
ctttttctgc ttcaactaggc tcatgagcct ctaactcttc tggagtaact cctagagcaa 540
acacaaactg cttccacaaa tcaatatgat tagggtaaac gttctcttca tccatacaag 600
tatctacaaa taacttaccg gctcttaaat catcgcaacg actatgaatc gcagataaaat 660
atttaggaaa ggcctttgata tgttaataat agtctttggc atacgctctg aattgctctt 720
tagtaagctc ccccttgacg catttcacat aaaacgtgtg ttctagcata tgcttatttt 780
gaataattaa atctaactga tctaaaaaat tcaataaac cctccatcatt tctttttctg 840
actccacgta accgcttgca aaaaaggtcc gtataaag 877

```

<210> 15

<211> 396

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 15

```

tgtaccaaat atgagcttag atcaatctgt tgttgaactt tacacagata ctgctctctc 60
ttggagcgtg ggogctogag cagcttttgg ggagtcgga tgtgcgaact taggggcttc 120
tttccaatcc gctcaatcta aacctaaagt cgaagaatta aacgttctct gtaacgcagc 180
tgagtttact atcaataagc cttaaaggata tgtagggcaa gaattccctc tgcactctat 240
agcagggaact gatgcagcga cgggcaactaa agatgcctct attgattacc atgagtggca 300
agcaagtgtt gcctctctct acagattgaa tatgttcaact cctcaacttg gagttaaagt 360
gtctcgacga agttttgatg cggatacagat tegtat 396

```

<210> 16

<211> 516

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 16

```

ctcaaaattt gaagatttct cagaatacag ggaatgttct gttttataac aacgtggcct 60
ctctggggagg agctgttctg atagaggatc atggtaattg tcttttagaa gcttttggag 120
gagatattgt ttttaaaggaa aattcttctt tcagagcaca aggatccgat gccactatt 180
ttgcaggtaa agaatcgcat attacagccc tgaatgctac ggaaggacat gctattgttt 240
tccacgcgagc attagttttt gaaaatctag aagaaggaaa atctgctgaa gatttggtaa 300
tcaatagctg agaaaaatcca ggttacactg gatctattcg atttttagaa gcagaaagta 360
aagttctcca atgtattcat gtacaacaag gaagccttga gttgctaaat ggagctacat 420
tatgtagtta tgggttttaa caagatgctg gagctaagtt ggtattggct totggatcta 480
aactgaagat tttagattca ggaactcctg tacaag 516

```

<210> 17

<211> 723

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 17

```

ctccttttaa gggggacgat gtttaactga atggagactg cgtttttgtc aatgtctatg 60
caggggcaga gaacggccta attatctcag ctaatggcga caatttaacg attaccggac 120
aaaaccatac attatcattt acagattctc aagggccagt tcttcaaaat tatgccttca 180
tttccagcag agagacactt actotgaaag atttttcogt ttgtatgttc tcaaaaaatg 240
tttcttgccg agaaaaaggga atgatctcag ggaaaaacgt gagtatttcg ggagcaggcg 300
aagtgatatt ttgggataac tctgtggggt attctctttt gtctattgtg ccagcatcga 360
ctccaactcc tccagcacca gcaccagctc ctgctgtctc aagctcttta tctcaaacag 420
ttagtgtgct tgggaaaggg tctatttttt ctgtagagac tagtttggag atctcaggcg 480
tcaaaaaagg ggtcatgttc gataataatg ccgggaattt tggaaacagt ttctcaggta 540
atagtaataa taatgctggt agtgggggta gtgggtctgc tacaacacca agttttacag 600
ttaaaaaactg taaagggaaa gtttctttca cagataacgt agctctctg ggagcggcag 660
tagtctacaa aggaactgtg cttttcaaa acaatgaagg aggcattatc ttccgaggga 720
aca 723

```

<210> 18
 <211> 1377
 <212> DNA
 <213> *Chlamydia trachomatis* serovar E

```

<400> 18
aaacagcttaa tcgtcactac gctcacgtgg actgcctctg tcacgctgac tatgttataaa 60
acatgatcac cggtgcggct caaatggagc gggctattct agtagtttct gcaacagacag 120
gagcatatgcc tcaaaactaaa gagcatattc ttttggcaag acaagittggg gttccttaca 180
tcgtgttttt tctcaataaaa attgacatga tttccgaaga agacgctgaa ttggtcgact 240
tgggtgagat ggagttggct gagcttcttg aagagaaagg atacaaaggg tgtccaata 300
tcagaggttc tgccttgaaa gotttggaa gggatgctgc atacatagag aaagttcgag 360
agctaatgca agcctgcgat gataatatoc ctactccaga aagagaaatt gacaagcctt 420
tcttaatgcc tattgaggac gtgttctcta tctccggagc aggaactgta gtaactggac 480
gtattgagcg tggattgttt aaagtttccg ataaagttoa gttgtgtggt cttagagata 540
ctaaagaaac gattgttact ggggttgaaa tgttcagaaa agaactccca gaaggtcgtg 600
caggagagaa cgttggattg ctctccagag gtattggtta gaacgatgtg gaagaggaa 660
tgggtgtttt cttgccaaac agtggttaaac ctacatacaa gtttaagtgt gctgtttacg 720
ttctgcataa agaagaaggt ggaacacata agcctttctt cacaggatat agactccaat 780
tcttctccgg tacaaacagc gttacagggt tggtaactct gctgaggaga gttgagatgg 840
tcctgcctgg ggataacggt gagtttgaa gtcgaattgt tagcctctgt gottttagaa 900
aaggtatgag atttgogatt cgtgaagggt gtctacaatt cgtgtcgtga actatttcta 960
agatcactgc ataaattaa gttgtgtgtt gogaggctga aaagccttgc ctttgggtgt 1020
gtagctttaga ttgtagagca gtggcctcca aagccgcggg tgcggggtgc gaatccctcc 1080
gcactcgtat tagtaaacgt aaagaagaat tcgcttatgg ggcgaagtca ccgaagaaa 1140
tttcttaaga aagtatcttt tgcataaaaa caagcagctt ttcgggtaa ctttatcgaa 1200
gaaattaaag agattgagtg ggttaataag cgaacttta aaagatacgt caagatttgt 1260
ttgatgaata ttttggctt tgaatttccc atctattgtg tggatttagc tctctgaag 1320
tcctttctat tgttcggtta agtaacaagc ttttctttg tctgattcat gtttaag 1377

```

<210> 19
 <211> 1736
 <212> DNA
 <213> *Chlamydia trachomatis* serovar E

```

<400> 19
gtagcggaac aaagccggac caccaggcct catagaatat aaaaatacga ggagctttaa 60
catgtccgat caagcaacga cctccaagat taaccttttg ggagatagaa ttttagttaa 120
aagagaagaa gaagcttcca ctgcaagagg cggatcattt ctctctgaca ctgccaagaa 180
aaagcaagat agagctgaag ttttagctct aggaacagcg aaaaagatgt atcaagagca 240
gcaactctct tttgaagttc aggttgggtga catcgtttta atgtataaat attctggcca 300
agaactcact ttgaagagtg aagagtaagt catcgttcaa atgacgcaag tatcgcagct 360
tctgcaataa aaactaagag agtgaagtaa gatttaaagg agcgatcaa tggctgctaa 420
aaacattaaa tacaacgaag aagccagaaa gaaaattcaa aaaggagtta agactttagc 480
tgaagctgta aagtcactc tagggcctaa aggaagacat gttgtcatag ataaaagatt 540
cggtaccctt caagtaacta aagatggtgt taccgttgcg aaagaagtgt agcttgcgca 600
caaacatgaa aatattggcg ctcaaatggt caaagaagtc gccagcaaaa ctgctgacaa 660
agctggagac ggaactacaa cagctactgt tcttgcgtga gctatctata cagaaggatt 720
acgcaatgta acagctggag caaatccaat ggacctcaaa cgaggtattg ataaagctgt 780
taaggttgtt gttgatcaaa taagaaaaat cagcaaaact gttcagcatc gttcagcatt 840
tgctcaagtt gcaacaattt ctgctaataa tgatgcagaa atcgggaatc tgattgtgta 900
agcaatggag aagatttgga aaaaacggctc tatcactgtt gaagaagcaa aagatttga 960
aaccggtttg gatgttgttg aaggaatgaa tttcaataga ggttacctct ctgactactt 1020
cgcaacaaat ccagaaactc aagaattgtg attagaagac gctttggttc taactcacga 1080
taagaaaatt tctgggagca aagatttctt tctgttttta caacaaagtg ctgaactcgg 1140
ccgtctcttt ttattatag cagaagacat tgaaggcgaa gctttagcta ctttggctgt 1200
gaacagaatt cgtggagagt tccgggtttg cgcagttaaa gctccaggct ttggagatag 1260
aagaaaagct atgttggaag acatcgtcat cttaactggc gctcaactca tttaggaaga 1320
gttgggcatg aaattagaaa acgctaactt agctatggtt ggtaagacta aaaaagatt 1380
cgtttctaa ggaacacga ccaatcgtga aggaatgggt gaaaaagaag ctttagaac 1440
tcgttgcgaa agcatcaaaa aacaaattga agacagctct tctgattacg ataaagaaa 1500

```

10

```

actccaagag cgtcttgcta agctctctgg tggagtagca gtcattcgcg ttggagctgc 1560
aacagagatt gagatgaaga agaaaaaaga tcgtgtagac gatgctcaac atgctacaat 1620
cgctgctggt gaagaaggaa ttcttcctgg tggaggaa caattaatcc ttgttatccc 1680
tactcttgag gctctcttgc caatgtttgac taatgaagat gagcaaatg gagctc 1736

```

<210> 20

<211> 1135

<212> DNA

<213> *Chlamydia trachomatis* serovar E

<400> 20

```

ggctcttgat gaaaaagagc ggcaggttat ggctctttat tactatgatg acttgggtatt 60
aaaaagaaatt gggaagattt taggagtgag cgagtcocga gtttctcaga tacactccaa 120
agcttttattg aagttacagag gtacattgtc cagtcctgct tagtaacctg ctccagaaga 180
tcctctttgtt atttttctta tcaatattct attggagaag cgcgtcggtt ttttgacagag 240
gtgtctgctga tcgcttgctc tgcataaaaa agaacaggat agataagatg ttgotagata 300
agtttataatg gatagatttt tatgcaacag ttaatcgata accttaagaa acgggggtatt 360
ctagataaatt cttctgcagg attagaaact cgtgccgaag tttgtggaga agagaaagaa 420
atctctctag cagactttcg tggtaagtat gtatgtctct tcttttatcc taagaatttc 480
acctatgtgt gctctacaga attgcatgct tttcaagata gattggtaga ttttgaagag 540
cggggtgcag tcgtgcttgg ttgctccgtt gacgacattg agacacatto tcgttggctc 600
cgtgtagcga gaaatgcagg aggaatagag ggaacagaa atctctctgt agcagacctc 660
tcttttaaaa tatcagaagc ttttgggtgt ttgaatcctg aaggatcgct cgtcttaaga 720
gogactttcc ttatcgataa acatgggggtt gtctgtoatg cggttatcaa tgatctctct 780
ttaggcggtt ccattgacga ggaattgcgt atttttagat cattgatctt ctttgagaa 840
caoggaattg ttgtccagc taactggcgt tctggagagc gtgggaatgt gcttctcaga 900
gagggtattaa aagaatattt ccagacagatg gattaagcat ctttgaagat aagaagatgc 960
tacaagattct gatctgaaaa gagaagaagg ctttttaatt ttctgcagag agccagcgag 1020
tgctcaattaa tgttgaagtc tccgccacca ggcaatgcta agcgcatgat attgatgat 1080
gaaatctgag tgtttaaggaa ataaaggcca aagaagtagc tatcaataaa gaagc 1135

```

<210> 21

<211> 731

<212> DNA

<213> *Chlamydia trachomatis* serovar E

<400> 21

```

ttgaagacac tctttctccc ggagtcacag ttcttgaagc tgcaggagct caaatcttct 60
gtaataaagt agtttggact gtgaaagac tgaatcctgg agagtcctcta cagtataaag 120
ttctagttaag agcacaaact ctgggcaact tacaataaaa tgttggtttg aagagctgct 180
ctgactgtggt tacttctgact tcttgcgcag aagcgcaaac tactgtggaa ggagttgctg 240
ctctcataat gtgcgtagta gatacttctg acctgttttg tgtaggagaa aatactgttt 300
acogtatttt tgcaccaaac agaggtttctg cagaagatac aaatgtttct ttaatgcta 360
aatctctctaa agaactgcac cctgtatcct tctctggacc aactaaagga acgattacag 420
gcgaatacagt agtattcgat tctgtaccota gattagggttc taaagaacct gttagatttt 480
ctgttaacatt gaaagcagtt acagctggag atgctcgtgg ggaagcgatt cttctctcog 540
atacatctgac gtcttcagtt tctgatacac agaatacaca catctataa tcttttgatt 600
tactogattg taggtgcggt ccagggtatc ctggcgctgt ttttttgtt ctctataga 660
aaataaaaga gttcatcttc ggtctcagag catattctag acgggttttt gaaaaaata 720
agtgtttgtg t

```

<210> 22

<211> 1181

<212> DNA

<213> *Chlamydia trachomatis* serovar E

<400> 22

```

ctatcgtctg aatgctgaac tgaacaatct ttttgattta gacgcgttag cogatgctat 60
ggatctatct cgagatctac agtttttcta catgggtatt caaaatctgt atgatcgcta 120
ttttatacac cagaagaatt gcggtttaga aactcccaaa attttttgga tgcgctgtgt 180
tatgggggtg gcaatgaatg agcaagacaa gactctcttg gctattactt tttataatt 240

```



```

gctttogaca ttcgatata caccagctac gccaaccttg ttcaattcag gtatgcggca 300
ttctcagtta agctcttgct atctttccac tgtacaagat aatttggtea atactataaa 360
gggtcattgct gataacgcta tgcattctaa gtgggcaggga gggataggta atgattggac 420
ggcggttctgt gcaacaggggg othtaattaa aggaaccaat ggaagaagtc agggagtaat 480
tccttttatt aaggtgacaa atgatacagc agtcgcagtg aatcaaggtg gttaacgcga 540
gggagctgta tgcgtctatt tagaagtttg gcacctcgac tacgaagatt tccctgaatt 600
gagaagaagt acaggggagtg agcgtcgagc ggcctcatgat gtcaatatag ctactgtgag 660
tccagatctt ttcttcaaac gtttacagca aaaagggaca tggactctat tcagcccgaa 720
tggagagaaat ggattacacg atgcttatgt ggaagaattt gagcgtttgt acgaagaata 780
tgaagcggag gttgatcacg gagagattog gttattcaag aaggtagaag ctgaagatct 840
gtggagagaaa atgctcagca tgccttttga aacgggacac ccatggatga cttttaaaga 900
tccatccaac atccgttcggt ctcaagatga taaaggcggt gtgcgtttgt ccaatctgtg 960
tacggagatt ttgttaaact gctcggagac agaaactgct gtttgtaatt taggatcgat 1020
taacttagtt caacatatcg taggggatgg gttagatgag gaaaaactct ctgagacgat 1080
ctctatagca gtcogtatgt tggataacgt gattgatatt aacttttacc caacaaagga 1140
agctaaagag gcgaactttg ctccacgcgc tattggatta g 1181

```

<210> 23

<211> 167

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 23

```

ttaaaaagat tttaaactaa aaagaagatt ttttaattata gtttttcaaa atcattttga 60
tatttttaat gctgagataa acaagaaaag cggaaactcc ttggacaaaa gattttctgc 120
tcgagccctc ttccctgagg atttttttag ggagatccat tcttcca 167

```

<210> 24

<211> 1265

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 24

```

cagggttcttt ctagacgaac aaagaataat cctatgttga taggggagcc cggagttggg 60
aaaaacagcaa tgcgtgaagg acttgccttt cgcatagtgc aaggggatgt tccagagagt 120
ttaaagaaaa agcatctgta tgtactggat atgggagctt tgattgcagg tgccaagtat 180
cgagaggaagt ttgaagagcg gttaaaaagt gtattgaagg gtgtagaagc ttctgaagcc 240
gagttgtatcc tattcattga tgaagtgcac acttttagtag gagcgggagc tacagatgga 300
gctatggatg cagcgaatct attaaagcct gotttagcac gaggcacttt gcatttgaatt 360
ggcgctacga ctttgaatga ataccaaaaa tatatagaga aagacgcggc ttgtgaacgg 420
cgtttccagc ctatttttgt aacagaaact tctttggagc atgctgtatt caattctccg 480
gggttaaggg aaaaaatgta aatttttcat gggtgcccga ctacagaagg ggctttgaat 540
gcagctgttag ttcttttcta tegtatcacat acagaccgat ttcttctgta taaggcgatt 600
gacctaaatt atgagcgctg gagtttaact cgtatgcaaa taggaagttt accctgcctc 660
attgatgaaa aggaagagga attatcagct ttaatcgtga aacagaagc tattaaagc 720
gagcaagcac cagcttatca ggaagaggct gaagacatgc aaaaagcaat accctgcctc 780
aaggaagagc tggccgcttt acgcttgcgc tgggatgaag aaaaagatt aatgcagga 840
ttaaaaagaa aagaagaatgc tttagaaaaa ttaaaatttg ccgaagagga agctgcagct 900
actgcgattt acaatcgggt agcagaacta cgotatagtt tgattccttc ttgtggaggaa 960
gaaattcatt tagctgagga agctttaaat caaagagatg ggccgctgct tcaagaggaa 1020
gttgatgagc ggttgatgac gcaagttgtt gcgaattgga ctgggaatcc ttgtcaaaaa 1080
atgttggagg gagaactgta aaagtatttg gtgttgagga gtctttagaa gaaagggttg 1140
tcggacagcc tttcgctatt gccgcagcta gtgattcgat tcgagctgct cgagttagat 1200
tgagtgatcc gcagcgtctc cctcacaagg gaattattagc tggcggcgcc aaccgctggc 1260
gaaac 1265

```

<210> 25

<211> 463

<212> DNA

<213> Chlamydia trachomatis serovar E

```

<400> 25
atgacgaaca accccatggt tctcgataag gaaagtcgct cttaaagcga gcgatccctc 60
aggattcaaa accacaaaag ottotgatat tttaaaagaa gggctctgta acagaggata 120
ttctgtccoc tctattcttc ctgcattttct cgtacacgag agccaacgag aatgtgtctc 180
aatgtctgta acggagcaac caagcacgac tgcaccccg ctttcaaaat tcccaaatct 240
atcttgaaaa gcatgcaatt ctgtaggaca cactagagtg aaattcttta gataaaaaga 300
gcgactaca tacttaccac gaaagtctgc tagagagatt ttttctcttc ctccacaaa 360
aacgctgtta ccagaaaaat ccggagcctg ttttccaatt agtgatocca taatactcct 420
octagaaga aacaacgcac cagagaggat ttgaacctot gac 463

```

```

<210> 26
<211> 636
<212> DNA
<213> Chlamydia trachomatis serovar E

```

```

<400> 26
ggtagaaat tctctgaagt cgaatctgaa attaaaacag tccctacaa agttgtctct 60
aactcgaag gagatgcggt ctttgatgtg gaacaaaac tgtacactcc agaagaaatc 120
ggcgctcaga tctcatgaa gatgaaggaa actgctgagg ctctatctcg gaaacagta 180
acggaagcag tctattaccgt accagcttac tttaacgatt ctcaagagc tcttcaaaaa 240
gatctggagc gtatcgcagg attagatggt aaacgcatta tctctgaac aacagcgccg 300
gctcttgctt atggtattga taaggaagga gataaaaaa tcccgctctt cgcacttaga 360
ggaggaactt tgcgatttct tatottggaa atcgggtgag gaggttttga agttctctca 420
accaacgggg atactcactt gggaggagac gaactcgaag gacttcacat caactggatg 480
cttgatgaat tcaaaaaaca agaagccatt gatctaagca aagataacat ggctttgcaa 540
agattgaaag atgctgctga aaaagcaaaa atagaattgt ctggtgtatc gttacttgaa 600
atcaatcagc cattcatcac tatcgacgct aatgga 636

```

```

<210> 27
<211> 1797
<212> DNA
<213> Chlamydia trachomatis sere

```

```

<400> 27
atgcctacac atcaccatca catgagcctc aggggagtag gaggcaacgg gaatagtcca 60
atccctctct ataatgggga tggatcgaat cgcgaagalc aaaaacgaa gggtataaat 120
aaagtgtgaag atcgagtttg ttctctatat tcatctcgta gtaacgaaaa tagagaatct 180
cttatgacag tagtagacgt cagctctatg atcgagcaga ccccaacgag tggagagcac 240
acaaagtcgt ccgctggagt gctcagctcag ttccaagag gtttagtacc aatagctgac 300
aaagtgaagc gacgtgttca gtgtgcgtgg agtcaagctc ctacaagcag atcgctctga 360
acaagagccg cagaatccgg atcaagtagt cgtactgctc ttggtgcaag ttctgggata 420
aggagatatt ctctctcagc agctagaggg ctgcgtctta tgttcaaga ttctgggata 480
actcgggttt tacgcacagc ctctcctatg cgtggagttt ttgggaatct tgatctgac 540
gaggtcgtgt tgatggtctg gtacacaaat gagtgcggcg atcatttaga agcgaagag 600
ttggctggcc ctgacggggt agcggccgcc cgggaaattg ctaaaagatg ggagaaagaa 660
gttagagact tacaagataa aggtgtgcga cgaataatt taaatgatcc tttaggcga 720
cgacaaccta attatcacag caaaaaatcca ggtgagtata ctgtagggaa tccactgtt 780
tacgatgtct ctacagtagc gaattctccag aacgtgcaga ctggtttttg gctggacgt 840
agcaatctct cagaagttgt attatccaga gagattcaca caggacttcg agcaagact 900
actttggaag aatccatgcc gatgttagag aatttagaag acggttttag acgtttgcaa 960
gaaacttgtg atcggcctcg tactgagata gaagaatcgg gatgactcg agagtccga 1020
tcaagaattg aagcgcatga ggcgcgaaga cttctagata tacaacaagc ttttcagaga 1080
tttgtaaatg aatgtaacag catcgagttc tcaattggga cgtttggaga gcatgtgcca 1140
ttctctctgc ctgagtagtc acgagtagta gctgcgcgag gagaggcgat tccgctgtgc 1200
ttctcttggt gtaaaagatc cagcgcctgc tactctcttc cgttagactc atctctgaa 1260
ggtgcactgt tagcagagac tttggctaga ttgcagagat atatgggaat agagcaggt 1320
gctgatggaa ctacagatat tcttttgta gatgattgga gaagaggggt tctcatgtat 1380
gaaggagaag gatctgactc gatctatgaa atcatgatgc ctatctatga agttatgaat 1440
atgatatctg aaacacgaag atcttttgcc gtacagcaag gcacactaca ggaccaaga 1500
gcttcagatt atgacctccc acgtgctagc gactatgatt tgcctagaag cccatctcct 1560

```

13

```

actocaccctt  tgcctcotag  atatcagcta  cagaatatgg  atgtagaagc  aggggtccgt  1620
gaggcagtttt  atgctctottt  tgtagcagga  atgtacaatt  atgtagtgc  acagcgccaa  1680
gagcgtatttc  coaatagtc  gcaggtggaa  gggattctgc  gtgatgtct  taccacgggg  1740
tcacagacat  tttagaacct  gatgaagcgt  tggatagag  aagtcgatg  ggaataa  1797

```

<210> 28

<211> 1983

<212> DNA

<213> Chlamydia trachomatis serE

<400> 28

```

atgoatcacc  atcaccatca  catggaatca  ggaccagaat  cagttttctc  taatcagagc  60
tcgatgaatc  caattattaa  tgggcaaatc  gcttctaatt  cggagaccaa  agagtccacg  120
aaggcgctcc  aagcgagttc  ttcagcatcg  tctctgttaa  gcagctggag  ttttttatcc  180
tcagcaaga  atgcattaat  ctctcttcgt  gatgccatct  tgaataaaaa  ttccagtcac  240
acagactctc  tctctcaatt  agaggctct  acttctacct  ctacgggtac  acgtgtagcg  300
gcaaaagatt  atgatgagc  taaatcgaa  tttgatacgg  cgaaaagtgg  attagagaac  360
ctaaagacac  ttgctgaata  cgaacgaaa  atggctgatt  tgatggcagc  tctccaagat  420
atggagcggt  tagctaattc  agatcctagt  aacaatoata  ccgaagaagt  aaataatatt  480
aagaatcgcc  togaagcaca  aaaagatact  attgataagc  tgaataaact  cgttacgctg  540
caaaatcaga  ataaattctt  aacagaaagt  ttgaaaacaa  ctgactctgc  agatcagatt  600
ccagcgatta  atagtcagtt  agagatcaac  aaaaattctg  cagatcaaat  tatcaaaagt  660
ctggaaagac  aaacataag  ttatgaagct  ttgtctcata  acgcaggaga  gtttatccaa  720
gctctctctg  aagcgggaat  taagttagga  caagctttgc  agtctattgt  ggatgtcggg  780
gaccaaagtc  agcgtgcagt  tctgcagca  cagcaaaaata  atagcccaga  taattatgca  840
gcocaggaag  aattaattga  tgctgtgtaa  acgaaggtaa  acagattaaa  acaagagcat  900
acagggctaa  cggactcgcc  tttagtgaag  aaagctgaag  agcagattag  tcaagcacaa  960
aaagatattc  aagagataca  aactagtgt  toggattatc  ctatcgttgg  tccgagttgg  1020
tcagctgtctt  cgcaggaag  tgcgcagga  gcgttgaaat  ctctatacaa  ttccaggaag  1080
atttctctgt  tgctgtatga  ttagacaaat  gaaatggcag  cgattgcaat  gcaaggtttt  1140
cgatctatga  tgaacaatt  taatgtaaac  aatctgcac  cagctaaaga  gctacaagct  1200
atggagggctc  agctgactgc  gatgtcagat  caactgggtg  gtgggagtg  cgagctccca  1260
gcgcgaatac  aagcaatcaa  agatgtctct  gcgcgaagct  tgaaacaacc  atcagagat  1320
ggtttggcta  cagctatggg  caaagtggt  ttgtcagctg  ccaaggttgg  aggagctctc  1380
gcaggaacag  ctggcaactgt  ccaagtgaa  gttaaacagc  gttctcagtg  aggtttttct  1440
tcgactctct  caagctctta  tgcagcagca  ottctcagtg  gatattctgc  ttacaaaaaa  1500
ctgaactctt  tatattcaga  aagcagaagc  ggcgtcagct  caagtattag  tcaaaactgca  1560
aatcccgctg  ttccagaag  ogttctcgt  tctggcatag  aaagtcgaag  agcagatgca  1620
gatgtcagcc  aaagagcagc  agaaaactatt  gtcagagata  gccaaacgtt  agggatgta  1680
tatagccgct  tacaggttct  ggattctttg  atgtctacga  ttgtgagcaa  tccgcaagca  1740
aatccagaag  agattatgca  gaagctcaag  gcatctatta  gcaagctcc  acaattggg  1800
tatcagtgct  ttcaaatc  tgcggatag  ttgcagaagt  ttgctgcgt  ttgtgaaaga  1860
gagtttgttg  atggggaacg  tagtctcgca  gaattccaag  agaatcgctt  tagaaaacag  1920
cccgtcttca  ttcaacaggt  gttggtaaac  attgctctc  tattctctg  ttatctttct  1980
taa  1983

```

<210> 29

<211> 1224

<212> DNA

<213> Chlamydia trachomatis serE

<400> 29

```

gtaacttttc  aacatttttc  acaatgacaa  gaataaaagc  aaaaagaag  gctgccgata  60
aaataaaagt  tttactcgca  gaacagaaga  ctaaaactat  ctggacgaat  aagccgagat  120
cgagagataa  ttgcgcataa  aacactttaa  tagagagtga  tttattgtct  aaacacacct  180
tatccatagc  taactctctg  catgggocag  tatataacgc  cgatgattat  gaactctttt  240
gttgctatat  agaataoact  ccagcgagct  cogttaaatt  cgaaactggat  aaagaaactg  300
gtatctcaaa  atgtgactcg  ccacaaaagt  ttttaacttt  ttgtctcttg  ttatagctcg  360
tgttaoctaa  gacttattgt  ggagatcttt  ctggagaata  cagtgggtcaa  caaagtaaca  420
gagagataat  caaagcgat  ggcgatctct  ttgatatctg  ttgtttaacg  aaaaaaata  480
ttacacaag  gaacatctct  ttgcaagcgc  gtctatctcg  agggattctg  attttagact  540

```

```

cggaagaagc  cogatgataa  atcatogctg  ttctagaaga  tgatttagtc  tatggcaata  600
taagaagatat  ttctgaatgc  ccaggcacag  ttctggacat  gatccaaac  tatttcttaa  660
cotataaagc  tactccagaa  agcttaattc  aagcaaaacc  agctaaaatt  gaaattgtag  720
gtttatacgg  caaaaaagaa  gctcaaaaag  tcatttgcgt  tgctccagaa  gactattgca  780
atctttttat  gtaaatogac  agaaaaagaa  aaggctgttg  tgggagattc  cacaacggcc  840
ctctctaac  aagttttttt  catctagggg  gactttatga  agcaaataga  taactttgaa  900
caaattcatc  tctogtgcgc  aattoggcac  gagattaaaa  caaagctoto  aaaaagatt  960
ggatcccgca  attcatcag  cagttccggc  tgccaaagtt  aaagagatac  gctttttatt  1020
aggaatgta  tggacgcaca  agaaaagaaa  taagacgcac  cagcoatcac  cgtttttaga  1080
ggattgcag  ctgttgctga  ggtctctgga  atgtacattg  gtgatacagg  agttacccga  1140
ttgatccact  tggtttatga  agtgggtgat  aacagtatgc  atgaggcaat  ggcgggtttt  1200
tgtaccgagg  tegtgtgttg  cata  1224

```

```

<210> 30
<211> 883
<212> DNA
<213> Chlamydia trachomatis serE

```

```

<400> 30
atgttgacta  acatggcgac  catcagaaac  totgtgaaga  cattgaacag  aattgaattg  60
gatcttgaag  cttctaatc  ttgtcttacg  aaaaagaga  tcgctttatt  aacgaaaaga  120
cgcgcgaact  tgcataca  cctgggaagt  gttctgta  tgaactctct  cccagggttt  180
ttaattgtaa  ttgacccggg  ctatgagcgc  attgctgtgc  cagaagctgg  aaaaataggc  240
attctgttaa  tggccttagt  tgatacaaac  tgcgatccaa  caccatcaaa  ccacgttatt  300
ccttgcaaac  atgattccat  taagagatgc  cgtctgggtg  tcaatgtact  taagaacgct  360
gttatgtgat  cgaagaagcg  ttoaggcacc  gaaattttat  ctccagtaag  tctgtgaa  420
agacotgcag  aagaagctgt  ggaagagttg  cctcttccaa  caggtgaagc  tcaagatgaa  480
gtcttotcta  aagaaggttt  tttactttgg  gcagatattg  acaattgagg  ggcattgaa  540
tgagcgaact  ctccatggaa  acattgaaaa  atttaagaca  gcagacaggt  tgaggcctga  600
ctaaatgtaa  agaggtctta  gagcatgcta  agggcaattt  agagatgct  gttgtttatt  660
taagtgaact  tggctttggc  ctgcaggcca  aaaaagaaca  cggagaaaca  aagaagcggt  720
taattgctgc  actogttgat  gaacgtgggt  cggcacttgt  tgaagtcaac  gttgaaactg  780
attttgttgc  taacaacagt  gtttccgag  caitcgltac  aggtttgita  tccatgtctc  840
tgtaccacaa  gcttagcgat  gttgaagctt  tagctoggt  aat  883

```

```

<210> 31
<211> 393
<212> DNA
<213> Chlamydia trachomatis serE

```

```

<400> 31
agtgtaaaaa  ggotgtttct  tgcattcaaa  aaactatcga  gcaagagaga  tctattttgt  60
ttgttgtaac  aaaaaaacag  gcaaaacaga  tcattagaga  agctgtatc  gaaatgtggc  120
aatcttttgc  ttoagagaga  tggttgggtg  gcaatgtgac  taacatggcg  accatcagaa  180
actctgtgaa  gacattgaac  agaattgaat  tggatcttga  agctcttaat  tctgtgttta  240
cgaaaaaaga  gatgccttta  ttaacgaaaa  gacatcgcaa  gttgttcaa  aactgtgaa  300
gtgtgttcca  tatgaactct  ctccagggcg  ttttaattgt  aattgaccgc  ggcataagac  360
gcattgtcgt  cgcagaagct  ggaaaactag  gca  393

```

```

<210> 32
<211> 2577
<212> DNA
<213> Chlamydia trachomatis serE

```

```

<400> 32
attacggagg  ccatacggta  tcttctcgag  gaggatttca  agggatatgc  gtacgaatag  60
ccgatttatt  ccgtaactgt  ttctctcgta  atagaggcac  tactactacg  ccactctgaa  120
ctgttatcac  taaggcagat  atttatcacc  cgaatatttc  tggacaagga  gctcaacct  180
ttgtctctac  aggaagataa  aaattagata  gcgcaattat  tcaagcagat  ttgctgtggc  240
agaaataaca  gactttggct  acacatatcc  aaagtaagct  aggttctatg  gaggacacat  300
ctctcaaga  ttataaagct  ggtgcgtata  gtgcgctaag  attgatgctg  ttactccag  360

```

```

gcgaactac  tgtgagtagc  gagcgggaac  gtcaagcggt  cgttacgggt  cgggatctct  420
gggaacagcg  tgcaggagat  cttgctacca  atgggaatac  agatggggct  atgttaatag  480
ctaactatc  tgtggggagg  aagcatgtgc  ctgcggggca  ttaagagaa  tacatggata  540
ctgtaaagg  tacgtttact  gatgagaacg  aggtacagaa  tctacaggta  gatgccattt  600
tagatttagc  agcaaaaatc  gatcgacgg  aattctctag  tcttggttca  gggaagatca  660
ttcttaatta  tataggaaat  tatggacaag  tctgttttag  aaacgaggag  atgaaccttc  720
ttgttttaga  agatcaaaat  ggcaagatc  ctcaacgtgt  tcaagataac  tcaaaagagt  780
tacaanaact  gttagaaaat  gctcgaanaa  cagatcctga  gttatatct  caaacactaa  840
ctgtcataac  ttctctgtgt  ttcttagact  aaagagaagg  tatcgggtgt  tcggttcott  900
caactattaa  gaggaagtag  tggtagtagt  cataagccct  atagggggga  attctgggcc  960
agagggattt  tctagtgcac  ctcgaggcga  tgagattgat  gatgtaccag  atagtattga  1020
gggagagcta  gaagagcgcg  ttctggatca  tgcagagtct  atcattacgc  agagctcgga  1080
aacgtgtttt  cgtactactt  cttcatcagg  ggtcagtgaa  gatcttcagc  aacacgttag  1140
cttggaggaa  ttccacgac  aacgaggttt  ccttgacgg  atccgtgatg  cagttagctc  1200
tatttggaa  cgtcgtgttg  cacgaaggaa  tgaaaactat  gatgtgaaaa  aagcagaaga  1260
cgacgaagg  attgtgcaat  atctgcagga  ttcgaaaatg  cctgctttaa  cgcgtgccta  1320
tcgccatctc  cgtgctttca  attctgcatg  cttacgtacg  attcgtgagt  ttttgcctac  1380
catttttctg  gataaaagg  atgcttatta  tcgacattgt  acacgttctg  ggatcaactt  1440
tttggagct  gataaagact  ctttagaagt  ctttgttgog  tggggtttgc  gttgttgatt  1500
ggotacotta  cgtcttttgc  aacatgtcgg  tgggaattac  gaagatcgat  tagtaataaa  1560
tgatgtccog  tgcacagggt  cggggagaac  tctgttgtag  gatgctgtag  acgatattga  1620
atcgatttta  aatcagagaa  ccaactggcc  tcaacatgct  atgatagggg  tttctcgtag  1680
ttctgttcaa  ttatgtgcga  ctcttataaa  tgcgactctc  caagaatggt  tcaagtogat  1740
tggtgtttaa  gaaaaaagg  acccttcttc  agattattct  caagctttat  ttatagcagg  1800
gataatagat  cgtctggcgg  agaaagccoc  tatggctgca  aagtattgtt  ttggtcaaat  1860
gcgtgttcga  actctggagc  tcataggaga  actcaattat  ctogattttg  ttctcctgtg  1920
atgggaaggt  ggcggggagg  gogtattccc  tctgtgaaat  gagcagctcg  ttgtgcaaat  1980
tggttaagca  aacgtagaac  gattgcatc  cactttogct  catgagccac  aagcttattt  2040
gcgtatgac  gaaggtttgg  taacaaattt  ctttttctta  cctagcgagg  aagatccttc  2100
ttcgttggg  aatatctaa  aacattttct  aatagggaag  aggataaata  gogtgaataa  2160
atactgata  tggtagaat  aggcataaag  acctaaatcc  ttatgtcta  ttgattctca  2220
gtttccctac  agattattat  ttaogtatcc  tagaatagc  catccgggat  gcttcttgta  2280
aattggtata  taacgcagcg  ctgcattgtg  tggagcgagt  cctcttgat  caaaaacttt  2340
ctactgatca  agagggggaa  tcaagtattt  taocgaaagt  gattagcgag  ctacttgocg  2400
aattcgggga  aagttatgag  atttcagctc  aattacttgc  cgtaatcgat  atttatttaa  2460
aacaagagca  acgctogaat  tcatggttgc  ctogaatctt  tgcgaagaga  gacggggcta  2520
gaaaacgaca  aacaattaat  aagtgtcttt  tgttaaaaa  tatctattt  ttggaac  2577

```

<210> 33

<211> 554

<212> DNA

<213> Chlamydia trachomatis serE

<400> 33

```

ttcttttatta aaaaaaactt tctctttct ctcagacttc ttatgatgta agaaaactcaa 60
cgaagtcttgg tgtatggaga aggatttttt agaaaatgt ttctgtcaat tccgttacgc 120
ttttttttaa ttaagtgtac ttccagctct tctcggactc tggctatttt ttactcctaa 180
tattcttaac tatttggatt cttctgttat ttatcagat aaaaatttgc gogtcttttt 240
aatttttata tcagctttat ctttttataa tctcgtttat ttgcaactag gactttttat 300
tgggtctcggg gtttttttct tttctgttgc ttccgacctc ttcttttag tttttctca 360
tgattcgcga tcaggttttg ccacactagc tattattttt ctactcctca atcgtctcga 420
agatctagaa gttggttcta ctattccaga aacttgcctt ttaatactct tctccggagg 480
gaaaagagct cgggttctta tttttgctt tgtagtgatg ttcaaaagtc gctacttaac 540
ttccgcgca cgag

```

<210> 34

<211> 1433

<212> DNA

<213> Chlamydia trachomatis serE

<400> 34

```

ctgcacgaaa attattaaat gatcctttag gccgacgaac acctaatatt cagagcaaaa 60
atccaggtga gtatactgta gggaaattcca tgttttaacga tggtoctocag gtacgcgaac 120
tcagaaacgt cgacactggt ttttggtctgg acatgagcaaa tctctocagac ttgttatatt 180
ccagagagat tcaaacaggga cttcgagcacc gagctacttt ggaagaatcc atgcgcgatg 240
tagagaattt agaagagcgt tttgagcgtt tgcagaanaac ttgtgatcgc gctgcgatgc 300
agatagaaga atcgggatgg actcgagagt ccgcaiccaag aatggaagcg gatgagcgcg 360
aaggaccttc tagagcacaa caagccttttc agagccttgt aatgaatgt aacagcatcg 420
agttctcatt tgggagcttt ggagagcatg tgcgagttct ctgcgcgtaga gtatcacagc 480
gattagctgc cgcaggagag gcgattcgcc gttgctttct ttgttgtaaa ggatcgacgc 540
atcgctacgc tctctcgcat gacctatctc ctgaaggtgc atcgtttaga gagactttgg 600
ctagattcgc agatgatag ggaatagagc gaggtgctga tggaaacctac gatattcctt 660
ttgttagatga ttggagaaga ggggttctca gtattgaagc agaaggaatc gactcgatct 720
atgaatcat gatgcctatc latgaagtta tgaatatgga ctagaaaca cgaagatctt 780
ttgcggtaca gcaaggcgac tatcaggacc caagagcttc agattatgac ctcccaactg 840
ctagcgacta tgatttgcct agaagcccat atcctactcc accttgcct cctagatctc 900
agctacagaa tatggatgta gaagcagggg tctcgtaggc agtttatgct tcttttgtag 960
caggaatgta caattatgta gtgacacagc cgcaagagcg tattcccaat agtcagcagg 1020
ttggaagggat tctcgtgatg atgcttaccg accggtcaca gacatttaga gacctgatga 1080
agcgttgaaa tagaagaagtc gataggggat aaactggtat ctaccatagg ttgtagcaa 1140
aaaaactaagc ccaaccaaga gaaattctct ttggtggcgt tcttttttta ttcaaaaaag 1200
aagacccctct tcaagattat accaagatgg gatgtataat ctgaaaggaa gcgcttttat 1260
ttctatacca tatgatgggt gtggttatct cctttagagg agcagcagtc tccatgacgt 1320
tctttgaagc agcaactcaa gaagtttagc cagaccataa cccagcgatc tccggtactc 1380
acataagctg cttgtgtcca catggttctc tcaccaagca ggtgagtaag tag 1433

```

<210> 35

<211> 196

<212> DNA

<213> Chlamydia trachomatis

<400> 35

```

ctcgtgcgca tgatacagca gtgcgcagta atcaaggtgg taaacgcaag ggagctgtat 60
ggtctatttt agaagtttgg caactcgact acgaagattt ccttgaatgt agaaagaata 120
caggggatga gcgtcgacgg gctcatgatg tcaatatagc tagctggatt ccagatcttt 180
tttccaagcg tttaaa

```

196

<210> 36

<211> 1990

<212> DNA

<213> Chlamydia trachomatis

<400> 36

```

ttcactaggc tcatgagcct ctaactcttc tggagtaact cctagagcaa acacaaactg 60
ttctccacaa tcaatatgat tagggtaacc gttctctcca tccatcaagt tatctaaaca 120
taactctacg cctctctaat catcgcaacg actatgaatc gcagataaat atttaggaaa 180
gggttttgata tgtaaaatat agtctttggc atacgcctgt aattgctctt tagtaagctc 240
ccctctcgac catttcacat aaaaagtggt tcttagcata tgcatttttt gaataattaa 300
atctaactga tctaaaaaat tcataaacac ctccatcatt tcttttcttg actccacgta 360
acgcgttgca aaaaaggctc gtataagctc tctgtttcat ctatgcgcaa agaaacaatc 420
tctcttcogag aagtaggatg tgaatggtag accatattag gtgcctgctc tatccacgct 480
aacggtgttt gctcattccc ctctcccata caaacaacag ccgcaactgc taagcgatct 540
acaagattac ttgtcgctat ctgtagaaga cgacgcgaac aatctagcga tcttatatag 600
ttgtgtaatg tggtaaaatc ataccaacac agcccgatcc ccgactctcc agccgcattt 660
ggagtatgat ggctatctgt aatgattacg cctagctctt tcaactgaaa atcaattctt 720
aacattcttc cgtatcgatt acacgatccc aaaaatatct taggatataa acaaaaaagg 780
tgytccgfat togatctatc aatccctgca gaaggaaatc aaataacttc ttttttgctt 840
agatatatcc gctttttctc acaaaacaaa taagcatcgc ctcttttttt tctaacgctc 900
gctttgcaca tctctgcatc agcgacagcg ccttccacata aactccacat ctttgaagag 960
acaactacca caactcgttc ttgcagagcg ggcaaacgct cttgcaagat cctgtgaagc 1020
gaatcatgtg caaatctttt acgtgttttt atcggagtta ttcttcaaat aataaatact 1080

```

gaaatctctt	gtattacaaa	tacatttcct	cttccatctt	gataatcgcg	tgtatgggaa	1140
gaaagtctatg	ccccataatt	cttttttgat	atgtgttgaca	aaacaagctt	tcagaaagtt	1200
ttgttggaaa	aaactttcaa	agagctccgc	tcctccaact	aaaaacgact	gattccaaaga	1260
tagtgtccca	tactctgcac	aggaagaac	tcctattgat	tgtgtgtgat	gcattctctg	1320
agaaaagaca	acgatatccc	gccatgctt	atacttggct	ggaagagact	cccaagctct	1380
tgcgtccata	atgtatggat	gatttgcgat	ggttttctga	aaaaaacgt	gatctctggg	1440
ataactccaa	ggagctctgc	ctaaagctcc	catcactctc	ctgggatcaa	tagcaaacgt	1500
acctgttctg	tggatcatat	aaacatacca	gcccaagcag	cagcggtctaa	ggcaactctg	1560
ttactctcaa	ccgtatgcac	gcgtagataa	tcaactcttc	gatcatgaag	agatacaaga	1620
cagcgtatgc	tttcccaatc	acgatcgcta	ctattaatct	ggcccaacat	actcaaacac	1680
tgctttctag	ataaggctat	taatacagga	cactctaaaa	cagcttttaa	tcgtttact	1740
ccatccatca	ataacatcga	ctgaacggga	gtcttcccaa	atcttatctc	tggatcgaaa	1800
acaaactggc	aacttgtatc	taaacctact	tgagcaaat	gttctaaact	ggactctccc	1860
caacgcacaa	tttgcctaat	aggagattct	ctataagaaa	gtacacaact	tggctcttga	1920
ggcagcgac	acgaatgatt	tattaatagc	cgtagcccaa	actccttctg	caaatgagcc	1980
atttccaaag						1990

<210> 37

<211> 2093

<212> DNA

<213> Chlamydia trachomatis

<400> 37

cagaaactct	atccgcatac	cttcttcggc	aaattgtatc	caattttgcc	tcttctcagg	60
aagctactat	agctcagtat	attgtaggca	actctctccc	agaaggactc	tttttagaaa	120
atctcagttc	tgtggctgca	gatttaaagc	tttcggaaca	ccctttccac	aaggtatggc	180
aacgtatcca	acaattacat	cttttaggag	tgggagcgcc	ttccctacag	tcctactggg	240
tatcgctact	acagacatct	ccccataagg	aggcttttagc	tattatttgc	aaccattccc	300
ctcagtttagc	tgtttgtgat	ttoactacta	tgcgtaggaa	aatgcgatga	accaacaacg	360
agattcttacc	attttctaga	caagctttttg	cttccatccc	ttgtgtgtcca	gcagcagctct	420
tttccgagac	acigcaccccc	ccgtctccag	cgcttctctga	tgcctacott	tcctttctgc	480
gaactcttta	ttgggatgtc	tctattaata	aagatttgtct	ccctctatt	agactcaaacg	540
acacgctact	agatatctat	ccctctcttc	ctcgtgaaga	gaagagccac	ctatcgcaac	600
aaactcagagc	agcaatacaaa	ttgcttgcga	atgtaaaaaa	acgagaagaa	acgatttttg	660
ctatcctctcg	agttctcatc	ccctacccaag	aagagtttct	tccttaaaaa	gccactcttc	720
ctaaagcttt	tttgttaaaa	caaatagctc	gggaactctc	ttctcatgaa	gcacagcttt	780
gtcgcccatc	tgataataaaa	acgttagcaa	ccctgtttgg	attactccct	attgcgatgc	840
tatttccaca	agcggtttga	tcctggcccg	atccacttaa	agcaactatt	ttgcattgga	900
tccacccagt	gatttctaca	gaaaaacatc	ctctcttga	tcagctatt	agccaaaaaa	960
tatttagaaa	ggcactcccc	tgcgcacgac	gcacagtgcg	caaatatcgt	tgcgaactga	1020
atatccacc	tgcgcaccaa	cgcaaacacc	tatgtctgtg	tttaaacaca	acacgacagc	1080
agaattctcg	acatactatc	taatcggata	tgtaaagctg	otttcaatcc	cttgaactag	1140
aaataaaagt	gaaataaaaa	gcccagaaca	agagaagttg	ttctgggctg	acagaagctg	1200
tcagatcatt	ttaataagat	tgatgacaa	tacgacaagt	tcctggatcc	aaaaagaagt	1260
ctaaaaagcc	atacaaatgt	tgcgttaact	cttgcgatgc	ctctaaacat	ttatcagctt	1320
catcttttag	aagcatctca	atgagcgctt	ttttctctct	agcatcgccg	acatccgctt	1380
cttctctgtc	tgtgaaatat	gcattagttt	caggatttga	aaatccaaag	tactcagctca	1440
atccacgaat	ttttctctca	cgatagctg	gaatttgact	ctataagaa	tacaaaagac	1500
ccactctctg	agctaaagaa	tctctgtac	acacccgcac	gaagtagctc	actttctgct	1560
cttctcttgc	actaggtcca	tgaagctcta	actcttctg	agtaactctc	agagcaacaa	1620
caaatctgct	ccacaaatca	atatgattag	ggtaacgctt	ctcttctatc	atcaagtatt	1680
ctaacacata	cttaacgccc	tctaaatcat	cgcaagcact	atgaatcgca	gataaatatt	1740
taggaagagc	tttgatatgt	aaataatagt	ctttggcata	cgctgtaat	tgtcttttag	1800
taagctcccc	cttcgacat	ttccacataa	acgtgtgttc	tagcatatgc	ttattttgaa	1860
taattaaatc	taactgatct	aaaaaatcca	taaacacctc	catcattttc	tttcttgact	1920
ccacgttaacc	gttgcacaaa	aaggtccgta	taagttctct	gtttcatcta	tcgcgaagaa	1980
acaaactctc	tctcgagaag	taggatgtga	atggtagacc	atatttagtg	ctgctctat	2040
cacgcgtaac	ggtgtttgct	catctccctc	tcccatcaaa	acaacagccg	caa	2093

<210> 38

<211> 1834

18

<212> DNA

<213> Chlamydia trachomatis

<400> 38

```

ctctactctct acctctacgg ttacacgtgt agcggcaaaa gattatgatg aggcataaat 60
gaatttttgat aogggcaaaa gtggattaga gaacgtctaag acacttgctg aatagcaaac 120
gaaaatggctt gattttgatg cagctctcca agatatggag cgtttagcta attocagatcc 180
tagtaacaat cataccgaag aagtaataaa tattaagaaa gcgctogaag cacaataaaa 240
tactattgat aagctgaata aactcgttac gctgcaaaat cagaataaat ctttaacaga 300
agtggttgaaa acaactgact ctgcagatca gattccagcg attaatagtc agttaagatg 360
caacaaaaaat actgcagatc aaattatcaa agatctggaa agacaaaaaa taagttagta 420
agctgttctc actaacgcag gagaggttat caaagctctc tctgaagcgg gaattaaagt 480
aggacaagct ttgcagtcta ttgtggatgc tggggaccaa agtcaggctg cagttctgca 540
tgacagaccaa aataatagcc cagataatat tgcagccacg aaggaattaa ttgatgctgc 600
tgaacacgaag ttaaacagag taaaacaaga gcatacaggg ctaacggact ccgctttagt 660
gaaaaaagctt gaggagcaga ttagtcaacg acaaaaagat attcaagaga tcaaacctag 720
tggttcggat attcctatcg ttggtccgag tgggtcagct gcttcgcgag gaagtgcggc 780
aggagcgttg aaatcctcta acaattcagg aagaatttcc ttgttgctg atgatgtaga 840
caatgaaatg gcagcgattg cactgcgaag ttctogact atgatcgaac aatttaagt 900
aacaacactct gcaacagcta aagagctaca agctatggag gctcagctga ctgcgatgtc 960
agatcaactcg gtgtgtgcgg atggcgagct cccagccgaa atacaagcaa tcaaaagatg 1020
tcttgcgcaa gctttgaaac aaccatcagc agatggttgg gctacagcta tgggaacaag 1080
ggcttttgca gctgcgaagg ttggaggagg ctccgcagga cagctggcca cgttcagatg 1140
gaattgaaaa cagotttaca agacagcgtt tctctcgact tcttcagct cttatgcagc 1200
agcactttcc gatggatatt ctgcttacia acaactgaac tctttattt ttcttattt 1260
aagcggcggtt cagtcagcta ttagtcaaac tgcataatcc gcgctttcca gaagcgtttc 1320
tcgtttcggc atagaaaagtc aaggaagcag ttcatagctt agcacaagaa cagcagaaac 1380
tatgtccaga gatagcaaaa cgttaggtga tglatatagc gccttcacgg ttcttgatct 1440
tttgatgtct acagttcga gcaatccgca tgggttatcc gctgttcaga attctgcgga 1500
cagcgactct attagcaaa ctccacaatt tgggttatcc gctgttcaga attctgcgga 1560
tagctttcga aagtttgctg cgcgaattga aagagagttt gttgatgggg agcagatctc 1620
cgcagaatct caagagaatg cgttttagaa acagcccgct ttcattcaac aggtgttggt 1680
aaacatttgt tctctattct ctggttatct ttcttaagc gtgattgaag ttgtgaatg 1740
aggggcgccc aaaaaagaat ttcttttttg gctctttttt cttttcaagg gaactctgtg 1800
tctacagaag ttttttcagc acgagoggca ctag

```

<210> 39

<211> 1180

<212> DNA

<213> Chlamydia trachomatis

<400> 39

```

agaaaatttct caaaaaacaa agttttttac atttaagggg catcttacca ccacaacaac 60
ctctatagag caaactatcc ccatiaaata aagaataatta aatatacaaa aacatctgtg 120
attatttttg ttaaaagaaa tacttaatga gttttattta attaagcaaa cgaagaagctt 180
gctaagtgaat atatttcaca cagctatcga atttgctccg gtaatcaaaa cggagagcct 240
gggagagcgg ctataccgag tagcaaaagc tttagccgct aatcacacaa cgggaagtggt 300
aatcccttta taccttaaat tatttacttt gcccaagaaa caagactottt gctogatoca 360
aaaattatct tatttttttg ctggagagca agaagcaact gctttctcct aettttatga 420
aggaattaaa gtaactctat tcaaacctga cacacagcca gaggttatct ctgagcggga 480
aacaatctac acaagcgatg atgccttcgg tttttgcgct tttttctcgt ctggcgccctc 540
ctacatccaa aaagaaggag ocaatatcgt tcatttacac gattggcata cgaatttagt 600
tgctggagcta ctcaaacac agccctgctc tcaattacaa aagattgttc ttaccttaca 660
taattttggg tatcaggact atacaacagc agaaattata gaagcctcct taagcttaga 720
attttatatac agccagttac aactatttgc cgatccacaa actgtgtgt tgctaaaagg 780
agctttatata tgttcagatt tctgtactac gttttctcct acatcgcga aagaattctc 840
tgaagatttat tccgatttag aaattcacga tgcatttact gctagacaac atcatctcgc 900
cgggatttata aatggaatgc acacagcaat ttgggggctt gaaacggatc caaatttagc 960
gaaaaactac actaaaagagc ttttcgagac cccttcaatt ttttttgaag ctcaagcgca 1020
gaataaaaaa gcctgttagc aaagatttag cctctcttta gaacactctc tctcgtgtgt 1080
cattatttctt agaattgctg agcagaaggc tctctcattt atgaacagc cacttctcca 1140

```


tgactagaa aacgcttaca cgctcattat tataggtacc

1180

<210> 40

<211> 1297

<212> DNA

<213> *Chlamydia trachomatis*

<400> 40

agaaactct ataggagggt atgtgatcga cataggtacg tgtgagttat gggatatcga 60
 atttgggtat aatggataag aaattctctg aagataaaga ggctcctcca actaaaagac 120
 catcaacatc agggcagagg gcaagtgcgc gagcattatc ggctttcaca gatcctcgt 180
 aaagaatggg ggtgcgttcc gcaatattct tggaaaagag agaagcaatc gtttttctac 240
 agaaaagcatg ggtttcctga actagatcag gatgagctac tttccgggtg cctatagccc 300
 agactgggttc ataagctaga atgaagaggc ctgcctcagg gagtttagat aatcctatag 360
 tcagtttgatt taaaagaata tcttgagttg ctccagattc ttgttctctc aaagtctctc 420
 caatacacag aactggaatc attccactat ggatagctgc agcagctttt tcagcaagta 480
 caggattttg ttcattgaaag atatgacgtc tttcggaaag tccgatgaga acaaaatcga 540
 ctccgatgac tttgagcatt ggggctgaaa tctcaccagt aaaagctcct gagtcagctt 600
 catgagtggt tgggctccca agaaagatgg gggaaatcgt lacagcttgt tgacaagctg 660
 acagcagttg gaaaggagga atgattcctg taatgatttg gggattagac agaattgcac 720
 tagagatgaa actttttaa aaggtctgag ctctggtaag cgtcttggtc attttccaat 780
 tacocgaaac aaattgcttt gatggctcag agtgggaaag gtggggcccaa gtgggaaatg 840
 gttttctgtg agtttctttg tctgtaaaac tgagattttg tgaataacct gtgcagtgtat 900
 ttgttttgta agatagatca aaggttaata ctgattttct tgcaaggaaag gcttattttt 960
 alatagttta ttctctattg ctttgatata aatctcttgg atatgctaat ctctctgctc 1020
 tacttttttc tgtgaatttg cttaaatagt tggtttttag cctcttggtt tatgaaggtg 1080
 aaaaattttg gtattacgca tctctgalgat gcctgggaag ctgcccgaagc gggagccgat 1140
 tacattggca tgaatttttg taagaattct cgaagatgtg tagtagaaga aaaagcaaat 1200
 tatatcgtag aggtctatca ggaagggaat tcggaacctg ttggagtagt cccagagact 1260
 tcagtagaag aaatttttagc tattactgag acgacag 1297

<210> 41

<211> 1141

<212> DNA

<213> *Chlamydia trachomatis*

<400> 41

ctttccataa gttctttctt tcaatgattc tagcttattc ttgtctgtct ttaagtgggg 60
 gggggatgac agcagaatcc atgattcttc aaggaattta cgtatgggag acgttaaactg 120
 tatcatcttc ctatactgtt ataggagalc cgatgggagc tactgttttt tctgcaggag 180
 agttaacgctt aaaaaattct gacaattcta ttgcagcttt gcttttaagt tgttttggga 240
 acttattagg ggtttttact gttttaggga gaggacatcc gttagacttc gagaacatac 300
 ggaactctac aaatggagct gcaactaagt acagcgctaa tagcgggtta ttactattg 360
 aggggtttta agaattatct ttttccaalt gcaactcatt acctggccga ctgctgtcgt 420
 caacgactaa taatggtagc cagactccga cgacaacatc tacaccgtct aatggtaacta 480
 ttatctctaa aacagatctt ttgttactca ataagagaaa gtctelcaltc tatagtaatt 540
 tagttctcgg agatggggga gctatagatg ctaagagctt aacggttcaa ggaattagca 600
 agctttgtgt cttccaagaa aatactgctc aagctgatgg gggagcttgt caagtactca 660
 ccagtttctc tgotatggct aacgaggctc ctattgcttt tatagcgaat gttgcaggag 720
 taagaggggg agggattgct gctgttcagg atgggcagca gggagtgtca tcatctactt 780
 caacaggaaga tccagtagta agtttttcca gaaatactgc ggttagattt gatgggaacg 840
 tagcccgagt aggaggaggg atttactcct accgggaactg tgccttctgc aataatggaa 900
 aaacottggt tctcaacaat gttgctcttc ctgtttacat tgcctgtgag caaccaacaa 960
 atggacagcg tcttaatacg agtgataatt accggagatgg aggagctatc tctgttaaga 1020
 atggtagcga agcagcagga tccaaatact ctggatcagt tctctttgat ggagaggagg 1080
 tagttttctt tagtagcaat gtagctgctg ggaagggggg agctatttat gccaaaagac 1140
 t 1141

<210> 42

<211> 822

<212> DNA

<213> Chlamydia trachomatis

<400> 42

```

cgccacagagt gtagctgaa caagcagaag ggccactga gaacgagcct ctgagaaaaa 60
aagcttttat taaaaaatta aaaaaaatac ttacaaaact ttactgtag ttgagaaaag 120
agotccaagc taagcatcc aaagctccgt acttacaata ttattgcgga tagagcgaaat 180
taattctctt ttttagtgat gaagaggttt ttggggctg aagcgagcca aaagatcttt 240
atgcgcaact tgacgagcta actctaacac ccgttcgata tcgggttttt tgaattcaac 300
aaagtctctg cgctttttag aacctcgagg agctcgtggt ttagggtctaa tggatctggg 360
agtgatagaa togatocaaa acgtcttttaa catttttaac agttgctcag gagcagagtt 420
cttcatTTTT tttaaagtaa aatgatgcac gtgacccctt gttggccctg ggagataaac 480
acaaagatca ttttctttac ttctccgacg tttgtcaatc gotttagtta tgagctgtcc 540
tatttctctt tggatagtaa tctgtccggt agcatgaat agctccttag tgggtagctc 600
agttctacag atggtagttt ttgctttatt aattgtaata gtcaactaag tctgtttttt 660
togatattat ttctagtcga aataaaaatc aattagtgtt tatcttttgg tgatttttat 720
agtggttttt cgttttttcc caatctcatt tttagatttt ttttgatttg gacaaaagaa 780
aataaagtat ttcagattgt tttctaagtt tgtttgcata aa 822

```

<210> 43

<211> 1634

<212> DNA

<213> Chlamydia trachomatis

<400> 43

```

ataaaaattt aaattttggc tactccctgc tcttaataga atttccaccg aggagcttgc 60
tactgtattt gcattttctc tagggagagt agctgacgta ctggtagcat ttgcatctag 120
tactagtatt acaatatctt cattaaaaat aatagatcgg ccgttcggag aaattttggg 180
gttaactggtt ccgtctatat agatagcgcc ccccttatta tggttataaa 240
gtaggtaggg caattatcca ctagggtaac taacaggacg taatatagctt cgcataaatt 300
tttttgata ttgtcactaa aaaagatcct accacgattg cctgttaact cttaggcagat 360
agttacttta attgctcctc catcagaagc tctgaagaa gctgtttcta catttttaaa 420
gcagcagattg ttatagaaaa cgaattgtacc acgatttctt gttagaagac agtataggga 480
gaagatcgct cctcctgcac aacaggcggt attgatgaag aagagatcgc agttattact 540
ctcaaaaagaa ttgctcgttc cagcatagat agcgccaact ttctcgtctg tattagtttg 600
aatcacagatg ttgtccataa agagaaaaca agaactgatt tcgtccacaa caaagctatt 660
agcggtaacta atggctcctc cttggaacta agaaaagttc ttcaataatc cgaaccatc 720
atgattatga tttatgtaa gattttgagc atgaattggt ccgcctcttc ttattttatc 780
agcagcatata ggattttctc atgtaaatag tctgcaacaa gtattatttt caaagattac 840
aggaactatg gtatcacgaa tctccacggt aggaatattg ggaactgcac aaccaatcgc 900
accaccactt tcaggggtga gattttttgc aaaataaata cctctttttt gtgtatcaaa 960
aaagcttagg taactctgta ttgtgacagc agctccttca ttgggagttt ttgtagatat 1020
agccagtagt tagcgtaggt tatcgagata gcagtttagt agattgtgag tgtctcctgt 1080
caactcaatt tttttgata gogaactctt cgtaggatct ggaactgagt tgggcaataag 1140
aaagattcta gaaggaacct ctctagctag tctctgtagg gattttccga taagaaaaaa 1200
gaaaaacgct tttttcataa taaaagaacc agagctcctc ctgcattgat gtatgtgag 1260
acagtggaag, tagccaactt tgcttgatag tttagcaata gtttcagatg agaaaaattg 1320
agggagttag aacctctccc ataaaaggaa ttgttagcta atggggtatt tgtggtgacc 1380
gaagaacogt tattttgat taatagtgtg ttgagttagg gaagtttcca gttaggggtg 1440
cgttggtgaa gtgtgtccat ttcccaagag agtgtgtggc atgtatcaga agaataagct 1500
cctttgattc ctattggaga gacaaacgca gtatgggctt gctcataagt aaataatcta 1560
gctagatcac cgtttctctg gatagaagct ggttctgttc gagagaataa agcctgagca 1620
aatgggtgta gcat 1634

```

<210> 44

<211> 1862

<212> DNA

<213> Chlamydia trachomatis

<400> 44

```

gttagctttc cctccaggga ttgcaattt aatgatttta atgatttttt tattcgacat 60
attcaacctt ttcatttggc aagacatgga gtcatagtta gagggctcat gtatgctct 120

```

```

ctaacaagca atatagaagt atatggccat ggaagatat agtatcgaga tacttctoga 180
ggttatgggt tgaagtgcagg aagtaaagtc oggttctaaa aatatttggtt agatagttaa 240
gtgttagoga tgcccttttc tttagagatc acatcaattt gtttttttagc ttgtttgtgt 300
tccatttcgt atggatttcgc gagctctcct caagtgttaa caccataagt aaccaotcct 360
tttaaggggg acgatgttta ctgaaatgga gactgcgctt ttgtoaatgt ctatgcaggg 420
gcagagaacg gctcaattat ctocagtaat ggcgacaatt taacgattac cggcaaaaaac 480
calacattat cactttacaga ttctcaaggg ccagttcttc aaaaattatgc ctctacatta 540
gcaggagaga caactactct ctacaggaaa acogtgagta ttctcgagac agggcaagtg 600
tgcgagaaaa agggaagtgt ctacaggaaa acogtgagta ttctcgagac agggcaagtg 660
attttttggg ataactctgt ggggtattct ccttctgtct lttgtccagc atcgaotcca 720
actctccagc accacgcacc agctcctgct gottcaagct ctctatctcc aacagttagt 780
gatgtcggga aagggtctat ttttctgta gagactagt lggagatctc aggcgtcaaa 840
aaaaggggtca tggtcgataa taatgcgggg aattttgtaa cagtttttgc aggttaagt 900
aataaataatg ctggtagtgg gggtagtggg tctgtacaaa caccagattt tacagttaaa 960
aactgtaaaag ggaaagtctc tttoacagat aacgtagctc cctgtggagg cggagtagtc 1020
tacaanaagaa ctgtgtcttt caaagacat gaaggaggca tattcttcog agggaaacaca 1080
gcataagatg atttagggat tottgctgct acagtgtggg atcagaatcc ggagacagga 1140
ggcggtggag gaggatttgg ctctccagat gattctgtaa agtttgaag caataaaggt 1200
tctattctgt ttgattacaaa atttgcacaaa ggcagaggcg gaagcatctc aacgaagaa 1260
ttctctctgt tagcagatga ttoggttctc tttagtaaca atacacagga aaaaagcggt 1320
ggagctattt atgtctctac tatcgatata agcacgaatg gagagtcgat totgtttgaa 1380
agaaacccag atgtccgaag aggcgcacac tgcgtgagtg aagcaagctc tgggttaoact 1440
ggaaactctt cttaagcgc ttctgatggg gatattgttt ttctgggaa tatgcaggt 1500
gatgctctg gagagccgag ccgacagaga attcttaagt atggaacgac tggttcttta 1560
aatgcttcog gaactacgaa gctgatcttt tatgatctgc tagtacaaaa taattcaaga 1620
gcgggtgcat cgcaccatc acocatctct tottctatgc ctggtgctgt cacgattaa 1680
cagtcoggtta atggatctgt gatttttacc ccogagtoat tgactctctc agaaaaaact 1740
caagttctta actctacttc taacttccca ggagctctga ctgtgtcagc aggggagttg 1800
gtgtgtaggg aaggagctac cttaactact gggaccatta cagccaactc tggctcgtgc 1860
gg 1862

```

<210> 45

<211> 1668

<212> DNA

<213> Chlamydia trachomatis

<400> 45

```

agaaaaatcc atagcagaaa tagaagaatt cgtatgtggt gcgaacaaag ctcaagattg 60
ggatgtcgtat gtagctatgt caaattcttt tggtttttggc ggacacaaatt caacgatatt 120
atttttaggg tatgaacctt cattatgatg aaaaactaagc acgaatatctc ttgttgcgtt 180
attcctatcca gatttttttg tactcoggat agaagtaacct taaaggcttg ttctatctgc 240
catcacagatg gaaaacattg gggtttccct agagggcgat gggagaaaaa aaggggccct 300
caggaagcgtc ctgagagaga actttagaaa gaaactgggt tggggattgt taatttttct 360
ccaaaaaatat ttgtggaaaa ttattccttt aatgacaaaag aagaatctct tgtacgtaaa 420
gaggtgaactt attttcttgc agaggtttaa ggcgaagtaac atgctgatcc tgatgatagc 480
tgttatgtgc agtggctaag ctctcaagaa ggtttacgoc ttftaaattt occagaaatt 540
ogtaaatgtt ttaocggaagc agatgaattt gtctaaagtt atctatttgc ttcataaagt 600
cccatagatt gaaaaaaaact tggttaggag atcaaatctc ggaatctccc aacacagcct 660
ttcttttttc tctcogattta catataaaga ttgcaatagt ctctgtgagc aagacgaagt 720
actttttgag ctctctttttt gcogtataaa cctacaattt caatttttag tggttttgtg 780
tgaattlaagc ttctgtggat agottttatg gttaaagaatt agtgttgagt catgtccaaa 840
actgtgctgt ggcattcaaga aatatctctc atattgcoat agatcaaatc atctctotaga 900
acagagatga ttatcatcgc gggtctctcc gagtctaaaa taogaatccc tccagataga 960
cgogcttgca agaggtatgt cctctgtgta atatttttct cgttaaacac acagatatca 1020
agagatgctc catcgctctt gatattctct atatttttct gttgacacct gtattctcca 1080
gaagatgctc cacaataagt cttaggtaac agcccgata atcaaggga aaattagaaa 1140
aaattttgtc gcogatccac tttaggata ttttaggata caaagagatt cataatcact cgtgttact 1200
gagtgctgtc gaggtagttc tatatagcaa aatgtgttgt tagacataag atcaactctc 1320
actggcccat ccgaaggatg agctatggat gcatcoggtc tatgttccca gatagtttta 1380
gtctctgttt ctocagtaaa aacttttatt ttatcggcag ctctctttt ttgttttaatt 1440
cttgtcattg tgaaaaatgt tgaaaagtta ctctgtgcaa cctttacagc aggttttttg 1500

```

tacgaagaag	gagagtgatt	gtactgcmaa	ataatatgag	ccggacgtag	gatatgaatt	1560
actcttttgc	aatagaagac	ctacatatgt	aaggatatga	acaggttttt	aaagttaact	1620
gogagtgctg	acaggttagt	gctgtaattg	ctattcatca	gacaaaag		1668

<210> 46
 <211> 2010
 <212> DNA
 <213> Chlamydia trachomatis

<400> 46

atatcaaaagt	tgggcaaatg	acagagccgc	tcaaggacca	gcaaataatc	cttgggacaa	60
catcaacacc	tgtcgcagcc	aaaatgacag	cttctgatgg	aatatcttta	acagtcacca	120
ataattcatc	aaccaatgct	tctattacaa	ttggttttga	tgcgaaaaaa	gcttaccagc	180
tattctctaga	aaagtgtgga	gatcaaatcc	ttgatggaa	tgctgatact	attgttgata	240
gtacagtcga	agatatctta	gacaaaatca	aaacagaccc	ttctctaggt	ttggtgaaag	300
cttttaacaa	ctttccaatc	actaataaaa	ttcaatgcaa	cggtgtattc	actccacgta	360
acattgaaac	tttattagga	ggaactgaaa	taggaaaatt	cacagtcaca	cccaaaagct	420
ctgggagcat	gtctctagtc	tcagcagata	ttattgcatc	aagaatggaa	ggcgggtgtg	480
ttctgacttt	ggtagcagaa	ggtgattcta	agccctgcgc	gattagttaa	gtgactactc	540
caggcatctc	taattttatg	agtctaagaa	ccagttattc	taatacagga	ttgactcgca	600
caactgtatc	attactgtga	ggcggtttag	aaagcgggtt	ttgatgggtt	aatgcccctt	660
ctaatggcga	tgatatttta	ggaataacaa	atacttctaa	tgatatcttt	ttagagttaa	720
taactcaaac	aaacgcttaa	acaattttta	ttggattttt	cttatagggtt	ttatattagt	780
agaaaacagt	tgcgaattag	gggtttgtta	tgcaaaataa	aaagaaaagt	agggacagtt	840
tttataaat	tgttaagaat	gtgaaaaag	atttcccgca	attagacctc	aaaaacagag	900
taaaaacaga	aaaagtaact	ttcttaaat	ctcccttaga	actctacact	aaaagtgtct	960
caactaatct	aggactgctt	caacaaatag	aaaactcttc	aggattattc	ccagactctc	1020
ctgttcttga	aaaattagag	gataaacagt	taaaagctta	aaagcgtttg	attatgctta	1080
ctttgtctag	aaaagacatg	ttttcccaag	ctgaatagac	aacttaactc	aatctgtggag	1140
ttgatcttga	caccttagtt	ttttgtctct	tttagggagg	aacttgaaaa	acaacacttt	1200
ctctaaacgt	gggatgcaac	ttggcccaat	ttttaggaaa	aaaagtgtta	cttgcgtacc	1260
tagaccgcga	atccaattta	tcttctggat	tgggggctag	gtgcagaagt	gaccaaaaag	1320
gcttgcacga	catagtatac	acatcaaacg	atttaaaatc	aatcatttgc	gaacaaaaaa	1380
aagatagtgt	ggacctaatt	ctgcatacat	tttcatccga	acagttttag	gaattgtgata	1440
ttcatagagg	acctagttaac	aacttaaaat	tatttctgaa	tgagtactgc	gtctcttttt	1500
atgacatctg	caataatagac	actccaccta	gcttagggag	gttaaocgaaa	gaagcgtttt	1560
ttcggcagga	caaattaatt	gcttgtttta	ctccagaacc	tttttctatt	ctaggggttac	1620
aaaagatacg	tgaattotta	agttcggtgc	gaaaacctga	agaagaacac	attcttggaa	1680
tagctttgtc	tttttgggat	gatcgtaact	cgactaaaca	aatgtatata	gacattatcg	1740
agtctattta	caaaaacaa	cttttttcaa	caaaaattcg	tcgagatatc	cttctcagcc	1800
gtctctctct	taaagaagat	ctgttagata	atgtctatcc	taactcttag	gccgcagaag	1860
atattctcga	gttaacgcgt	gaatatgcaa	atatttttga	catcgaatat	gaacgagatt	1920
actctcagag	gacaaactga	acaaactaaa	aaaagaagcg	gatgtctttt	ttaaaaaaaa	1980
tcaaacctgc	gcttctctag	atttttaaga				2010

<210> 47
 <211> 2044
 <212> DNA
 <213> Chlamydia trachomatis

<400> 47

gtcatcaaga	aaagattggg	aacotatcgc	tatgtttggt	aaagacatgc	gtatgcgaca	60
gtgtcagcgt	attggtcatag	cttcgcagac	gaagatctcc	aacattatag	gagtaactca	120
atctatttgg	ccaacgtaca	aaacatttgtt	tgtaagatct	aatgttccgc	gagaattcac	180
gattccaagt	gtgataatga	ttgagaagtt	gaagaataa	ggtatctggg	atgctgatata	240
gttagatgac	ctgaataatt	ttgttgggtc	tttatttgaa	atcgagcgta	taccagatca	300
cttaaaaact	attttcttga	cagcttttga	gaattgaacca	gaatggatta	tcgaaatggc	360
gtctcgaaga	caaaaatdga	ttgatattgg	gcaattcttc	aacctttatc	ttgcccagcc	420
agacgggaaa	aaactgtcga	attgtatttt	acgggtttga	aaaagaggtt	tgaaactaac	480
gttatattctg	agatctctcat	cagcaacgac	cggttaaaaa	ctcttttgag	atcttaataa	540
gagaggaatt	cagctctggt	ggatgaagaa	taagtctgct	tcggcagaaa	ttctgttgta	600
aagagcgaa	aaagcacctg	tctgttcttt	ggaagaaggg	tgtgaagcat	gtcagtaatt	660

```

aatcatataa attaacaata aaattaacgg ttcttatgca agcagatatt ttagatggaa 720
aacagaacgg cggttaatacta aatagcaagc gtctagttaa ctgcaaccag gtccagtgtca 780
accaaatgtg tcttaataag tacaaatggg cttgggaaca ttatttgaat ggtctgcgoaa 840
ataactggct ccctacagag atccccatgg ggaagacat cgaattatgg aagtcggatc 900
gtotctotga agatgagcgg cgaatcattc ttttgaattt aggttttttc agccacgoag 960
agagcttggt tgggaataat attgtctag caatttttaa caatgtaact aatccggaa 1020
cgagacaata tcttttaaga caagcttttg aagaagcggg tcacagcagc acatttttgg 1080
atatttttga gtcaactogga ttgacagaga aagaaatctt caatgcctat aacgagcgtg 1140
ctgcgattaa ggccaaagat gattccaga tggaaatcac tggcaaggta ttggtatccta 1200
attttgcac ggactctgtt gaggtctac aggaatttgg tggcaaggta gtgagatcct 1260
acatcattat ggaagggatt ttctctata gtgggtttgt gatgatcctt tcttccaca 1320
gacaaaataa gatgattggt attgggaac aatatcaata catcttaaga gatgagaca 1380
tccactgaa ctttggatt gatttgatca acgggataaa agaagagaac ccgggtatt 1440
ggactccaga gttacagcaa gaaattgtog aattaataa gcgagctgtc gatttagaaa 1500
ttgagtatgc gaaagactgt ctccctagag ggattttggg attgagagct tcgattgtca 1560
togattatgt gcagcatatt gcagacgctc gtttggaaag aatcgagta aaactatt 1620
atcatcagaa aaaccatttc ccttgatga gcgaacaat agacctaatt aaagagaaaa 1680
acttctttga aacaagggtt atagaatata aacatgcagc aagcttaact tggtagctct 1740
gattcaaaaa taggagaaag cctcaacctat agagtggag ctttttttgg ttctatcgta 1800
acctgataag aatttttga ttttcaggtt agaagtaaat gtatttacc atgaattttt 1860
tttatttttc ctaataatc ttgtagccct ttattaaaa ttgaaagggc tagtcaactc 1920
tctatgact actgttagag tggtagagtt tggggttggg gcaggtgtag cctttcgat 1980
acgaagtatt tctctgtgaa accacaagat ttgaaacttc cctatttttg ggaagaactg 2040
tctc
2044

```

<210> 48

<211> 3734

<212> DNA

<213> Chlamydia trachomatis

<400> 48

```

gttattcgct tctaactcat tagaagtcac taatgtcaaa ctaccattt ttctccttt 60
ccgttaaaac aggaagaataa ttgtacagaa acattttttt aaagaaatca aaaagccatt 120
tgcaggcaga tatcaggcca ttatatcaa aaacagaaag aatgattagg ataaaacttt 180
gtcttgcctat cggtccagag agoattgaga agccgttttt attataaata caattgcaata 240
agaattcttaa attcgaacag acaacacaat ggctcgaaca gactgatcca cagcactaaa 300
ttcaaatgca aaaaactctt aaaatgaaca cagcaagctt gatataaaca tataaagaaa 360
ttggatcata gagctttaag agaagggcgg cactgcaatc tgtctcgacc aaatagcaat 420
gcaaacagat aaataccctt aatcattggg aaaaatttag tgtagaatag cctctttctc 480
ttctctatt ttgtgttag ctaacgcgat ttctcttata gagatactgc agagctctgt 540
cttatccaaa aagccttgct totcattttc caatacaaat ctgtccagag aaactttttt 600
tggctctcca ccatalagct aaattctagt aagaacagca ccatgatcca cagatccaaa 660
acaacacagg gtaaacacta cgtcaatcat aggaagcgta gtccaaactg cccaataaaa 720
taaggctgct cctgtaacta tgaataaaat actaagaata ccgagcgcaa gcacagcaat 780
cagttcgcta caacaagaaa ctctcgcttt agaagcgcta tccaccaag gagcctctgg 840
catataactt ctgaaggata cactatctcc acaaaaactc attggaatcc ctcttaagta 900
aaagagaagc ttctctctaa atagaagaag gtatcgtcaa ctcttttata gatctaaaaa 960
catataactt ccttaatccc acccatgaaa tttagatata aaacatacca acatattccc 1020
agcgtctttt aaaaggccta ttctcattt ttcttgagtc tctaaaaacc tataatggct 1080
ggaaattttt ogcgcaactt ccttggcttc ttgtaatag ttgtatgaat tgggtatcac 1140
agataacagg taagaaacta atccaaaagg toctatacaa gaaccaataa ttgcagctct 1200
ccactactc ctaaaactaa ggaagaatag aactccccaa aactcctctc aactcctctc 1260
aaagctgcga gccaaactgt tagaacaact acaaataact ggtatgtttt agaagcgtga 1320
ataaaggagt tgttagccac atttctaat ttatctagtt ttgtcgtaac acaactttcc 1380
atatcagact tcttccaat aaacgttcat aattactcta ctggaaacc attaataatt 1440
gcagcaagctc taactotttg acccttttg acccatttgc aagctgtgtc atttttttct 1500
ctctctgttc atatagaagt tcttgataag ataacactaa tcaaggagtc aagcgtctgt 1560
gagctcaact agatgaact gaagggagtc atacacacta tcaagagtc ccaagagtc 1620
aaactagact tctctctata caggcagcac cagctgtgat taaagaaac ccaatgaat 1680
gagcaaaaag agagacaca cttctcgaaa caagagcagc aagagaccca ctaaaaaactg 1740
ctagtacccc taataacaa gcaactattg ccaacaattg ctcttagtg cttgtagtag 1800

```

tctgagcacc	gatagtttca	gtatgatcgg	cacgcaatgg	tttgctggaa	ttacaacaaa	1920
aagaataatt	aaacatggcg	oatctatttc	gcaaaaaaaa	ggccaacatg	ctacagggaa	1980
gctaatttaa	gtaaaaattt	ttatatattt	caatggtagt	taataaccta	atctaccocaa	2040
ccaaaagatg	tctaaatgac	aaaaaataaa	togtatttat	attatcatga	gacacttata	2100
gtcacgtctg	ctcaattcag	ctcaaatctt	aatgaaaaat	cggatttaga	agaaaataga	2160
ctcggaaggt	cagaactagc	caaaaattgt	gtctaatctc	tattttgcaa	tcccgagacta	2220
caagaccaat	agagaaacgt	taacctactc	octaagccca	cagaaccaat	cataatcgt	2280
ccaataacct	aaccggcaaa	cacaagcgac	gatccccgcg	aaagcaacaa	aagcaaggct	2340
acacaactta	aaatagcaaa	aattcctaag	gaacggccaa	attctatat	tcctcttcgt	2400
ttgcaataaa	tatgcgtctt	atacagacac	aactctcgcg	ggctctccag	agttggagcg	2460
caagaggaac	aaaaaagata	agacattgtc	gaactccgac	caaaaaaagg	cagatagata	2520
cgcgagatgg	taaaaaatac	gaatatattt	tgcacatgaa	aaacctaac	ctcctttcat	2580
cgctgagac	tagagtgtaa	acaaagatgc	gaagcaagg	ttcgctatg	ttggaaccaa	2640
acotccacac	ggtcccgat	tatcaaaaca	agctctccag	ggatattgta	gagaaactgc	2700
tatccatacc	aaagcaaat	atagacgtct	ttgtgaaaa	gaactgaatg	aggaactcaa	2760
gaactctggg	tagcgtctat	agatgcttta	agagcagctt	ttctcttttc	agcactatcc	2820
aaacatcttg	tgtagctaga	taaaaactaag	cgacatcgg	acaataaagg	ttgctcattt	2880
ttctctcaatc	tgtccaaaca	atcaatctca	actcttattg	cttagtcttc	caaaagctgg	2940
agactctcgg	taagaactcg	cagaaacatc	ttattaatga	aagagacgga	gaccacaagc	3000
tcctctctct	ctgaagaagt	acgcaaacgt	tgtctagcca	aaactttttt	tgtctctaa	3060
ctagacataag	agactcgaca	catagagcga	gatatctccg	accocacaca	acgacatcca	3120
ataattaatg	cagcaatacc	tattgcagta	aataatgcat	tgtctagcca	caaaaaaaaa	3180
gctaatccoc	cagcgacaa	aactaaagcg	oatcgatag	ctaaagctat	atccaaaagt	3240
tttgaacagg	tattctcttt	tggtgaagac	gaagtgaat	ctaaagctat	caggaagct	3300
gttgcgcaatg	taaaagaaga	agcgtctcgc	ctaatagtat	tactaatctt	tcocactttt	3360
tattttttaa	acggaaaaac	tgtatcagaa	cgcgctttta	ttcgcaaatc	attataaatc	3420
cgcaactatgc	agaactaaag	cgccgttaag	aaaaggaaac	oatcaactct	agatgcaata	3480
ttctgaggagt	ctttaattat	ttttacagac	gggatgcgtg	caactcgacg	cgctctgata	3540
atgtcttatt	ctcagatctc	aatttacaca	actctgctgt	taattgaactg	caagtgatct	3600
gaacttcttg	caaccgtctg	ttaaacocct	ctgtctgatg	aaactttct	tgtctcagat	3660
cctcctcaat	ggagcaaaat	gtttcggcat	aacgcttaca	caaatctaat	atttgttctt	3720
ccaactcttg	gcaa					3734

<210> 49

<211> 2937

<212> DNA

<213> Chlamydia pneumoniae

<400> 49

atgcctcttt	ctttcaaaatc	ttcatctttt	tgtctacttg	cctgtttatg	tagtgcgaat	60
tgcgcgtttg	ctgagactag	actcggaggg	aacttggctc	ctccaattac	ctcagcagg	120
gaagagatct	tactcaacttc	agattttgtt	tgttcaaaat	ctctgggggc	gagtttttca	180
agttctcttta	tcaatagttc	cagcaatctc	tccttattag	ggaaggcctc	tcctttaacg	240
tttaacctct	gtcaagctcc	tacaataagt	aactatgcgc	taactttcgc	cgcagagact	300
cttcaactca	agaatttttc	ttctataaac	tttaacggga	accaatcgac	aggaacttgc	360
ggcctcatct	acggaaaaga	tattgttttc	caactatcca	aagatttgat	cttcaactag	420
aaacgtgttg	cttattctcc	agcatctgta	actaactgct	caactccgcg	cgctctgata	480
gtcaactacg	gagcctctgc	tctccaactc	acagactcac	tcactgtcga	aaacatctcc	540
caatcgatca	agtttttttg	gaacottggc	aacttggctc	tgtcaattag	cgctctgata	600
agggcagctg	ttaattctat	caatacaacc	gtcaccatga	gctcttccca	taactttact	660
tcgtcaggag	ggcggtgat	ttatggagga	agctctctcc	ttttgaaaa	caactctgga	720
tgcatactct	tcacgcgcaa	ctcctgtgtg	acaagcttaa	aaggcgtcac	ccctctacca	780
ggcaactatg	cttttagaag	tggcggagcc	atctgcctcc	ctacgggaac	tttcgaatta	840
aaaaacaatc	aggggaagtg	cacctctctc	tataatggta	caccaaatga	tgcgggtgag	900
atctcagcgt	aaacctgcga	catcgttagg	aaccaggttg	ctacgggttg	aggaagcgtg	960
ctctattgaat	ctctagaaaa	cgccggggag	gtcaaaagtgc	tcaaatatca	aggaagcgtg	1020
gtttggagac	ctcggaagca	aacatcgaca	aggggtggag	ctattttcat	agtcctctct	1080
cgcttccaa	gaaacatgct	caatacaaaa	cttgcagattt	tgctctcaga	aggtgatatt	1140
gcagggggag	agattgtgtc	cttatctgca	cctggaaatcc	gcaatcgatcc	caactgataa	1200
ccattataca	atagctccc	aaccacaagt	cagtgcttata	caactgtgtg	attttatgat	1260
				aagcaactac	aatcaacgct	1320

```

aatggcgctt caggatctgt agtctttaca agtaaggagc tctcctctac agaactcctg 1380
ttgcctgcga acaagacaac tatactttca ggaacagtcg agatcgctag tggagaactg 1440
aaagattactg acaatgcggg tgtcaatggt cttggctctcg ctactcaggg ctcaggctcag 1500
cttaccctcgg gctctggagg aacctttagg ctggcaacac ccacgggagc acctgcgcct 1560
gttagacttta cgattggaaa gttagcattc gatccttttt ccttctcaaa aagagatttt 1620
gtttcagcat cagtaaatcg aggcacaaaa aacgtcaact taacaggagc tctggttatt 1680
gatgaacatg acgtttacaga tctttatgat atggtgtcat taooatctcc agtagcaatt 1740
ctatctcgctg ttttcaaaag agcaacogtt actaagacag gatttctcga tggggagatt 1800
gogactccaa gccactacgg ctaccaagga aagtgttctt acacatggtc ccgtccctcg 1860
ttaattccag ctctcgatgg aggtatttct ggagggtcct ctctcagcgc aaatactctc 1920
tatgctgtat ggaattcaga cactctcgct cgttctacct atattctaga tcccgagcgt 1980
tacggagaaa ttgtcagcaa cagcttatgg atttctctct taggaatatc gccattctct 2040
gatattctcc aagatgttct tttgatagat catcccggtt tgtccataac cgcgaaagct 2100
ttaggagcct atgtcgaaac cacaccaaga caaggacatg agggcttttc aggtcgctat 2160
ggaggctaac aagctgcgct atctatgaac tacacggacc acactacgtt aggactttct 2220
cttggcgagc tttatggaaa aactaacgcc aacccctacg attcagcttg ctacgaacaa 2280
atgtattttac tctcgttctt tgggtcaatt cctatcgtga ctcaaaagag cgaggcctta 2340
acttctcgtga aagcagctta tggttattcc aaaaatcacc taataaccac ctacctcaga 2400
cctgacaaaag ctccaaaatc tcaagggcaa tggcataaca atagttacta tgttctatt 2460
cttcgacaga atctcttctct aaactgggtt cttcttaaca gacctctggc tcaagcttgg 2520
gatctttcag gttttatttc cgcagaattc ctagggtggt ggcaaaagta gttcacagaa 2580
actggagatc tgcacagtag ctttagtaga ggttaaagggt acaatgttct cctacgcata 2640
ggatgttctt ctcaatgggt cacaccattt aagaaggctc ctctacact gaccatcaaa 2700
cttgcctaca agcctgatat ctatcgtgtc aacctcaca atattgtgac tgtcgtctca 2760
aaccaagaga gaccttcgat ctcaggagca aatctacgcc gccacgggtt gttttacaa 2820
atccatgatg tagtagatct caccggagac actcaggcct ttctaaacta taccttttag 2880
gggaaaaatg gatttcaaaa ccaccgagtg tctacaggac taaaatccac attttta 2937

```

<210> 50

<211> 801

<212> DNA

<213> Chlamydia pneumoniae

<400> 50

```

atgcattcaa aatttctttc tcgaagaaaa aaaaatagtt ctcataagga ggaacctctc 60
tgggatttga tagctccaag ttacaataag atagtccaag ataaaggcca ctactactat 120
agagaaacta tctctcccca actcctgcct tcaactcaact taggttcaaa aagttctgta 180
ttggtatttg gctgcggtca aggtttttta gaaggggccc tctctaagga atgtcgttat 240
ctaggcctag atatctcttc tagattgatt gctctagaca agaaaatgcy atcggtaaac 300
tctcatcagt ttaaggttgc agatcttlagc aaacgcttag agttcgtaga accgacatta 360
ttctctcagt cagtagcaat cctctccctt caaaatattg aattccccgg agagcgtata 420
cgtaatacag ctacgctcct cgaaccactc gggcaatttt ttatagtttt aaacctcctc 480
gtttttcgtg tctcatgggc atcatcctgg cactatgatg aaaaataaaa agctatctct 540
cgctatatag atcgttatct ctccccaatg aaaaatccca ctaatgctca cccaggacaa 600
aaagatttgc cttctacctc ctcccttcaac tttctcttaa gctatgtggt taagaagactg 660
tcttctcatg gattctttag ttacggtctt gaggaatgga catctcaaaa aacctcaaca 720
ggaaaacgag ctaaggcaga aaacctttgt cgaagggaat ttccattatt ccttatgatt 780
tcatgcatta agataaaaa a

```

<210> 51

<211> 252

<212> DNA

<213> Chlamydia pneumoniae

<400> 51

```

atgaacaac aacacaatcg taaggcttta tctcgcaaga ttggcacagt gaaaaaacaa 60
gccaaatttg gattgaagctt tttatagtag attaaaaaaa ttgaatgggt aagcaagac 120
gatcttaaga aatcacataa agtagttctt atcagtattt ttggttttgg atttgcctatt 180

```

tatttcgttag atcttctgtgtt gcgttaagtc atcacatgtt tagatggtag aacaaccttt 240
 ttgttcgtgtt aa 252

<210> 52

<211> 1185

<212> DNA

<213> Chlamydia pneumoniae

<400> 52

atgtcaaaa aaacttttca acgtaataag ccccatatca atattgggac gatcggggcac 60
 gtgaccatg gtaaaactac gctaacagcg gcaattacac gcgcgtctatc aggggatgga 120
 ttggcctctt tccgtgacta tagttcaatt gacaatactc cagaagaaaa ggctcgtgga 180
 attactatca acgcttctca cgttgaatac gaaccccaa atcgtcacta cgctcaagta 240
 gactgccttg gtcacgtgta ctatgtttaa aatatgatta caggcgccgc tcaaatggac 300
 ggagctatcc tagtctgttc agctacagac ggagctatgc cacaactcaa agaacatctc 360
 ttgctagctc gccaggttgg agttccttat atcgtttgtt tcttgaataa agtagatatg 420
 atctctcaag aagatgctga acttattgac cttgttgaga tggaaacttag tgagctctctt 480
 gaagaaaaag gctacaaaag atgcctctatt atccgttggtt ctgctttgaa agctctttgaa 540
 ggtgatgcaa attatatcga aaaagtctga gaacttatgc aagctgtgga tgacaacatc 600
 cctacacacg aaagagaaat tgataagcct ttcttaatgc ctatcgaaga cgtattctca 660
 atctctgttc tgggtactgt ggttacagga agaactcgag gtggaatcgt taaagtcttc 720
 gataaagttc agctcgtggg attaggagag actaaagaaa caatcgttac tggagtcgaa 780
 atgtctcaaga aagaactctc tgaaggtcgt gcaggagaaa acgttgggtt actcctcaga 840
 ggatattgaa agaactgatg tgaagaggtt atggttggtt gtcagcctaa cagcgtggaag 900
 cctcatcaga aatttaagtc agctgtttac gttctcaga aagaagaagg cggaactcgt 960
 aagcctttct tcagcggata cagacctcag ttcttctctc gtactacaga cgtgacagat 1020
 tgcgttaact tctcgaagg aactgaaatg gtaattgcct gagataaagt tgagcttgat 1080
 gttagagctca ttggaacagt tgcctttgaa gaaggaatga gatttgcact tcgtgaaggt 1140
 ggtgcactca tcggcgctgg aacgatttca aagatcaatg ctttaa 1185

<210> 53

<211> 1431

<212> DNA

<213> Chlamydia pneumoniae

<400> 53

atgagaatcg tacaagtcgc tgtagaattc actccaatcg ttaagtagg cggtctaggc 60
 gatcgtgtag ctagtctatc taaggagtta gcgaacaaa atgatgtgga agtactcttc 120
 cctcattatc ctttaatttc caaattctct tcgtctcaag ttcttctcga gggtctcttc 180
 tattatgaat ttittaggcaa gcagcaagcc tctgcaattt ctattcttta cgagggttct 240
 acgctctacta taattacgtt ggattccaaa atagagcttt tctcaaccac gtccgtgtac 300
 tctgagaata atgtgttagc ttctctgtct tttagcgtgc cagctgcagc ttactctcaa 360
 gaagcggatc ctgctgacat tgtgcacttg catgactggc atgtaggttt acttgcgggt 420
 ttattaaaaa accctttaa cctgtgcatt tgcgaagttg ttttactat ccatattt 480
 ggttatcgag ggtattgtag tacgcagcta ttgacagcgt cgcaaatgga tgatttcat 540
 ttgagctact accaactatt tcgcgatccg caaactctg tctaatgaa gggagctctc 600
 tattgtctgg attacattac gacagtgtct ctacttatg tgcaggaaat tataaacgac 660
 tattctgatt acgaacttca tgcgtgcatt ctgcaagaa attctgtatt ttctgggact 720
 atcaatggca ttgatgaaga cgttggaaac ccgaagacag atctcgtctt agctgtacag 780
 tacgatgcga gccctatta gcgaacctgac gttctcttta ctaaaaaga agagaacaga 840
 gcggtattat atgagaagtt ggggatcagt tcagactatt ttctcttgat ttctgtgatc 900
 tcaagcattg ttgaggaaaa ggttctgaa tttatgaaag agatattct ccatgctatg 960
 gagcacagtt atgcctttat cttgattggg acaagtcaaa atgaggttct tcttaatgag 1020
 ttccgttaact tacaagattg tttagcgagc tcccccaaca ttgctttgat tcttgggtat 1080
 aatgatctct tagccaggct aacttatgct gctgcgcata tgactgtcat cctctcaact 1140
 agggaggtct gtagcattac ccagctgata gcgattgctt atgagcagct tcttttatgt 1200
 cgtaaaaact gagggcttgc tgatacagtg attcctgggg taaatgggtt caactctctt 1260
 gatcaaaaa attttaatga atttcgggct atgcttagca acgctgtaac gacgtatcgt 1320
 caggagcctg acgtttgtgt gaatttgatt gagtcgggaa tgctctgggc ctctcgctta 1380
 gatgcactga ctaagactta cgtaaattct tatcaattct tactctcatg a 1431

<210> 54
 <211> 1041
 <212> DNA
 <213> Chlamydia pneumoniae

<400> 54
 atggaagcag atatttttaga tggaaagctc aaacgggttg aggtaagtaa aaaaggattg 60
 gtgaattgta atcaagtaga tgtcaatcag ctagtcccta toaagtataa atgggcttgg 120
 gaacattacc tcaatggatg tgcacaacac tgggtcccta ctgaagttcc tatggcaaga 180
 gatatacgagt tgtggaatcc agatgaactg tctgaagacg aacgcagggtt catttttgta 240
 aacctaggat ttctcagtag cgcggaaagc ctagtggaa ataacatogt tottgotatc 300
 ttcaaacata tcaacaaccc tgaagcaaga cagtatttacc tgcgtcaagc ttttgaggaa 360
 gcgtacata cctatacatt tctctatatt attgagctta aggaacttga tgaaggcgaa 420
 gtattcaatg atgtctctga tctcattttt tctgtacagt attgaggtta tcaaaatgaca 480
 ttaaacgtcg atgtctctga tctcattttt tctgtacagt attgaggtta tcaaaatgaca 540
 ttoattaaaa acitagttag atactatata attatggaa gaactctctt ctatgtgggt 600
 tttgttaata ttctctcttt cactagacaa aataaaatga caggaaattg agaacagtac 660
 caatacatcc tcagagatga saacatacat ttaaattttg gaatcgatct tatocataga 720
 attaaagaag aaaacccaga agtttggact acggaaactac aagaagaatc ogtogctctt 780
 attgaaaaag ctgtagagct tgaattggag taocgttaag attgottacc tgcaggaatc 840
 ttgggattaa gatottcgat gtttatagat taocgttgct atattgcaga tctgtgtta 900
 gagagaattg gggtgaagcc tatctatcac tcaagaatc otttccctg gatgagcaga 960
 acoatggatc tgaataaaga aaagaatttc tttgaaaccc gggttaccga ataccacac 1020
 gctggtaatt taagttggta a 1041

<210> 55
 <211> 3135
 <212> DNA
 <213> Chlamydia pneumoniae

<400> 55
 atggtcgaag ttgaagaaaa gcattacacc atcgtcaaac gtaatggaat gtttgtacca 60
 ttttaacaa atcggatttt ccaggcttgg gaggcagctt ttogagatac gcgtagctta 120
 gaaactagtt ctccactacc taaagactta gaagaatcta ttgcgcaaat tactcataaa 180
 gtcgtgaagg aagtctctgc taaaatttca gaaggtcagg tagtcaactg agagagaatc 240
 caggatcttg tagaaagta cgtctatatt agcgggttgc aggatgtgc tgcgattatc 300
 attgtttaca gggaccacag caaggcagag cgcggttaact ctctgtccat aattgccatc 360
 atacgtagag acgggggaag cgtcaaatat aatcctatga agatctctgc agctctcgaa 420
 aaagacttca gagcgagcgt ccaaattaat gggatgactc ctctgcacac actatccgaa 480
 attaatgacc taccctctag gatcgttgaa gatgtcctat gctctcatg ttgaagaagct 540
 attaatctgc aagagatcca agatatttgt gaaaagcaac ttatggttgc cggctattat 600
 gatgtggcca gaattatata tttatataga gaagctctgc cagcagcccg cgtcaataaa 660
 gatcaagatg gacaaagaaga gtttgtcccc caagaggaaa cgtacgttgt tcaaaaagaa 720
 gacgcccaca cctcctctct gagaaaaaca gatttagaaa agagggtttc ttgggcatgc 780
 aaacgcttcc ctaaaactac agattctcaa ctgcttcgag atatggcatt tatgaatttg 840
 tattcagaaa tcaaaagaaga cgaggtccac acagcatgca tcatggcggc agctgcacat 900
 atcagagagag aacctgatta cgtctttatc gcacgcagac tctccacgag ttctctgtat 960
 gaagagacct taggatgcag ctctcaagac ccaattttat cagaataaca taaaacaact 1020
 tttaaagaat acatctccaa tggagaagag tatcgttga abctccaatt aaaggattat 1080
 gatctcgatg cctcttagta agtctagac ctctctatga accaacagt ttctctatag 1140
 gggatccaaa atctctacga tctgtatttt aatctgaat aaggacgag tttagagact 1200
 gcgcagatct ttgttagtgc gttttctatg ggttagcgtt taaatgaag taatagacag 1260
 aatttttggg caatcacttt ctataactcg ttatccaact tctcgtatca cccagactat 1320
 ctccactgtt ttaactccgg aatgcgtcat tcccaactca tctcatgcta tctttccaca 1380
 gtaaaagatg acctaatgca catttataag gtgatttctg ataactgctt gctttctcaa 1440
 tggggcaggg gaattggaaa tgattggaaa gctgtctgtg ctacaggagc tgataactga 1500
 ggaaccaaag gaaagatgca aggcgtcttt cctctcatga aggttgcaca tgatactgca 1560
 atttgcagtg atcagggggg caaacgttaa ggtgctatgt gctgtatatt gtaaatga 1620
 cacttggatt acgaagaact tttagaattg cggaagaata caggagatga gcgtgtaga 1680
 actccagata tcaatacaga aagctggatt cctgatctct gttttaagag actagaaaaa 1740
 aaaggcatgt ggacactctt tagcccgat gatgtccag gtttaacaga agctctatgg 1800

```

ttagagtgttg aaaagcttta tgaagaatat gaacgtaagg ttgaatctgg ggaatccgt 1860
ctttataaaa aagtagaagc cgaagtgcgt tggcgtaaaa tgtaagcat ggtttacgaa 1920
acaggggcatc ttgggttac atttaaatgat ccttcgaata ttgcgtcaaa ccaagatcat 1980
gttggtcgctg tacgtgtctt taatctatgt acagagattt tattgaactg ttcggaatca 2040
gagaactcgag ttgttaattt aggttccata aacttggtag aacatatacg taatgacaag 2100
ttagatgaag aaaattataa agaaactatc tcaatagcca tccgtatttt ggataacgt 2160
attgaactga actctacoc tacaccagag gctaaacaag ccaactaac tcacagagct 2220
gtgggtgttg ggtgttatgg attccagatg gtcttttaag agttgaaact tagctatgcc 2280
tcacaagaag ctgtcgaatt ttctgaagag tgcctggaga tcatgcata ctacgtatct 2340
ctagctcgcg gottactcgc cttagatact atcgagcttc tcaaaagaac tcgctggaga 2400
tggggtcgtg ggtatctacc cttagatact atcgagcttc tcaaaagaac tcgctggaga 2460
cataatgttc ttgtagacac atcaagtaaa aaagattgga ctccagttcg tgatactatc 2520
cagaaatacg gaatgagaaa tagccaggtc atggcaattg ctctcaagc aacgatctcg 2580
aatatcatag ggtgcaccca atctatagag cccatgtata aacatctctt tgtaagctcc 2640
aaccttttcg gagagtttac gatccccaac acctacctga ttaaaaaact taaggaaata 2700
ggacttttgg atgcagaatg gttatgatgt ctataatatt ttgacggatc tctatttgaa 2760
attgaaagga tccctaatac attgaaaaag cttttcttta cggcatttga aatcgaaacc 2820
gagtgttata tagagtgtac ctctagaaga cagaaatgga ttgatattgg agtttctcta 2880
aatctgtatc ttgctgagcc agatggttaa aaactctcca atatgtatct acaggtctgg 2940
aaaaaaggat taaagactac ctattattta agatctcaag ctgcaacata agtagaaaa 3000
tcttttatag atatcaataa acggggcatt cggactcggt ggatgaaaaa taactcagcg 3060
tcacaaagta ttgtgtgcga aagaaaaaca accccgcttt gttcaatgga agaaggttgc 3120
gaatctgtgc aataa 3135

```

<210> 56

<211> 1386

<212> DNA

<213> Chlamydia pneumoniae

<400> 56

```

atgatgagct ctaagcgtao ctogaaaaata ggggtgcttt caatttttatt aacatttact 60
cactctatag ggttcgcaaa tgcgaattcg tccgtaggctc ttggcagcgt ctacattaca 120
tcogaggttgg taaagaagcc tcagaagaaga tcagaagaaga aacaagccaa aagaagacct 180
cgtgtctcga aaggatactt agtcctctct tcaaggactc ttccagctcg agcccaaaaag 240
atgaaaaact cctctgttaa agagtcttca ggtggttcta acgaaatttc tgcaaatct 300
acacccagct ctgtaaaaat acgaaagaac aaactgtcag aacnaaaggc agctcaaaaa 360
ggattttcag ctattttcaa cctaactttg aaaagcctac ttctaaactc tcttcaaaa 420
caaaaaactt caattcacga gagagaaaaa gcaactccta gatttgttaa tgagtctcag 480
cttagttccg cagcaaaaag ctactgcaca ccaactctcag ccgtctcttc cctattttta 540
gaaacagaaa tctgttcgagc tctgttagaa agaactaaag aactcaaga taatgaatt 600
catattctcg tagtgcaagt ccaaacgaac cccaagaac aaaaatacaa gacaaactaa 660
cagttggcat cccaagcctc gattcaacaa tctgaagaga ccgagcaact attgcyagaag 720
ctcgcccaag gtgctagact acctgtctta gtgcgtctta atccgtgaag gtctgtcaaa 780
agacaaaaag aagagttatt aaaagaactc ttagctgaac gttagacaatg taagaagaag 840
tctgtaagac aagctcttga agctgtctct ttaactaaga aagtgctgag aggcgggtct 900
gtgacctcga ctttacgata cgatccagaa aaagcggcgg aaatcaaaag tagacgcaat 960
tgcataagtaa gtctgaagc acgtgaacaa aaatattctat ctgtcaaaa agatgtctgc 1020
gctaatggga gactcgtacg caaactcct aaatctcagg ttgcagtaa tgctcagac 1140
tctacacgaa attctaaaaa tacaacata gatagctatc ttacagctaa ccaatacagc 1200
tgtagttctg aagaacacga ttggcctatg tcttctctgc ctctcaaaag cagaactcac 1260
aacagtatat ctgtatgtac catggtagtt actgtcatct cgtatgctgt agggggttgc 1320
attatagcta atgctacaga atctcaaaac acatcagatc caactctcc aactctact 1380
ccatag 1386

```

<210> 57

<211> 1731

<212> DNA

<213> Chlamydia pneumoniae

<400> 57

```

atgacagatt ttctactca cttcaaagga cccaaactta acccoattaa agtaaattcca 60
aacctttttt agaggaatcc taaagtogca agggtaactgc aaattacagc cgtagtcttta 120
ggaaatcattg cctcttatlc cgttatagta ctcattatag gcacccctct cggagctcct 180
ataagtatga tctctggcgg atgtctttta gctctggag gcgccttatt tgttgggtgt 240
acgattgcta cgtatttgca agctagaat agttataaga aggcgtgaa ccaaaagaaa 300
ctctcagagc ctttgatgga acgcccogaa ttgaagcct tagattatc cctagatctg 360
aaagaggtat gggacctaca tcaattctgt ttcaaacatc agacctgaat 420
ctttccaaaa cccaaaggga agttctaaat caaatcaaaa ttgatgatga gggaccctcc 480
ctaggggaat gcgcgctat gatttcagaa aactacgacg catgcttaaa gatgctgcg 540
tatcgtgagg agctcctgaa agaacaacc caataccaag agacacgatt caatcagaac 600
ctcactcata gaataaaagt ttgtctctcc atcctctcaa ggatocagga caatattct 660
aaagcggggc ggtcttttcc ttgaaattt tccacgttaa gctcgcggat gtcaagaa 720
cataccacca ccaactgtgat tctggcttta agtgccgttg ttctgtcat ggtcgtagca 780
gctctaattc caggtggcat tttagcacta cctatactt tggctgttg tattctgca 840
ggagtgtatt tcacggact ttctatcta gttctgaaga tttaaagtaa cccaagcgt 900
aatcgtcagg atttttataa agattttgta aaaaatgtag atatagagct tcttaacca 960
acggtaaactt tacagcagatt cctcttgtaa atgctcaaa ggttctgaa agagaagaa 1020
gaagctcctc tagaaggtca agatttggtat acacaataca taaccaatgc accatagaa 1080
aaaagatgta tcgaagagat cagagttacc cttagaaaa gagggtgcgt tcgagtctca gaccaaaaa 1140
atgaaagacg acttgagtt cttagaaaa gagggtgcgt cggggagact gtctgtagcg 1200
tcccgctcgg aagatccaa gaaactctct attttactc aagtaagga gtttgcaaa 1260
ttacgtcgcc aaaaactctca gaatatatcc acgatttatg gtccggacaa tgaaaattat 1320
gatcccgaa ttctcttacc ctggatgcct aaaaaagaa agaaatatga ccttagctta 1380
gaactcgtta caaagtgtga acccggtcca agagaagagt tgtgtgtgtg agagggggct 1440
aaaccaactc taagaagact caatatgaga attgcacttc tacaacaaca actatacagt 1500
gtccgaaat ggagacacc tcagggggaa cttacgggga atgttatcta ttacagatca 1560
gaactcgatc gtattcagat gctagaaagg gcattttata atcactcag ggaagctcaa 1620
gaggaataca ccaagctctc cggagacctt gttgacattc aaaaacgtat tttagggatc 1680
atagttgaag gggactcaga ttcaagaaca gaagaagagc ctcaggaata g 1731

```

<210> 58

<211> 1086

<212> DNA

<213> Chlamydia pneumoniae

<400> 58

```

atgcaacaaa ctgtaattgt agcaatgtca gggagcgtg attcttctgt cgttgcoatt 60
ttattccaaa aatttaccaa ttataaggtt attgguctct tcatgaagaa ttgggaagag 120
gatagcgaa ggcggccttg ctcgtctact aaagattatg aagatgtcga gaggttatgt 180
cttcagctcg atatccctta ttacacogta tcttttgcta aagaatatag agaaagatg 240
ttcgctcgtt tctcaagga atactcttta ggctacactc ctaaccocga cattctttgt 300
aacccagaaa tcaaaattga cctctacaaa aagaagcttg agaaacttgg cggagattac 360
ctcgtcatcag ggcactactg ccgattaaat acogagctcc aagaaaccca actccttaga 420
ggttgcgata ctcaaaagaa ttcagagctat tttttatcag gaactcctaa aagtgctct 480
cacaatgtgc tctttctctc tggggaaatg aataagactg aagttcgtgc gatgtcag 540
caagcagctc tctccacagc agaaaaaaaa gatagtacag gcaattgtct tatagggaa 600
cgccctttta aagagtctct agagaagttt cttcccaata aaacaggcaa cgttatcgt 660
tgggatacca aggaattgt agggcaacat caggagctc actattatc tatagggcag 720
cggcgaggag ttgatcttgg aggatccgag aaacccgtgt atgttgtggg aaaaaatata 780
gaggaataa cgtatttatat tgtgagggg gaagacacat cccagctcta cctacgggaa 840
ttaacagcta gagagctcaa ttgttttacc cctcotaat cgggatgtca cgttagcgt 900
aaagtccgct accgttctoc tgatgaagct tgaacgatag attatagctc aggtgacgag 960
gtcaaggtgc gattttcaaa acccgtcaag gcggttaact caggacaac aatagcgtt 1020
tatcaaggag atacctgct tggtagtgga gtatcgacg tctctatgat tccaagtga 1080
ggctag 1086

```

<210> 59

<211> 4830

<212> DNA

<213> Chlamydia pneumoniae

<400> 59

atggttagcga aaaaaacagt acgatctttat aggtcttctat ttctctcatc cgtaatagta 60
 cgaatatgtg cagcaggcat tgccttttgaa gcacattctct tacacagctc agaactagat 120
 ttagggtgat tcaataaaaa gtttgaggaaa cactctctctc atgttgtaaga ggctcaaaa 180
 tctgttttaa agggatcaga tctctgtaaat cctctcaga aagaatccga gaagttttg 240
 tacactcaag tgcctcttca ccaaggaaag cctggagaga gtttggatct cgcgcgtgct 300
 aattctctag agcattttca gcatcttttt agatactag aatttttccc aaccocctca agaacctgat 420
 acaagttaag ctgtaagtga gaaaatctcc tcagatacca aagagaatag aaaaagacct 480
 gacagtgaag atccttcaaa aaaaagtggc cttaaagaag ttctatcaga tctcctcaaa 540
 agtctctgaaa ctgcagtagc agctatttct gaagatcttg aaatctcaga aacatttca 600
 gcaagagatc ctcttcaggg tttagcattt tttataaaa atacattctc tcagttcttc 660
 tctgaaaagg attcttcatt tcaaggaaat atctttcttg gttcaggagc taattcaggg 720
 ctggtttttg aaaaatttaa ggccgcgaaa tctggggctg cagttttatc tgatcgaagt 780
 attgtttttg aaaaatttgt taaaggattg agttttatat ctgtgaaac tttagaagat 840
 ggctctgcgc cagggtgtaa catgttgttg acccatttgt gtgatgtaac tctcactgat 900
 tgtgcacgtg gtttagacat tgaagottta cgtctgggta aagatttttc tctgtgagga 960
 gctgttttca ctgctgacat ccatgaaagt caaaaatacc ttgcaggagt aattctatcc 1020
 gtttaggaga atcaaggagc ttgtgttgta gaaaaaacc gtgctgagaa gtccaatgga 1080
 ggagcttttg ctctcggaag tttgttttga agtaaacacc aaacaccgcg ctgtgtgaaa 1140
 gaaaatcaag cattaacagc agggacata tctctcagaa gtgatattga ttctcaagg 1200
 aactgtacgc ctattgaaat ttcaggaaac ttcagctctga ttgctctgag agacatata 1260
 gggtctacag atttttgtag tggaggagct ttgctctc agggagcgt tactcttaaga 1320
 aataatgcaag tagtgcaatg tgtlaaaaaa actctcaaaa cacttggtgg agctatttga 1380
 cgaagtaact ttgatctcaa cgaacaatt agcgaaagtg ctttttagca gaatacaga 1440
 gctctaacgt gagggtcatt aagtgcgaat gataagggtta taattgcaaa taacttttgg 1500
 gaaattcttt tttagcaaaa cgaagttagg aatcacggag gagcacttta ttgtgtagt 1560
 cgtcttaatc ctaagttaga acaaaaggat tctggagaga acatacaat ttgtgaaac 1620
 tccggagcta tcaatttttt aaaaaataag gctctgtttt tagaagtgat gacacaagct 1680
 gaagattatg ctggtggagg cgttttatgg ggccataatg ttctctaga ttccaatgtg 1740
 ggaattatc aatttatagg aaatataggt ggaagtacct tctggatagg agaattatgc 1800
 ggtggtgggt cgttctcttc tactgataga gtgacaattt ctaataaact tggagatgt 1860
 gtttttaag gaacaaaagg ccaatgtctt gctcaaaaat atgtgatctc tcaagaaaac 1920
 gctccgtggt aatcagatgc ttcatctaca aataaagacg agaagagcct taatgtctgt 1980
 agtcatggag atcattatcc tcttaaaact gtgaagaagg aagtgcocac ttcatgttta 2040
 gaagaacatc ctgttgtttc ttgcacagat attcgtgggt ttggggccat tctagctcaa 2100
 catattcttta ttacagataa tacaggaaat ctgagattct ctgggaacct tggttgtgt 2160
 gaagagctct ctactgcgg tgatttagct atcgtaggag gagggtcctt gctttctact 2220
 aatgaagtta atgtttgcag taacaaaaat gttgttttt ctgataacgt gacttcaaat 2280
 ggtgtgatt cagggggagc tatttttagct aaaaaagtag atatctcgc gacacactct 2340
 ttgtaatttg ttctaatgg ttccaggaaa tctgggtggt ccgttttgcc tttaaacgaa 2400
 tctgtaaaaa ttacgagcaa tggctcgcca gtatcattct ctaaaaatag aacacgtctt 2460
 ggcggtgctg gagtgcagc tctcaaaagg tctgtaacga ttgttgaaa cagggaaac 2520
 atagacttta aagagaactt tgtttttggc tctgaaaaatc aaagatcagg ttgagagct 2580
 atcatttgcta actctctgtg aaatattcaag gataacgcag gagatattct atttgaagt 2640
 aactctacgg gactcttagg aggtgctatt ttgttaggag ttgttagtgc atttggatg 2700
 agcaacccac gaacgcttac aattacaggc aacagtggtg atatctctt ttgtcaaaa 2760
 agcaacgaaa cagccgcttc ttatcagaa aagattctct ttgttgagag gccaactact 2820
 acacaaiaac tcaaaattgt aaagaatgca gggaacgttt ctctctatgg caacagagct 2880
 cctggtgtgct cgtgtgtcca aattgcagac gggaggaact ttgttttaga ggtcttttga 2940
 ggagtatctt tatttgaagg gaattatcaat ttgatggga gtttcaatgc gattoactta 3000
 tgcgggaatg actcaaaat cgtagagctt tctgtgtctc aagataaaaa atttatttc 3060
 caagtagcaa ttacttatga agagaacaca attctgtgct tccagataaa agatgtcagt 3120
 cctttaagtg ccoccttaatt aatttttaac tccaagccac gcttcaaacat 3180
 catgaaggga cgtacaggtt ttctcggagg gtatctaaaa tctctcagat tgcgtctata 3240
 caagagggaa ccttagcttt atcacaiaac gcagagcttt ggttggcagg gtttgaacag 3300
 gaaacaggaa gtcttatcgt attgtctcgg ggatctattc tctgtatttt tgatctccag 3360
 ttgtgatgca tgcgcctctc tctacagaa aataaaggag agactctgt tctctgcaga 3420
 gttcaaatta acatgagctc tctcaacccc aataaagata aagctgtaga taactcagta 3480

cttgcagata	tcataagtat	tactgtagat	ttgtcttcac	ttgttctgta	gcaagacgga	3540
actctctctc	ttctctctga	aattatcatt	cctaaggaaa	caaaatatac	ttcttaattg	3600
atagatctta	agattataga	ttctaccaat	gtgggatatg	aaaatcatgc	ttcttlaagt	3660
ttctcataag	atattccatt	aatttctctt	aagacagcgg	aaggaaatgc	aggagcgctc	3720
acagcagatg	ctctctctac	taataataaa	atagatgtat	ctttaccctc	gatcacacca	3780
gcaacgtatg	gtcacacagg	agtttggctt	gaaagtaaaa	tggaagatgg	aagactttga	3840
gtcggttggc	aacctacagg	atataagtta	aatcctgaga	agcaaggggc	cttagttttg	3900
aataatctct	ggagtoatat	tacagatctt	agagctctta	agcaggagat	ctttgtctac	3960
catacgatag	ctcaaaagat	ggagttagat	ttctcgacaa	atgtctgggg	ctcaggatta	4020
gggtgtgtgt	aagattgtca	gaacatcgga	gagtttgatg	gggtcaaaac	tcattctcaa	4080
gggtatggcc	taggcttggg	tacacaacta	gttgaagact	ttcttaattg	aggatgtttc	4140
tcacagttct	ttggtaaaac	tgaaagccaa	ttctacaaag	ctaagaacga	tgtgaagagt	4200
tatatgggag	ctgcttatgc	ggggatttta	gcaggctcct	ggttaataaa	aggagctttt	4260
ggttaacgta	atataaaaca	cgatttgact	acagatttac	gtacttttag	tatttcaaca	4320
ggttcatgga	taggaaaagg	gtttatcgca	ggcacaagca	ttgattacgc	ctattattga	4380
aatcctcgac	ggtttatata	ggcaatcgta	ttccacgtgg	ttccttttgt	agaagccgag	4440
tatgtctgac	tattatcttc	agaaatttag	gaacacggta	aagaggttag	aacgttccaa	4500
aaaactcggt	ttgagaatgt	cgccatttct	tttggatttg	ctttagaaca	tgcttattgc	4560
ttgtgctcac	gtcgtgaagt	gaacagtgtg	caagcttgctt	acgtctttga	tgtatatagt	4620
aaggagctgc	ttctcttgat	tacactcaag	gatgtctgct	attcttggaa	gagttatggg	4680
gtagatattc	cttgtaaagc	ttggaagagt	cgcttgagca	ataatacaga	atggaattca	4740
tatttaagta	ogtatttagc	gttttaattat	gaatggagag	aagatctgat	agcttatgac	4800
ttcaatgggt	gtatccgtat	tattttctag				4830

<210> 60

<211> 591

<212> DNA

<213> Chlamydia pneumoniae

<400> 60

atgacactct	ccotagtttg	aaaggaaagg	cctgattttg	ttgcgcaagc	tggtgttaat	60
ggcgaaaagc	gtaccgtatc	tttaaaaagt	tatttaggaa	agtagtttgt	gctttctctc	120
tatcotaaga	attttactta	cgtgtgtctc	acggaattgc	acgcatttca	agatgtotta	180
ggagaattcc	acaccggagg	agctgaagtc	ataggctgtt	cogtggatga	cattgocacc	240
catcaacagt	ggtttagctac	taagaaaaag	caaggttgta	tcgaaggtat	tacctatctc	300
cttctctcag	acgaagataa	agtcatttca	agaaagttat	atgtgttaaa	acccgaagaa	360
gaattattct	tcagaggagt	tttctcgatt	gataaagggt	gaatcatccg	tactctttga	420
gtgaatgac	ttctcttagc	cogtctctata	gaagaagaac	ttagaacctc	agatgtctta	480
atcttctttg	aaactaatgg	cttagtctgt	cctgcaaaat	ggcatgaagg	agagcgagcg	540
atggctccaa	atgaagaagg	actgcaaaat	tatttcggga	ctatagacta	g	591

<210> 61

<211> 1983

<212> DNA

<213> Chlamydia pneumoniae

<400> 61

atgagtgaa	acaaaaaatc	aagcaaaat	ataggatag	acttaggcac	aacaaactcc	60
ttgcttatct	ttatggaagg	aggacaagct	aaagttaata	catcatccga	aggacaagaa	120
accacgccat	cgatcgttgc	cttcaaaagt	aatgagaaat	tagtggggat	tcagacaaaa	180
cgtcaagcag	tgacaaatcc	agaaaaaact	ctcggctcta	caaaacgcgt	tattggcgct	240
aaagtactct	aaagtactct	ggaaatccaa	acgcttcttc	atcacgtcac	ctccgattct	300
aaaggtgatg	cogtttctga	agttgatggc	aaacaataca	ctccagaaga	aattggcgca	360
caaatcttaa	tgaaaatgaa	agagacagca	gaagctttat	taggcgaaac	ttgcacagaa	420
gcagtgatca	cogtcccccgc	atacttcaat	gatcttcaac	gagcatccac	aaaagatgct	480
ggacgcattg	caggtctaga	tgtaaaaacg	atcaattccag	aacctccgcg	agcagctctt	540
gcttaccgaa	togataaagt	cogtgateaa	aaaatcgctg	ttcttcgact	tggtggagga	600
accttttgata	ttctcatctc	agaaatcggt	gatggcgctc	tcgaagttct	actcacaaat	660
ggagatactc	ttctcgggtg	agacagactt	gatgaagtca	ttatcaaatg	gatgatcgaa	720

gaattcaaaa	aacaagaagg	cattgatctt	agcaaaagata	atatggcctt	acaaagactt	780
aaagatgctg	ctgagaagaag	aaaaatagaa	ctttcaggag	ttctotccac	agaaatcaat	840
cagcattcca	tcaacaatgga	tgcaacaagg	ctcaaacacc	ttgcattgac	actcaacagt	900
gcgcacattg	agaaactcgc	agcctctcta	atcgaaagaa	caaaactctc	atgcactaaa	960
gcactcagtg	gcgcaaaact	ttcogctaag	gatattcgatg	atgttctctt	agttggaagt	1020
atgtcgaagaa	tgccgcagtg	gcaagaacct	gtaaaagaac	ttcttggcaa	agagcctaatt	1080
aaagagctca	acoccgacga	agttgttgct	atggagagoc	caattcaagg	tggtgttctt	1140
ggcggaagag	ttaaagatgt	ttacttctta	gaogttatoc	ccctatctct	gggtatogaa	1200
actctaggag	gcgtcatgac	gaotctggta	gagagaaata	ctacaatccc	tacacagaaa	1260
aaacaaattct	ttctccacagc	tgctgataac	cagcctcgcg	ttaccatogt	agtttccaaa	1320
ggagagcgtc	ctactggccaa	agataacaag	gaaactcgaa	gattogattct	tacagatctc	1380
ctctcggctc	ctcagggcca	ttctcaaaat	gaagtctctc	togatatoga	tgcaaacagg	1440
attttccatg	ttctcagctaa	agatgttgcc	agcggtaaa	aacagaaaat	togtatogaa	1500
gcaagctcag	gaactcaaga	agatgaaatc	caaagattgg	ttcgagatgc	cgaaattaat	1560
aaggagaaga	ataaaaaaacg	togtgaagct	tcagatgcta	aaaatgaagc	cgatagcatg	1620
atcttcagag	cogaaaaaacg	tattaaagat	tataaggagc	aaattctcga	aaacttagtt	1680
aaagaaatcg	aagagcgcaat	cgaaaaocgtg	cgcaacgcac	tcaaaagatg	cgcttctatt	1740
gaaaaaatcta	aagaggctaac	tgaagacctta	agcaagcata	tgcaaaaaat	tgagagatct	1800
atgcaaatcgc	agctctgcac	agcaagcagca	ctactcggag	coaatcgtaa	agttggaact	1860
aacatcaata	cagaagatttt	gaaaaaaact	agttctcagta	cgaagcctcc	ttcaataaac	1920
ggttcttcag	aagaccatat	cgaagaagct	gatgtagaaa	tatttgataa	cgaagataag	1980
ttaa						1983

<210> 62

<211> 1860

<212> DNA

<213> *Chlamydia pneumoniae*

<400> 62

atgaaaaaag	ggaaattagg	agocatagtt	tttggccttc	tatttacaag	tagttgtgct	60
ggttttttcta	aggatttgac	taaagacaac	gcttatcaag	attttaaattg	catagagcat	120
tttaatatcgt	taaaatatgc	tcotttacca	tggaaggaa	tattattttgg	ttgggattta	180
ttctcagcaaa	cacagacaag	tcgcttgcaa	ctggctttag	aagaaaaacc	aacaaccaac	240
tactgcocaga	aggtactctc	taactacgtg	agatcattaa	acgattatca	tgacgggatt	300
acgtttttatc	tactcgaagag	tgogtatatc	cottacogtat	tgaagtttaag	tgaagatggg	360
catgtcttttg	tagtgcacgt	acagactagc	caaggggata	ttactttagg	gagatgaatc	420
cttgaagattg	atggaattggg	gattcgtgag	gctatcgaaa	gccttcgctt	tgacggaggg	480
agtgccacagc	actattctgc	tgcaagttctg	tccttgacat	cgcttccgcg	tttatttgga	540
gatgcgggttc	cttcaggaaat	tgccattgttg	aaacttcgcc	gaccagctgg	tttgatcogt	600
tcgacacagtc	ctctgttggcg	ttatactcca	gagcatatcg	gagatttttc	tttagtctgt	660
ccctttgattc	ctgaacataa	acctcaalta	cctacacaaa	gtttgtgtct	atctcgtttc	720
ggggttaaatt	cacagttctc	tagtagctct	ttattcagtt	cctacatggg	gccttatttc	780
tgggaagaagt	tgccgggttca	aaataagcag	cgttttgaca	gtaatcacca	tatagggagc	840
cgttaatggat	ttttacotac	gtttgtgtct	attotttggg	aacaagacaa	ggggcccat	900
cgttctctata	tttttaaaag	aaaagattct	cagggcaact	cccatcgcat	aggattttta	960
agaattttctt	cttatgtttg	gaotgattta	gaaggacttg	aagaggatca	taaggatagt	1020
cccttggggagc	cttttgggaa	gatcatcgat	catttggaaa	aagagactga	tgcttttgatt	1080
attgatcaga	ccataatcc	tggaggcagt	gtttttctct	ttctattogt	actatcctag	1140
ttaacagatc	atcctttaga	tactctctaaa	catagaatga	ttttcactca	ggatgaagtc	1200
agctcggctt	tgcaactggca	agatctacta	gaagatgtct	tcacagatga	gcaggcagtt	1260
gcgctgctgat	gggaagactat	ggaaggatat	tgcatggata	tgcatgctgt	agcctctctt	1320
caaaaactctg	ctcagagtg	cttttctctc	ttgggttccg	gtgatattaa	actttcaaaa	1380
ccctatgcttt	tgctaggatt	tgcaacaggt	cgacotcctc	ctaaaactca	atatactaaa	1440
cccttgcctta	ttgttgataga	caggatgac	ttctcttctg	gagatttagc	gccttgaatt	1500
ttgaagagata	atggccgcgc	tactctctaat	ggaaagccaa	cagcaggagc	tgagaggttt	1560
gtattccaaag	ctacttttcc	taaccgttct	gtatttaaag	ttctttctct	aaacagatct	1620
tttagctgtta	ggaaagatgg	tgagtttatt	gaaactctag	gagtggtctc	tcatattgat	1680
tttaggttta	ctccaggga	tttgoaaact	tccaggttta	ctgattcaogt	tgaggcagtg	1740
aaaactatag	ttttaactct	tttgtctgag	aacgctaaga	agagtgaaag	gcagactctc	1800

cgcgaagaga cgcotgaagt tattogagtc tcttatccca caacgaattc tgcttcgtaa 1860

<210> 63
 <211> 1956
 <212> DNA
 <213> Chlamydia pneumoniae

<400> 63
 atgggttaabc ctattgggtcc aggtcctata gacgaacacg aacgcacacc tcccgagat 60
 cttttcgtctc aaggatttga ggcgagtgcg gcaataaaga gtgcggaagc tcaagaata 120
 gcagggtcggg aagctaagcc taagaatctc aagaccgatt ctgtagagcg atggagcctc 180
 ttgcgtttctg cagtgaatgc tctcatgagt ctggcagata agctgggtat tgcttctagt 240
 aacagctcgtt cttctactag cagatctgca gacgtggact caacgcacgc gaccgcacct 300
 acgctctctc caccacagtt tgatgattat aagactcaag cgcacaacgc ttacgatact 360
 atctttacct caacatcact agctgacata caggctgott tggtagagct ccaggatgct 420
 gtaactaata taaaggatgc agcggctact gatgaggaaa ccgcaatcgc tgcggatggt 480
 gtaactaaga atgcogagtc agttaagtt ggccgcgaaa ttacgaattc agcgaatat 540
 gcttcggata accaagcgat tcttgactct ttaggtaaac tgacttctct cgactctcta 600
 caggctgctc tcttccaatc tgtagcaaac aatacaaaag cagctgagct tcttaaaag 660
 atgcagata acccagtagt cccagggaac aogcctgcac ttgctcaatc tttagttgat 720
 cagcagagat gacacagcag acagatagag aagatggaaa atgcgattag ggatgoatat 780
 ttgcaagagc agaacgctag tggagctgta gaaaatgcta aatctaataa cagataaagc 840
 aacatagatt cagctcaaac cgcactctcc aattcttcaa gaagcggaac aaatggtaat acaggtctag 900
 aaaaagttcc cgcactctcc aattcttcaa gaagcggaac aaatggtaat acaggtctag 960
 aaagattctta aazatatcaa acctgcagat ggtctgtag ttccaaatcc aggaactaca 1020
 gttagagctt ccaagcaaca aggaagtagt atgtgttagt tctgtgttcc catgtctta 1080
 gtatgctctg aaaaagagac cgtctctaat ttgatgtctg ggtttgtctc gatgattac 1140
 atgttcaata cgaataatcc tgattctcaa gctgcocaa cgcagctcgc agcacaagct 1200
 agagcagcga aagcgcctgg agatgacagt gctgctgcag cgttcgcaag tgcctagaana 1260
 gctttagaag cctcgttagg taagctggg caacaacagc gctactcaaa gcttttagta 1320
 cagatgcctt ctgctgtctg ttgagcgcca ggagttctcc cgcgtgcagc aagttctata 1380
 gggtctatctg taaaacagct ttacaagacc tcaaaatota caggttctga ttataaaca 1440
 cagatatacg caggttatga tgccttcaaa tcoataatg atgctctatg tagggcaga 1500
 aatgatgcga ctogtgatgt gataaacaat gtaagtaacc cgcctctcac acgatccgtt 1560
 cctagagcac gaacagaagc tcgaggacca gaaaaaacag atcaagccct cgttaggggt 1620
 atttctggca atagcagaac tcttgagat gtctatagtc aagtttcgca acataatct 1680
 gtaatgcaga tcatccagtc gaatctctca gcgaataatg aggagatcag acaaaagcct 1740
 acatcgcgaag tgacaagaac tccacagttt ggctatcctt atgtgcaact tcttaatgac 1800
 tctacacaga agttcatagc taaattagaa agtttgggtt ctgaagatc taggacagca 1860
 gctgaataaa aagcacttcc ctttgaacg aactcctgtt ttattcagca ggtgctgctc 1920
 aatatcgctt ctctatatcc tggttatctc caataa 1956

<210> 64
 <211> 264
 <212> DNA
 <213> Chlamydia pneumoniae

<400> 64
 atgagtcaaa aaaataaaaa ctctgctttt atgcctcccg tgaatatctc cacagattta 60
 gcagttatag ttggcaaggg aactatgcct agaacogaaa ttgtaaaaga agtttgggaa 120
 tacattaaaa aacacaaactg taaggatcaa aaaaataaac gtaatatcct tccgatgcgc 180
 aatcttgcca aagttctttg ctctagtgt atctatgaca tgtttccaaat gaccaaaagc 240
 ctttccaaac atattgtaaa ataa 264

<210> 65
 <211> 978
 <212> PRT
 <213> Chlamydia pneumoniae

```

<400> 65
Met Pro Leu Ser Phe Lys Ser Ser Ser Phe Cys Leu Leu Ala Cys Leu
                    5
Cys Ser Ala Ser Cys Ala Phe Ala Glu Thr Arg Leu Gly Gly Asn Phe
                    20
Val Pro Pro Ile Thr Asn Gln Gly Glu Glu Ile Leu Leu Thr Ser Asp
                    35
Phe Val Cys Ser Asn Phe Leu Gly Ala Ser Phe Ser Ser Ser Phe Ile
                    50
Asn Ser Ser Ser Asn Leu Ser Leu Leu Gly Lys Gly Leu Ser Leu Thr
                    65
Phe Thr Ser Cys Gln Ala Pro Thr Asn Ser Asn Tyr Ala Leu Leu Ser
                    85
Ala Ala Glu Thr Leu Thr Phe Lys Asn Phe Ser Ser Ile Asn Phe Thr
                    100
Gly Asn Gln Ser Thr Gly Leu Gly Gly Leu Ile Tyr Gly Lys Asp Ile
                    115
Val Phe Gln Ser Ile Lys Asp Leu Ile Phe Thr Thr Asn Arg Val Ala
                    130
Tyr Ser Pro Ala Ser Val Thr Thr Ser Ala Thr Pro Ala Ile Thr Thr
                    145
Val Thr Thr Gly Ala Ser Ala Leu Gln Pro Thr Asp Ser Leu Thr Val
                    165
Glu Asn Ile Ser Gln Ser Ile Lys Phe Phe Gly Asn Leu Ala Asn Phe
                    180
Gly Ser Ala Ile Ser Ser Ser Pro Thr Ala Val Val Lys Phe Ile Asn
                    195
Asn Thr Ala Thr Met Ser Phe Ser His Asn Phe Thr Ser Ser Gly Gly
                    210
Gly Val Ile Tyr Gly Gly Ser Ser Leu Leu Phe Glu Asn Asn Ser Gly
                    225
Cys Ile Ile Phe Thr Ala Asn Ser Cys Val Asn Ser Leu Lys Gly Val
                    245
Thr Pro Ser Ser Gly Thr Tyr Ala Leu Gly Ser Gly Gly Ala Ile Cys
                    260
Ile Pro Thr Gly Thr Phe Glu Leu Lys Asn Asn Gln Gly Lys Cys Thr
                    275
Phe Ser Tyr Asn Gly Thr Pro Asn Asp Ala Gly Ala Ile Tyr Ala Glu
                    290
Thr Cys Asn Ile Val Gly Asn Gln Gly Ala Leu Leu Leu Asp Ser Asn
                    305
Thr Ala Ala Arg Asn Gly Gly Ala Ile Cys Ala Lys Val Leu Asn Ile
                    325
Gln Gly Arg Gly Pro Ile Glu Phe Ser Arg Asn Arg Ala Glu Lys Gly
                    340
Gly Ala Ile Phe Ile Gly Pro Ser Val Gly Asp Pro Ala Lys Gln Thr
                    355
Ser Thr Leu Thr Ile Leu Ala Ser Glu Gly Asp Ile Ala Phe Gln Gly
                    370
Asn Met Leu Asn Thr Lys Pro Gly Ile Arg Asn Ala Ile Thr Val Glu
                    385
Ala Gly Gly Glu Ile Val Ser Leu Ser Ala Gln Gly Gly Ser Arg Leu
                    405
Val Phe Tyr Asp Pro Ile Thr His Ser Leu Pro Thr Thr Ser Pro Ser
                    420
Asn Lys Asp Ile Thr Ile Asn Ala Asn Gly Ala Ser Gly Ser Val Val
                    435
Phe Thr Ser Lys Gly Leu Ser Ser Thr Glu Leu Leu Leu Pro Ala Asn
                    450
Thr Thr Thr Ile Leu Leu Gly Thr Val Lys Ile Ala Ser Gly Glu Leu

```


465	Lys	Ile	Thr	Asp	Asn	Ala	Val	Val	Asn	Val	Leu	Gly	Phe	Ala	Thr	Gln
					485					490						495
Thr	Ser	Gly	Gln	Leu	Thr	Leu	Gly	Ser	Gly	Gly	Thr	Leu	Gly	Leu	Ala	
				500				505							510	
Gly	Pro	Thr	Gly	Gly	Ala	Ala	Val	Asp	Phe	Thr	Ile	Gly	Lys	Leu		
				515			520					525				
Ala	Phe	Asp	Pro	Phe	Ser	Phe	Leu	Lys	Arg	Asp	Phe	Val	Ser	Ala	Ser	
				530			535				540					
Val	Asn	Ala	Gly	Thr	Lys	Asn	Val	Thr	Leu	Thr	Gly	Ala	Leu	Val	Leu	
				545			550				555					560
Asp	Glu	His	Asp	Val	Thr	Asp	Leu	Tyr	Asp	Met	Val	Ser	Leu	Gln	Ser	
				565				570								575
Pro	Val	Ala	Ile	Pro	Ile	Ala	Val	Phe	Lys	Gly	Ala	Thr	Val	Thr	Lys	
				580				585								590
Thr	Gly	Phe	Pro	Asp	Gly	Glu	Ile	Ala	Thr	Pro	Ser	His	Tyr	Gly	Tyr	
							600									
Gln	Gly	Lys	Trp	Ser	Tyr	Thr	Trp	Ser	Arg	Pro	Leu	Leu	Ile	Pro	Ala	
							615									
Pro	Asp	Gly	Gly	Phe	Pro	Gly	Gly	Pro	Ser	Pro	Ser	Ala	Asn	Thr	Leu	
							630				635					640
Tyr	Ala	Val	Trp	Asn	Ser	Asp	Thr	Leu	Val	Arg	Ser	Thr	Tyr	Ile	Leu	
				645					650							655
Asp	Pro	Glu	Arg	Tyr	Gly	Glu	Ile	Val	Ser	Asn	Ser	Leu	Trp	Ile	Ser	
				660				665								
Phe	Leu	Gly	Asn	Gln	Ala	Phe	Ser	Asp	Ile	Leu	Gln	Asp	Val	Leu	Leu	
				675			680					685				
Ile	Asp	His	Pro	Gly	Leu	Ser	Ile	Thr	Ala	Lys	Ala	Leu	Gly	Ala	Tyr	
							695					700				
Val	Glu	His	Thr	Pro	Arg	Gln	Gly	His	Glu	Gly	Phe	Ser	Gly	Arg	Tyr	
							710				715					720
Gly	Gly	Tyr	Gln	Ala	Ala	Leu	Ser	Met	Asn	Tyr	Thr	Asp	His	Thr	Thr	
				725						730						
Leu	Gly	Leu	Ser	Phe	Gly	Gln	Leu	Tyr	Gly	Lys	Thr	Asn	Ala	Asn	Pro	
				740					745							
Tyr	Asp	Ser	Arg	Cys	Ser	Glu	Gln	Met	Tyr	Leu	Leu	Ser	Ser	Phe	Phe	Gly
				755				760						765		
Gln	Phe	Pro	Ile	Val	Thr	Gln	Lys	Ser	Glu	Ala	Leu	Ile	Ser	Trp	Lys	
							775									
Ala	Ala	Tyr	Gly	Tyr	Ser	Lys	Asn	His	Leu	Asn	Thr	Thr	Tyr	Leu	Arg	
							790				795					800
Pro	Asp	Lys	Ala	Pro	Lys	Ser	Gln	Gly	Gln	Trp	His	Asn	Asn	Ser	Tyr	
				805					810							815
Tyr	Val	Leu	Ile	Ser	Ala	Glu	His	Pro	Phe	Leu	Asn					

Gly Lys Asn Gly Phe Thr Asn His Arg Val Ser Thr Gly Leu Lys Ser
 965 970 975
 Thr Phe

<210> 66
 <211> 266
 <212> PRT
 <213> Chlamydia pneumoniae

<400> 66
 Met His Ser Lys Phe Leu Ser Arg Arg Lys Lys Asn Ser Ser His Lys
 5 10 15
 Glu Glu Thr Ser Trp Asp Cys Ile Ala Ser Ser Tyr Asn Lys Ile Val
 20 25 30
 Gln Asp Lys Gly His Tyr Tyr His Arg Glu Thr Ile Leu Pro Gln Leu
 35 40 45
 Leu Pro Ser Leu Thr Leu Gly Ser Lys Ser Ser Val Leu Asp Ile Gly
 50 55 60
 Cys Gly Gln Gly Phe Leu Glu Arg Ala Leu Pro Lys Glu Cys Arg Tyr
 65 70 75 80
 Leu Gly Ile Asp Ile Ser Ser Arg Leu Ile Ala Leu Ala Lys Lys Met
 85 90 95
 Arg Ser Val Asn Ser His Gln Phe Lys Val Ala Asp Leu Ser Lys Arg
 100 105 110
 Leu Glu Phe Val Glu Pro Thr Leu Phe Ser His Ala Val Ala Ile Leu
 115 120 125
 Ser Leu Gln Asn Met Glu Phe Pro Gly Glu Ala Ile Arg Asn Thr Ala
 130 135 140
 Thr Leu Leu Glu Pro Leu Gly Gln Phe Phe Ile Val Leu Asn His Pro
 145 150 155 160
 Cys Phe Arg Ile Pro Arg Ala Ser Ser Trp His Tyr Asp Glu Asn Lys
 165 170 175
 Lys Ala Ile Ser Arg His Ile Asp Arg Tyr Leu Ser Pro Met Lys Ile
 180 185 190
 Pro Ile Met Ala His Pro Gly Gln Lys Asp Ser Pro Ser Thr Leu Ser
 195 200 205
 Phe His Phe Pro Leu Ser Tyr Trp Phe Lys Glu Leu Ser Ser His Gly
 210 215 220
 Phe Leu Val Ser Gly Leu Glu Glu Trp Thr Ser Ser Lys Thr Ser Thr
 225 230 235 240
 Gly Lys Arg Ala Lys Ala Glu Asn Leu Cys Arg Lys Glu Phe Pro Leu
 245 250 255
 Phe Leu Met Ile Ser Cys Ile Lys Ile Lys
 260 265

<210> 67
 <211> 83
 <212> PRT
 <213> Chlamydia pneumoniae

<400> 67
 Met Lys Lys Gln Gln His Asn Arg Lys Ala Leu Ser Arg Lys Ile Gly Thr
 5 10 15
 Val Lys Lys Gln Ala Lys Phe Ala Gly Ser Phe Leu Asp Glu Ile Lys
 20 25 30
 Lys Ile Glu Trp Val Ser Lys His Asp Leu Lys Lys Tyr Ile Lys Val
 35 40 45
 Val Leu Ile Ser Ile Phe Gly Phe Gly Phe Ala Ile Tyr Phe Val Asp
 50 55 60

Leu Val Leu Arg Lys Ser Ile Thr Cys Leu Asp Gly Ile Thr Thr Phe
 65 70 75 80
 Leu Phe Gly

<210> 68
 <211> 394
 <212> PRT
 <213> Chlamydia pneumoniae

<400> 68
 Met Ser Lys Glu Thr Phe Gln Arg Asn Lys Pro His Ile Asn Ile Gly
 5 10 15
 Thr Ile Gly His Val Asp His Gly Lys Thr Thr Leu Thr Ala Ala Ile
 20 25 30
 Thr Arg Ala Leu Ser Gly Asp Gly Leu Ala Ser Phe Arg Asp Tyr Ser
 35 40 45
 Ser Ile Asp Asn Thr Pro Glu Lys Ala Arg Gly Ile Thr Ile Asn
 50 55 60
 Ala Ser His Val Glu Tyr Glu Thr Pro Asn Arg His Tyr Ala His Val
 65 70 75 80
 Asp Cys Pro Gly His Ala Asp Tyr Val Lys Asn Met Ile Thr Gly Ala
 85 90 95
 Ala Gln Met Asp Gly Ala Ile Leu Val Val Ser Ala Thr Asp Gly Ala
 100 105 110
 Met Pro Gln Thr Lys Glu His Ile Leu Leu Ala Arg Gln Val Gly Val
 115 120 125
 Pro Tyr Ile Val Val Phe Leu Asn Lys Val Asp Met Ile Ser Gln Glu
 130 135 140
 Asp Ala Glu Leu Ile Asp Leu Val Glu Met Glu Leu Ser Glu Leu Leu
 145 150 155 160
 Glu Glu Lys Gly Tyr Lys Gly Cys Pro Ile Ile Arg Gly Ser Ala Leu
 165 170 175
 Lys Ala Leu Glu Gly Asp Ala Asn Tyr Ile Glu Lys Val Arg Glu Leu
 180 185 190
 Met Gln Ala Val Asp Asp Asn Ile Pro Thr Pro Glu Arg Glu Ile Asp
 195 200 205
 Lys Pro Phe Leu Met Pro Ile Glu Asp Val Phe Ser Ile Ser Gly Arg
 210 215 220
 Gly Thr Val Val Thr Gly Arg Ile Glu Arg Gly Ile Val Lys Val Ser
 225 230 235 240
 Asp Lys Val Gln Leu Val Gly Leu Gly Glu Thr Lys Glu Thr Ile Val
 245 250 255
 Thr Gly Val Glu Met Phe Arg Lys Glu Leu Pro Glu Gly Arg Ala Gly
 260 265 270
 Glu Asn Val Gly Leu Leu Leu Arg Gly Ile Gly Lys Asn Asp Val Glu
 275 280 285
 Arg Gly Met Val Val Cys Gln Pro Asn Ser Val Lys Pro His Thr Lys
 290 295 300
 Phe Lys Ser Ala Val Tyr Val Leu Gln Lys Glu Glu Gly Gly Arg His
 305 310 315 320
 Lys Pro Phe Phe Ser Gly Tyr Arg Pro Gln Phe Phe Arg Thr Thr
 325 330 335
 Asp Val Thr Gly Val Val Thr Leu Pro Glu Gly Thr Glu Met Val Met
 340 345 350
 Pro Gly Asp Asn Val Glu Leu Asp Val Glu Leu Ile Gly Thr Val Ala
 355 360 365
 Leu Glu Glu Gly Met Arg Phe Ala Ile Arg Glu Gly Gly Arg Thr Ile
 370 375 380
 Gly Ala Gly Thr Ile Ser Lys Ile Asn Ala
 385 390

```
<210> 69
<211> 476
<212> PRT
<213> Chlamydia pneumoniae
```

[illegible]

420 425 430
 Ser Asn Ala Val Thr Thr Tyr Arg Gln Glu Pro Asp Val Trp Leu Asn
 435 440 445
 Leu Ile Glu Ser Gly Met Leu Arg Ala Ser Gly Leu Asp Ala Met Ala
 450 455 460
 Lys His Tyr Val Asn Leu Tyr Gln Ser Leu Leu Ser
 465 470 475

<210> 70
 <211> 346
 <212> PRT
 <213> Chlamydia pneumoniae

<400> 70
 Met Glu Ala Asp Ile Leu Asp Gly Lys Leu Lys Arg Val Glu Val Ser
 5 10 15
 Lys Lys Gly Leu Val Asn Cys Asn Gln Val Asp Val Asn Gln Leu Val
 20 25 30
 Pro Ile Lys Tyr Lys Trp Ala Trp Glu His Tyr Leu Asn Gly Cys Ala
 35 40 45
 Asn Asn Trp Leu Pro Thr Glu Val Pro Met Ala Arg Asp Ile Glu Leu
 50 55 60
 Trp Lys Ser Asp Glu Leu Ser Glu Asp Glu Arg Arg Val Ile Leu Leu
 65 70 75 80
 Asn Leu Gly Phe Phe Ser Thr Ala Glu Ser Leu Val Gly Asn Asn Ile
 85 90 95
 Val Leu Ala Ile Phe Lys His Ile Thr Asn Pro Glu Ala Arg Gln Tyr
 100 105 110
 Leu Leu Arg Gln Ala Phe Glu Glu Ala Val His Thr His Thr Phe Leu
 115 120 125
 Tyr Ile Cys Glu Ser Leu Gly Leu Asp Glu Gly Glu Val Phe Asn Ala
 130 135 140
 Tyr Asn Glu Arg Ala Ser Ile Arg Ala Lys Asp Asp Phe Gln Met Thr
 145 150 155 160
 Leu Thr Val Asp Val Leu Asp Pro Asn Phe Ser Val Gln Ser Ser Glu
 165 170 175
 Gly Leu Gly Gln Phe Ile Lys Asn Leu Val Gly Tyr Tyr Ile Ile Met
 180 185 190
 Glu Gly Ile Phe Phe Tyr Ser Gly Phe Val Met Ile Leu Ser Phe His
 195 200 205
 Arg Gln Asn Lys Met Thr Gly Ile Gly Glu Gln Tyr Gln Tyr Ile Leu
 210 215 220
 Arg Asp Glu Thr Ile His Leu Asn Phe Gly Ile Asp Leu Ile Asn Gly
 225 230 235 240
 Ile Lys Glu Glu Asn Pro Glu Val Trp Thr Thr Glu Leu Gln Glu Glu
 245 250 255
 Ile Val Ala Leu Ile Glu Lys Ala Val Glu Leu Glu Ile Glu Tyr Ala
 260 265 270
 Lys Asp Cys Leu Pro Arg Gly Ile Leu Gly Leu Arg Ser Ser Met Phe
 275 280 285
 Ile Asp Tyr Val Arg His Ile Ala Asp Arg Arg Leu Glu Arg Ile Gly
 290 295 300
 Leu Lys Pro Ile Tyr His Ser Arg Asn Pro Phe Pro Trp Met Ser Glu
 305 310 315 320
 Thr Met Asp Leu Asn Lys Glu Lys Asn Phe Phe Glu Thr Arg Val Thr
 325 330 335
 Glu Tyr Gln Thr Ala Gly Asn Leu Ser Trp
 340 345

```
<210> 71
<211> 1044
<212> PRT
<213> Chlamydia pneumoniae
```

4000> 71	Met	Val	Glu	Val	Glu	Glu	Lys	His	Tyr	Thr	Ile	Val	Lys	Arg	Asn	Gly
					5					10					15	
Met	Phe	Val	Pro	Phe	Asn	Gln	Asp	Leu	Arg	Ile	Phe	Gln	Ala	Leu	Glu	Ala
			20					25						30		
Ala	Phe	Arg	Asp	Thr	Arg	Ser	Leu	Glu	Thr	Ser	Ser	Pro	Leu	Pro	Lys	
			35				40						45			
Asp	Leu	Glu	Glu	Ser	Ile	Ala	Gln	Ile	Thr	His	Lys	Val	Val	Lys	Glu	
			50			55					60					
Val	Leu	Ala	Lys	Ile	Ser	Glu	Gly	Gln	Val	Val	Thr	Val	Glu	Arg	Ile	
			65			70				75					80	
Gln	Asp	Leu	Val	Glu	Ser	Gln	Leu	Tyr	Ile	Ser	Gly	Leu	Gln	Asp	Val	
				85					90					95		
Ala	Arg	Asp	Tyr	Ile	Val	Tyr	Arg	Asp	Gln	Arg	Lys	Ala	Glu	Arg	Gly	
			100					105						110		
Asn	Ser	Ser	Ser	Ile	Ile	Ala	Ile	Ile	Arg	Arg	Asp	Gly	Gly	Ser	Ala	
			115				120					125				
Lys	Phe	Asn	Pro	Met	Lys	Ile	Ser	Ala	Ala	Leu	Glu	Lys	Ala	Phe	Arg	
			130			135					140					
Ala	Thr	Leu	Gln	Ile	Asn	Gly	Met	Thr	Pro	Pro	Ala	Thr	Leu	Ser	Arg	
			145		150					155					160	
Ile	Asn	Asp	Leu	Thr	Leu	Arg	Ile	Val	Glu	Asp	Val	Leu	Ser	Leu	His	
			165					170						175		
Gln	Glu	Glu	Ala	Ile	Asn	Leu	Glu	Glu	Ile	Gln	Asp	Ile	Val	Glu	Lys	
			180				185						190			
Gln	Leu	Met	Val	Ala	Gly	Tyr	Tyr	Asp	Val	Ala	Lys	Asn	Tyr	Ile	Leu	
			195			200						205				
Tyr	Arg	Glu	Ala	Arg	Ala	Arg	Ala	Arg	Ala	Asn	Lys	Asp	Gln	Asp	Gly	
			210		215					220						
Gln	Glu	Glu	Phe	Val	Pro	Gln	Glu	Glu	Thr	Thr	Val	Val	Gln	Lys	Glu	
			225		230				235					240		
Asp	Gly	Thr	Thr	Tyr	Leu	Leu	Arg	Lys	Thr	Asp	Leu	Glu	Lys	Arg	Phe	
			245					250					255			
Ser	Trp	Ala	Cys	Lys	Arg	Phe	Pro	Lys	Thr	Thr	Asp	Ser	Gln	Leu	Leu	
			260				265						270			
Ala	Asp	Met	Ala	Phe	Met	Asn	Leu	Tyr	Ser	Gly	Ile	Lys	Glu	Asp	Glu	
			275				280					285				
Val	Thr	Thr	Ala	Cys	Ile	Met	Ala	Ala	Arg	Ala	Asn	Ile	Glu	Arg	Glu	
			290			295					300					
Pro	Asp	Tyr	Ala	Phe	Ile	Ala	Ala	Glu	Leu	Leu	Thr	Ser	Ser	Leu	Tyr	
			305		310				315						320	
Glu	Glu	Thr	Leu	Gly	Cys	Ser	Ser	Gln	Asp	Pro	Asn	Leu	Ser	Glu	Ile	
			325						330					335		
His	Lys	Lys	His	Phe	Lys	Glu										

[illegible]

Lys Lys Leu Phe Leu Thr Ala Phe Glu Ile Glu Pro Glu Trp Ile Ile
 930 935 940
 Glu Cys Thr Ser Arg Arg Gln Lys Trp Ile Asp Met Gly Val Ser Leu
 945 950 955 960
 Asn Leu Tyr Leu Ala Glu Pro Asp Gly Lys Lys Leu Ser Asn Met Tyr
 965 970 975
 Leu Thr Ala Trp Lys Lys Gly Leu Lys Thr Thr Tyr Tyr Leu Arg Ser
 980 985 990
 Gln Ala Ala Thr Ser Val Glu Lys Ser Phe Ile Asp Ile Asn Lys Arg
 995 1000 1005
 Gly Ile Gln Pro Arg Trp Met Lys Asn Lys Ser Ala Ser Thr Ser Ile
 1010 1015 1020
 Val Val Glu Arg Lys Thr Thr Pro Val Cys Ser Met Glu Glu Gly Cys
 1025 1030 1035 1040
 Glu Ser Cys Gln

<210> 72

<211> 461

<212> PRT

<213> Chlamydia pneumoniae

<400> 72

Met Met Ser Ser Lys Arg Thr Ser Lys Ile Ala Val Leu Ser Ile Leu
 5 10 15
 Leu Thr Phe Thr His Ser Ile Gly Phe Ala Asn Ala Asn Ser Ser Val
 20 25 30
 Gly Leu Gly Thr Val Tyr Ile Thr Ser Glu Val Val Lys Lys Pro Gln
 35 40 45
 Lys Gly Ser Glu Arg Lys Gln Ala Lys Lys Glu Pro Arg Ala Arg Lys
 50 55 60
 Gly Tyr Leu Val Pro Ser Ser Arg Thr Leu Ser Ala Arg Ala Gln Lys
 65 70 75 80
 Met Lys Asn Ser Ser Arg Lys Glu Ser Ser Gly Gly Cys Asn Glu Ile
 85 90 95
 Ser Ala Asn Ser Thr Pro Arg Ser Val Lys Leu Arg Arg Asn Lys Arg
 100 105 110
 Ala Glu Gln Lys Ala Ala Lys Gln Gly Phe Ser Ala Phe Ser Asn Leu
 115 120 125
 Thr Leu Lys Ser Leu Leu Pro Lys Leu Pro Ser Lys Gln Lys Thr Ser
 130 135 140
 Ile His Glu Arg Glu Lys Ala Thr Ser Arg Phe Val Asn Glu Ser Gln
 145 150 155 160
 Leu Ser Ser Ala Arg Lys Arg Tyr Cys Thr Pro Ser Ser Ala Ala Pro
 165 170 175
 Ser Leu Phe Leu Glu Thr Glu Ile Val Arg Ala Pro Val Glu Arg Thr
 180 185 190
 Lys Glu Leu Gln Asp Asn Glu Ile His Ile Pro Val Val Gln Val Gln
 195 200 205
 Thr Asn Pro Lys Glu Gln Asn Thr Lys Thr Thr Lys Gln Leu Ala Ser
 210 215 220
 Gln Ala Ser Ile Gln Gln Ser Glu Gly Thr Glu Gln Ser Leu Arg Glu
 225 230 235 240
 Leu Ala Gln Gly Ala Ser Leu Pro Val Leu Val Arg Ser Asn Pro Glu
 245 250 255
 Val Ser Val Gln Arg Gln Lys Glu Glu Leu Lys Glu Leu Val Ala
 260 265 270
 Glu Arg Arg Gln Cys Lys Arg Lys Ser Val Arg Gln Ala Leu Glu Ala
 275 280 285
 Arg Ser Leu Thr Lys Lys Val Ala Arg Gly Gly Ser Val Thr Ser Thr


```

      290              295              300
Leu Arg Tyr Asp Pro Glu Lys Ala Ala Glu Ile Lys Ser Arg Arg Asn
305              310              315
Cys Lys Val Ser Pro Glu Ala Arg Glu Gln Lys Tyr Ser Ser Cys Lys
      325              330              335
Arg Asp Ala Arg Ala Asn Gly Lys Gln Asp Lys Thr Thr Pro Ser Glu
      340              345              350
Asp Ala Ser Gln Glu Glu Gln Gln Thr Gly Ala Gly Leu Val Arg Lys
      355              360              365
Thr Pro Lys Ser Gln Val Ala Ser Asn Ala Gln Asn Phe Tyr Arg Asn
      370              375              380
Ser Lys Asn Thr Asn Ile Asp Ser Tyr Leu Thr Ala Asn Gln Tyr Ser
385              390              395
Cys Ser Ser Glu Glu Thr Asp Trp Pro Cys Ser Ser Cys Val Ser Lys
      405              410              415
Arg Arg Thr His Asn Ser Ile Ser Val Cys Thr Met Val Val Thr Val
      420              425              430
Ile Ala Met Ile Val Gly Ala Leu Ile Ile Ala Asn Ala Thr Glu Ser
      435              440              445
Gln Thr Thr Ser Asp Pro Thr Pro Pro Thr Pro Thr Pro
      450              455              460

```

<210> 73

<211> 576

<212> PRT

<213> Chlamydia pneumoniae

<400> 73

```

Met Thr Asp Phe Pro Thr His Phe Lys Gly Pro Lys Leu Asn Pro Ile
      5              10              15
Lys Val Asn Pro Asn Phe Phe Glu Arg Asn Pro Lys Val Ala Arg Val
      20              25              30
Leu Gln Ile Thr Ala Val Val Leu Gly Ile Ile Ala Leu Leu Ser Gly
      35              40              45
Ile Val Leu Ile Ile Gly Thr Pro Leu Gly Ala Pro Ile Ser Met Ile
      50              55              60
Leu Gly Gly Cys Leu Leu Ala Ser Gly Gly Ala Leu Phe Val Gly Gly
      65              70              75              80
Thr Ile Ala Thr Ile Leu Gln Ala Arg Asn Ser Tyr Lys Lys Ala Val
      85              90              95
Asn Gln Lys Lys Leu Ser Glu Pro Leu Met Glu Arg Pro Glu Leu Lys
      100              105              110
Ala Leu Asp Tyr Ser Leu Asp Leu Lys Glu Val Trp Asp Leu His His
      115              120              125
Ser Val Val Lys His Leu Lys Lys Leu Asp Leu Asn Leu Ser Lys Thr
      130              135              140
Gln Arg Glu Val Leu Asn Gln Ile Lys Ile Asp Asp Glu Gly Pro Ser
      145              150              155              160
Leu Gly Glu Cys Ala Ala Met Ile Ser Glu Asn Tyr Asp Ala Cys Leu
      165              170              175
Lys Met Leu Ala Tyr Arg Glu Glu Leu Leu Lys Glu Gln Thr Gln Tyr
      180              185              190
Gln Glu Thr Arg Phe Asn Gln Asn Leu Thr His Arg Asn Lys Val Leu
      195              200              205
Leu Ser Ile Leu Ser Arg Ile Thr Asp Asn Ile Ser Lys Ala Gly Gly
      210              215              220
Val Phe Ser Leu Lys Phe Ser Thr Leu Ser Ser Arg Met Ser Arg Ile
      225              230              235              240
His Thr Thr Thr Thr Val Ile Leu Ala Leu Ser Ala Val Val Ser Val
      245              250              255

```

```

Met Val Val Ala Ala Leu Ile Pro Gly Gly Ile Leu Ala Leu Pro Ile
      260      265      270
Leu Leu Ala Val Ala Ile Ser Ala Gly Val Ile Val Thr Gly Leu Ser
      275      280      285
Tyr Leu Val Arg Gln Ile Leu Ser Asn Thr Lys Arg Asn Arg Gln Asp
      290      295      300
Phe Tyr Lys Asp Phe Val Lys Asn Val Asp Ile Glu Leu Leu Asn Gln
      305      310      315
Thr Val Thr Leu Gln Arg Phe Leu Phe Glu Met Leu Lys Gly Val Leu
      325      330      335
Lys Glu Glu Glu Glu Val Ser Leu Glu Gly Gln Asp Trp Tyr Thr Gln
      340      345      350
Tyr Ile Thr Asn Ala Pro Ile Glu Lys Arg Leu Ile Glu Glu Ile Arg
      355      360      365
Val Thr Tyr Lys Glu Ile Asp Ala Gln Thr Lys Lys Met Lys Thr Asp
      370      375      380
Leu Glu Phe Leu Glu Asn Glu Val Arg Ser Gly Arg Leu Ser Val Ala
      385      390      395
Ser Pro Ser Glu Asp Pro Ser Glu Thr Pro Ile Phe Thr Gln Gly Lys
      405      410      415
Glu Phe Ala Lys Leu Arg Arg Gln Thr Ser Gln Asn Ile Ser Thr Ile
      420      425      430
Tyr Gly Pro Asp Asn Glu Asn Ile Asp Pro Glu Phe Ser Leu Pro Trp
      435      440      445
Met Pro Lys Lys Glu Glu Glu Ile Asp His Ser Leu Glu Pro Val Thr
      450      455      460
Lys Leu Glu Pro Gly Ser Arg Glu Glu Leu Leu Val Glu Gly Val
      465      470      475
Asn Pro Thr Leu Arg Glu Leu Asn Met Arg Ile Ala Leu Leu Gln Gln
      485      490      495
Gln Leu Ser Ser Val Arg Lys Trp Arg His Pro Arg Gly Glu His Tyr
      500      505      510
Gly Asn Val Ile Tyr Ser Asp Thr Glu Leu Asp Arg Ile Gln Met Leu
      515      520      525
Glu Gly Ala Phe Tyr Asn His Leu Arg Glu Ala Gln Glu Glu Ile Thr
      530      535      540
Gln Ser Leu Gly Asp Leu Val Asp Ile Gln Asn Arg Ile Leu Gly Ile
      545      550      555
Ile Val Glu Gly Asp Ser Asp Ser Arg Thr Glu Glu Glu Pro Gln Glu
      565      570      575

```

<210> 74

<211> 361

<212> PRT

<213> Chlamydia pneumoniae

<400> 74

```

Met Gln Gln Thr Val Ile Val Ala Met Ser Gly Gly Val Asp Ser Ser
      5      10      15
Val Val Ala Tyr Leu Phe Lys Lys Phe Thr Asn Tyr Lys Val Ile Gly
      20      25      30
Leu Phe Met Lys Asn Trp Glu Glu Asp Ser Glu Gly Gly Leu Cys Ser
      35      40      45
Ser Thr Lys Asp Tyr Glu Asp Val Glu Arg Val Cys Leu Gln Leu Asp
      50      55      60
Ile Pro Tyr Tyr Thr Val Ser Phe Ala Lys Glu Tyr Arg Glu Arg Val
      65      70      75
Phe Ala Arg Phe Leu Lys Glu Tyr Ser Leu Gly Tyr Thr Pro Asn Pro
      80      85      90
Asp Ile Leu Cys Asn Arg Glu Ile Lys Phe Asp Leu Leu Gln Lys Lys

```

```

100      105      110
Val Gln Glu Leu Gly Gly Asp Tyr Leu Ala Thr Gly His Tyr Cys Arg
115
Leu Asn Thr Glu Leu Gln Glu Thr Gln Leu Leu Arg Gly Cys Asp Pro
130
Gln Lys Asp Gln Ser Tyr Phe Leu Ser Gly Thr Pro Lys Ser Ala Leu
145
His Asn Val Leu Phe Pro Leu Gly Glu Met Asn Lys Thr Glu Val Arg
165
Ala Ile Ala Ala Gln Ala Ala Leu Pro Thr Ala Glu Lys Lys Asp Ser
180
Thr Gly Ile Cys Phe Ile Gly Lys Arg Pro Phe Lys Glu Phe Leu Glu
195
Lys Phe Leu Pro Asn Lys Thr Gly Asn Val Ile Asp Trp Asp Thr Lys
210
Glu Ile Val Gly Gln His Gln Gly Ala His Tyr Tyr Thr Ile Gly Gln
225
Arg Arg Gly Leu Asp Leu Gly Gly Ser Glu Lys Pro Cys Tyr Val Val
245
Gly Lys Asn Ile Glu Glu Asn Ser Ile Tyr Ile Val Arg Gly Glu Asp
260
His Pro Gln Leu Tyr Leu Arg Glu Leu Thr Ala Arg Glu Leu Asn Trp
275
Phe Thr Pro Pro Lys Ser Gly Cys His Cys Ser Ala Lys Val Arg Tyr
290
Arg Ser Pro Asp Glu Ala Cys Thr Ile Asp Tyr Ser Ser Gly Asp Glu
305
Val Lys Val Arg Phe Ser Gln Pro Val Lys Ala Val Thr Pro Gly Gln
325
Thr Ile Ala Phe Tyr Gln Gly Asp Thr Cys Leu Gly Ser Gly Val Ile
340
Asp Val Pro Met Ile Pro Ser Glu Gly
355      360

```

<210> 75
 <211> 1609
 <212> PRT
 <213> Chlamydia pneumoniae

```

<400> 75
Met Val Ala Lys Lys Thr Val Arg Ser Tyr Arg Ser Ser Phe Ser His
5      10      15
Ser Val Ile Val Ala Ile Leu Ser Ala Gly Ile Ala Phe Glu Ala His
20      25      30
Ser Leu His Ser Ser Glu Leu Asp Leu Gly Val Phe Asn Lys Gln Phe
35      40      45
Glu Glu His Ser Ala His Val Glu Glu Ala Gln Thr Ser Val Leu Lys
50      55      60
Gly Ser Asp Pro Val Asn Pro Ser Gln Lys Glu Ser Glu Lys Val Leu
65      70      75      80
Tyr Thr Gln Val Pro Leu Thr Gln Gly Ser Ser Gly Glu Ser Leu Asp
85      90      95
Leu Ala Asp Ala Asn Phe Leu Glu His Phe Gln His Leu Phe Glu Glu
100      105      110
Thr Thr Val Phe Gly Ile Asp Gln Lys Leu Val Trp Ser Asp Leu Asp
115      120      125
Thr Arg Asn Phe Ser Gln Pro Thr Gln Glu Pro Asp Thr Ser Asn Ala
130      135      140
Val Ser Glu Lys Ile Ser Ser Asp Thr Lys Glu Asn Arg Lys Asp Leu
145      150      155      160

```

Glu Thr Glu Asp Pro Ser Lys Lys Ser Gly Leu Lys Glu Val Ser Ser
 165 170 175
 Asp Leu Pro Lys Ser Pro Glu Thr Ala Val Ala Ala Ile Ser Glu Asp
 180 185 190
 Leu Glu Ile Ser Glu Asn Ile Ser Ala Arg Asp Pro Leu Gln Gly Leu
 195 200 205
 Ala Phe Phe Tyr Lys Asn Thr Ser Ser Gln Ser Ile Ser Glu Lys Asp
 210 215 220
 Ser Ser Phe Gln Gly Ile Ile Phe Ser Gly Ser Gly Ala Asn Ser Gly
 225 230 235 240
 Leu Gly Phe Glu Asn Leu Lys Ala Pro Lys Ser Gly Ala Ala Val Tyr
 245 250 255
 Ser Asp Arg Asp Ile Val Phe Glu Asn Leu Val Lys Gly Leu Ser Phe
 260 265 270
 Ile Ser Cys Glu Ser Leu Glu Asp Gly Ser Ala Ala Gly Val Asn Ile
 275 280 285
 Val Val Thr His Cys Gly Asp Val Thr Leu Thr Asp Cys Ala Thr Gly
 290 295 300
 Leu Asp Leu Glu Ala Leu Arg Leu Val Lys Asp Phe Ser Arg Gly Gly
 305 310 315 320
 Ala Val Phe Thr Ala Arg Asn His Glu Val Gln Asn Asn Leu Ala Gly
 325 330 335
 Gly Ile Leu Ser Val Val Gly Asn Lys Gly Ala Ile Val Val Glu Lys
 340 345 350
 Asn Ser Ala Glu Lys Ser Asn Gly Gly Ala Phe Ala Cys Gly Ser Phe
 355 360 365
 Val Tyr Ser Asn Asn Glu Asn Thr Ala Leu Trp Lys Glu Asn Gln Ala
 370 375 380
 Leu Ser Gly Gly Ala Ile Ser Ser Ala Ser Asp Ile Asp Ile Gln Gly
 385 390 395 400
 Asn Cys Ser Ala Ile Glu Phe Ser Gly Asn Gln Ser Leu Ile Ala Leu
 405 410 415
 Gly Glu His Ile Gly Leu Thr Asp Phe Val Gly Gly Ala Leu Ala
 420 425 430
 Ala Gln Gly Thr Leu Thr Leu Arg Asn Asn Ala Val Val Gln Cys Val
 435 440 445
 Lys Asn Thr Ser Lys Thr His Gly Gly Ala Ile Leu Ala Gly Thr Val
 450 455 460
 Asp Leu Asn Glu Thr Ile Ser Glu Val Ala Phe Lys Gln Asn Thr Ala
 465 470 475 480
 Ala Leu Thr Gly Gly Ala Leu Ser Ala Asn Asp Lys Val Ile Ile Ala
 485 490 495
 Asn Asn Phe Gly Glu Ile Leu Phe Glu Gln Asn Glu Val Arg Asn His
 500 505 510
 Gly Gly Ala Ile Tyr Cys Gly Cys Arg Ser Asn Pro Lys Leu Glu Gln
 515 520 525
 Lys Asp Ser Gly Glu Asn Ile Asn Ile Ile Gly Asn Ser Gly Ala Ile
 530 535 540
 Thr Phe Leu Lys Asn Lys Ala Ser Val Leu Glu Val Met Thr Gln Ala
 545 550 555 560
 Glu Asp Tyr Ala Gly Gly Gly Ala Leu Trp Gly His Asn Val Leu Leu
 565 570 575
 Asp Ser Asn Ser Gly Asn Ile Gln Phe Ile Gly Asn Ile Gly Gly Ser
 580 585 590
 Thr Phe Trp Ile Gly Glu Tyr Val Gly Gly Gly Ala Ile Leu Ser Thr
 595 600 605
 Asp Arg Val Thr Ile Ser Asn Asn Ser Gly Asp Val Val Phe Lys Gly
 610 615 620
 Asn Lys Gly Gln Cys Leu Ala Gln Lys Tyr Val Ala Pro Gln Glu Thr
 625 630 635 640
 Ala Pro Val Glu Ser Asp Ala Ser Ser Thr Asn Lys Asp Glu Lys Ser

Leu	Asn	Ala	Cys	445	Ser	His	Gly	Asp	His	650	Tyr	Pro	Pro	Lys	Thr	Val	Glu
Glu	Glu	Val	Pro	660	Pro	Ser	Leu	Leu	665	Glu	His	Pro	Pro	Val	Val	Ser	Ser
Thr	Asp	675	Ile	Arg	Gly	Gly	Gly	Ala	680	Ile	Leu	Ala	Gln	His	Ile	Phe	Ile
Thr	Asp	690	Asn	Thr	Gly	Asn	Leu	Arg	695	Phe	Ser	Gly	700	Asn	Leu	Gly	Gly
705	Glu	Glu	Ser	Ser	Thr	Val	Gly	Asp	710	Leu	Ala	Ile	Val	Gly	Gly	Gly	720
Leu	Leu	Ser	Thr	725	Asn	Glu	Val	Asn	730	Val	Cys	Ser	Asn	Gln	Asn	Val	Val
Phe	Ser	Asp	740	Asn	Val	Thr	Ser	Asn	745	Cys	Asp	Ser	Gly	Gly	Ala	Ile	
Leu	Ala	Lys	755	Lys	Val	Asp	Ile	Ser	760	Ala	Asn	His	Ser	Val	Glu	Phe	Val
770	Ser	Asn	Gly	Ser	Gly	Lys	Phe	Gly	775	Gly	Ala	Val	Cys	Ala	Leu	Asn	Glu
785	Ser	Val	Asn	Ile	Thr	Asp	Asn	Gly	790	Ser	Ala	Val	Ser	Phe	Ser	Lys	Asn
Arg	Thr	Arg	805	Gly	Gly	Ala	Gly	Val	810	Ala	Ala	Pro	Gln	Gly	Ser	Val	
Thr	Ile	Cys	820	Gly	Asn	Gln	Gly	Asn	825	Ile	Ala	Phe	Lys	Glu	Asn	Phe	Val
Phe	Gly	Ser	835	Glu	Asn	Gln	Arg	Ser	840	Gly	Gly	Ala	Ile	Ile	Ala	Asn	
Ser	850	Ser	Val	Asn	Ile	Gln	Asp	Ser	855	Asn	Ala	Gly	Asp	Ile	Leu	Phe	Val
865	Asn	Ser	Thr	Gly	Ser	Tyr	Gly	Gly	870	Ala	Ile	Phe	Val	Gly	Ser	Leu	Val
Ala	Ser	Glu	885	Gly	Ser	Asn	Pro	Arg	890	Thr	Leu	Thr	Ile	Thr	Gly	Asn	Ser
Gly	Asp	Ile	900	Leu	Phe	Ala	Lys	Asn	905	Ser	Thr	Gln	Thr	Ala	Ala	Ser	Leu
Ser	Glu	Lys	915	Asp	Ser	Phe	Gly	Gly	920	Gly	Ala	Ile	Tyr	Thr	Gln	Asn	Leu
Lys	930	Ile	Val	Lys	Asn	Ala	Gly	Asn	935	Val	Ser	Phe	Tyr	Gly	Asn	Arg	Ala
945	Pro	Ser	Gly	Ala	Gly	Val	Gln	Ile	950	Ala	Asp	Gly	Gly	Thr	Val	Cys	Leu
Glu	Ala	Phe	955	Gly	Gly	Asp	Ile	Leu	960	Phe	Gly	Gly	Asn	Ile	Asn	Phe	Asp
Gly	Ser	Phe	965	Asn	Ala	Ile	His	Leu	970	Cys	Gly	Asn	Asp	Ser	Lys	Ile	Val
Glu	1000	Ser	Ala	Val	Gln	Asp	Lys	Asn	1005	Ile	Ile	Phe	Gln	Asp	Ala	Ile	
1010	Thr	Tyr	Glu	Glu	Asn	Thr	Ile	Arg	1015	Gly	Leu	Pro	Asp	Lys	Asp	Val	Ser
1025	Pro	Leu	Ser	Ala	Pro	Ser	Leu	Ile	1030	Phe	Asn	Ser	Lys	Pro	Gln	Asp	Asp
Ser	Ala	Gln	1045	His	His	Glu	Gly	Thr	1050	Ile	Arg	Phe	Ser	Arg	Gly	Val	Ser
Lys	1060	Ile	Pro	Gln	Ile	Ala	Ala	Ile	1065	Gln	Glu	Gly	Thr	Leu	Ala	Leu	Ser
Gln	1075	Asn	Ala	Glu	Leu	Trp	Leu	Ala	1080	Gly	Leu	Lys	Gln	Glu	Thr	Gly	Ser
Ser	1090	Ser	Val	Leu	Ser	Ala	Gly	Ser	1095	Ile	Leu	Arg	Ile	Phe	Asp	Ser	Gln
1105	Val	Asp	Ser	Ser	Ala	Pro	Leu	Pro	1110	Glu	Asn	Lys	Glu	Glu	Thr	Leu	
					1125				1130								

Val Ser Ala Gly Val Gln Ile Asn Met Ser Ser Pro Thr Pro Asn Lys
 1140 1145 1150
 Asp Lys Ala Val Asp Thr Pro Val Leu Ala Asp Ile Ile Ser Ile Thr
 1155 1160 1165
 Val Asp Leu Ser Ser Phe Val Pro Glu Gln Asp Gly Thr Leu Pro Leu
 1170 1175 1180
 Pro Pro Glu Ile Ile Ile Pro Lys Gly Thr Lys Leu His Ser Asn Ala
 1185 1190 1195 1200
 Ile Asp Leu Lys Ile Ile Asp Pro Thr Asn Val Gly Tyr Glu Asn His
 1205 1210 1215
 Ala Leu Leu Ser Ser His Lys Asp Ile Pro Leu Ile Ser Leu Lys Thr
 1220 1225 1230
 Ala Glu Gly Met Thr Gly Thr Pro Thr Ala Asp Ala Ser Leu Ser Asn
 1235 1240 1245
 Ile Lys Ile Asp Val Ser Leu Pro Ser Ile Thr Pro Ala Thr Tyr Gly
 1250 1255 1260
 His Thr Gly Val Trp Ser Glu Ser Lys Met Glu Asp Gly Arg Leu Val
 1265 1270 1275 1280
 Val Gly Trp Gln Pro Thr Gly Tyr Lys Leu Asn Pro Glu Lys Gln Gly
 1285 1290 1295
 Ala Leu Val Leu Asn Asn Leu Trp Ser His Tyr Thr Asp Leu Arg Ala
 1300 1305 1310
 Leu Lys Gln Glu Ile Phe Ala His His Thr Ile Ala Gln Arg Met Glu
 1315 1320 1325
 Leu Asp Phe Ser Thr Asn Val Trp Gly Ser Gly Leu Gly Val Val Glu
 1330 1335 1340
 Asp Cys Gln Asn Ile Gly Glu Phe Asp Gly Phe Lys His His Leu Thr
 1345 1350 1355 1360
 Gly Tyr Ala Leu Gly Leu Asp Thr Gln Leu Val Glu Asp Phe Leu Ile
 1365 1370 1375
 Gly Gly Cys Phe Ser Gln Phe Phe Gly Lys Thr Glu Ser Gln Ser Tyr
 1380 1385 1390
 Lys Ala Lys Asn Asp Val Lys Ser Tyr Met Gly Ala Ala Tyr Ala Gly
 1395 1400 1405
 Ile Leu Ala Gly Pro Trp Leu Ile Lys Gly Ala Phe Val Tyr Gly Asn
 1410 1415 1420
 Ile Asn Asn Asp Leu Thr Thr Asp Tyr Gly Thr Leu Gly Ile Ser Thr
 1425 1430 1435 1440
 Gly Ser Trp Ile Gly Lys Gly Phe Ile Ala Gly Thr Ser Ile Asp Tyr
 1445 1450 1455
 Arg Tyr Ile Val Asn Pro Arg Arg Phe Ile Ser Ala Ile Val Ser Thr
 1460 1465 1470
 Val Val Pro Phe Val Glu Ala Glu Tyr Val Arg Ile Asp Leu Pro Glu
 1475 1480 1485
 Ile Ser Glu Gln Gly Lys Glu Val Arg Thr Phe Gln Lys Thr Arg Phe
 1490 1495 1500
 Glu Asn Val Ala Ile Pro Phe Gly Phe Ala Leu Glu His Ala Tyr Ser
 1505 1510 1515 1520
 Arg Gly Ser Arg Ala Glu Val Asn Ser Val Gln Leu Ala Tyr Val Phe
 1525 1530 1535
 Asp Val Tyr Arg Lys Gly Pro Val Ser Leu Ile Thr Leu Lys Asp Ala
 1540 1545 1550
 Ala Tyr Ser Trp Lys Ser Tyr Gly Val Asp Ile Pro Cys Lys Ala Trp
 1555 1560 1565
 Lys Ala Arg Leu Ser Asn Asn Thr Glu Trp Asn Ser Tyr Leu Ser Thr
 1570 1575 1580
 Tyr Leu Ala Phe Asn Tyr Glu Trp Arg Glu Asp Leu Ile Ala Tyr Asp
 1585 1590 1595 1600
 Phe Asn Gly Gly Ile Arg Ile Ile Phe
 1605

<210> 76
 <211> 196
 <212> PRT
 <213> Chlamydia pneumoniae

<400> 76
 Met Thr Leu Ser Leu Val Gly Lys Glu Ala Pro Asp Phe Val Ala Gln
 5 10 15
 Ala Val Val Asn Gly Glu Thr Cys Thr Val Ser Leu Lys Asp Tyr Leu
 20 25 30
 Gly Lys Tyr Val Val Leu Phe Phe Tyr Pro Lys Asp Phe Thr Tyr Val
 35 40 45
 Cys Pro Thr Glu Leu His Ala Phe Gln Asp Ala Leu Gly Glu Phe His
 50 55 60
 Thr Arg Gly Ala Glu Val Ile Gly Cys Ser Val Asp Asp Ile Ala Thr
 65 70 75 80
 His Gln Gln Trp Leu Ala Thr Lys Lys Gln Gly Gly Ile Glu Gly
 85 90 95
 Ile Thr Tyr Pro Leu Leu Ser Asp Glu Asp Lys Val Ile Ser Arg Ser
 100 105 110
 Tyr His Val Leu Lys Pro Glu Glu Glu Leu Ser Phe Arg Gly Val Phe
 115 120 125
 Leu Ile Asp Lys Gly Gly Ile Ile Arg His Leu Val Val Asn Asp Leu
 130 135 140
 Pro Leu Gly Arg Ser Ile Glu Glu Glu Leu Arg Thr Leu Asp Ala Leu
 145 150 155
 Ile Phe Phe Glu Thr Asn Gly Leu Val Cys Pro Ala Asn Trp His Glu
 165 170 175
 Gly Glu Arg Ala Met Ala Pro Asn Glu Glu Gly Leu Gln Asn Tyr Phe
 180 185 190
 Gly Thr Ile Asp
 195

<210> 77
 <211> 619
 <212> PRT
 <213> Chlamydia pneumoniae

<400> 77
 Met Lys Lys Gly Lys Leu Gly Ala Ile Val Phe Gly Leu Leu Phe Thr
 5 10 15
 Ser Ser Val Ala Gly Phe Ser Lys Asp Leu Thr Lys Asp Asn Ala Tyr
 20 25 30
 Gln Asp Leu Asn Val Ile Glu His Leu Ile Ser Leu Lys Tyr Ala Pro
 35 40 45
 Leu Pro Trp Lys Glu Leu Leu Phe Gly Trp Asp Leu Ser Gln Gln Thr
 50 55 60
 Gln Gln Ala Arg Leu Gln Leu Val Leu Glu Glu Lys Pro Thr Thr Asn
 65 70 75 80
 Tyr Cys Gln Lys Val Leu Ser Asn Tyr Val Arg Ser Leu Asn Asp Tyr
 85 90 95
 His Ala Gly Ile Thr Phe Tyr Arg Thr Glu Ser Ala Tyr Ile Pro Tyr
 100 105 110
 Val Leu Lys Leu Ser Glu Asp Gly His Val Phe Val Val Asp Val Gln
 115 120 125
 Thr Ser Gln Gly Asp Ile Tyr Leu Gly Asp Glu Ile Leu Glu Val Asp
 130 135 140
 Gly Met Gly Ile Arg Glu Ala Ile Glu Ser Leu Arg Phe Gly Arg Gly
 145 150 155 160

Ser Ala Thr Asp Tyr Ser Ala Ala Val Arg Ser Leu Thr Ser Arg Ser
 165 170 175
 Ala Ala Phe Gly Asp Ala Val Pro Ser Gly Ile Ala Met Leu Lys Leu
 180 185 190
 Arg Arg Pro Ser Gly Leu Ile Arg Ser Thr Pro Val Arg Trp Arg Tyr
 195 200 205
 Thr Pro Glu His Ile Gly Asp Phe Ser Leu Val Ala Pro Leu Ile Pro
 210 215 220
 Glu His Lys Pro Gln Leu Pro Thr Gln Ser Cys Val Leu Phe Arg Ser
 225 230 235 240
 Gly Val Asn Ser Gln Ser Ser Ser Ser Ser Leu Phe Ser Ser Tyr Met
 245 250 255
 Val Pro Tyr Phe Trp Glu Glu Leu Arg Val Gln Asn Lys Gln Arg Phe
 260 265 270
 Asp Ser Asn His His Ile Gly Ser Arg Asn Gly Phe Leu Pro Thr Phe
 275 280 285
 Gly Pro Ile Leu Trp Glu Gln Asp Lys Gly Pro Tyr Arg Ser Tyr Ile
 290 295 300
 Phe Lys Ala Lys Asp Ser Gln Gly Asn Pro His Arg Ile Gly Phe Leu
 305 310 315 320
 Arg Ile Ser Ser Tyr Val Trp Thr Asp Leu Glu Gly Leu Glu Glu Asp
 325 330 335
 His Lys Asp Ser Pro Trp Glu Leu Phe Gly Glu Ile Ile Asp His Leu
 340 345 350
 Glu Lys Glu Thr Asp Ala Leu Ile Ile Asp Gln Thr His Asn Pro Gly
 355 360 365
 Gly Ser Val Phe Tyr Leu Tyr Ser Leu Leu Ser Met Leu Thr Asp His
 370 375 380
 Pro Leu Asp Thr Pro Lys His Arg Met Ile Phe Thr Gln Asp Glu Val
 385 390 395 400
 Ser Ser Ala Leu His Trp Gln Asp Leu Leu Glu Asp Val Phe Thr Asp
 405 410 415
 Glu Gln Ala Val Ala Val Leu Gly Glu Thr Met Glu Gly Tyr Cys Met
 420 425 430
 Asp Met His Ala Val Ala Ser Leu Gln Asn Phe Ser Gln Ser Val Leu
 435 440 445
 Ser Ser Trp Val Ser Gly Asp Ile Asn Leu Ser Lys Pro Met Pro Leu
 450 455 460
 Leu Gly Phe Ala Gln Val Arg Pro His Pro Lys His Gln Tyr Thr Lys
 465 470 475 480
 Pro Leu Phe Met Leu Ile Asp Glu Asp Asp Phe Ser Cys Gly Asp Leu
 485 490 495
 Ala Pro Ala Ile Leu Lys Asp Asn Gly Arg Ala Thr Leu Ile Gly Lys
 500 505 510
 Pro Thr Ala Gly Ala Gly Gly Phe Val Phe Gln Val Thr Phe Pro Asn
 515 520 525
 Arg Ser Gly Ile Lys Gly Leu Ser Leu Thr Gly Ser Leu Ala Val Arg
 530 535 540
 Lys Asp Gly Glu Phe Ile Glu Asn Leu Gly Val Ala Pro His Ile Asp
 545 550 555 560
 Leu Gly Phe Thr Ser Arg Asp Leu Gln Thr Ser Arg Phe Thr Asp Tyr
 565 570 575
 Val Glu Ala Val Lys Thr Ile Val Leu Thr Ser Leu Ser Glu Asn Ala
 580 585 590
 Lys Lys Ser Glu Glu Gln Thr Ser Pro Gln Glu Thr Pro Glu Val Ile
 595 600 605
 Arg Val Ser Tyr Pro Thr Thr Ser Ala Ser
 610 615

<211> 651

<212> PRT

<213> Chlamydia pneumoniae

<400> 78

```

Met Val Asn Pro Ile Gly Pro Gly Pro Ile Asp Glu Thr Glu Arg Thr
      5      10      15
Pro Pro Ala Asp Leu Ser Ala Gln Gly Leu Glu Ala Ser Ala Ala Asn
      20      25      30
Lys Ser Ala Glu Ala Gln Arg Ile Ala Gly Ala Glu Ala Lys Pro Lys
      35      40      45
Glu Ser Lys Thr Asp Ser Val Glu Arg Trp Ser Ile Leu Arg Ser Ala
      50      55      60
Val Asn Ala Leu Met Ser Leu Ala Asp Lys Leu Gly Ile Ala Ser Ser
      65      70      75      80
Asn Ser Ser Ser Ser Thr Ser Arg Ser Ala Asp Val Asp Ser Thr Thr
      85      90      95
Ala Thr Ala Pro Thr Pro Pro Pro Thr Phe Asp Asp Tyr Lys Thr
      100      105      110
Gln Ala Gln Thr Ala Tyr Asp Thr Ile Phe Thr Ser Thr Ser Leu Ala
      115      120      125
Asp Ile Gln Ala Ala Leu Val Ser Leu Gln Asp Ala Val Thr Asn Ile
      130      135      140
Lys Asp Thr Ala Ala Thr Asp Glu Glu Thr Ala Ile Ala Glu Trp
      145      150      155      160
Glu Thr Lys Asn Ala Asp Ala Val Lys Val Gly Ala Gln Ile Thr Glu
      165      170      175
Leu Ala Lys Tyr Ala Ser Asp Asn Gln Ala Ile Leu Asp Ser Leu Gly
      180      185      190
Lys Leu Thr Ser Phe Asp Leu Leu Gln Ala Ala Leu Leu Gln Ser Val
      195      200      205
Ala Asn Asn Asn Lys Ala Ala Glu Leu Leu Lys Glu Met Gln Asp Asn
      210      215      220
Pro Val Val Pro Gly Lys Thr Pro Ala Ile Ala Gln Ser Leu Val Asp
      225      230      235      240
Gln Thr Asp Ala Thr Ala Thr Gln Ile Glu Lys Asp Gly Asn Ala Ile
      245      250      255
Arg Asp Ala Tyr Phe Ala Gly Gln Asn Ala Ser Gly Ala Val Glu Asn
      260      265      270
Ala Lys Ser Asn Asn Ser Ile Ser Asn Ile Asp Ser Ala Lys Ala Ala
      275      280      285
Ile Ala Thr Ala Lys Thr Gln Ile Ala Glu Ala Gln Lys Lys Phe Pro
      290      295      300
Asp Ser Ser Ile Leu Gln Glu Ala Glu Gln Met Val Ile Gln Ala Glu
      305      310      315      320
Lys Asp Leu Lys Asn Ile Lys Pro Ala Asp Gly Ser Asp Val Pro Asn
      325      330      335
Pro Gly Thr Thr Val Gly Gly Ser Lys Gln Gln Gly Ser Ser Ile Gly
      340      345      350
Ser Ile Arg Val Ser Met Leu Leu Asp Asp Ala Glu Asn Glu Thr Ala
      355      360      365
Ser Ile Leu Met Ser Gly Phe Arg Gln Met Ile His Met Phe Asn Thr
      370      375      380
Glu Asn Pro Asp Ser Gln Ala Ala Gln Gln Glu Leu Ala Ala Gln Ala
      385      390      395      400
Arg Ala Ala Lys Ala Ala Gly Asp Asp Ser Ala Ala Ala Leu Ala
      405      410      415
Asp Ala Gln Lys Ala Leu Glu Ala Ala Leu Gly Lys Ala Gly Gln Gln
      420      425      430
Gln Gly Ile Leu Asn Ala Leu Gly Gln Ile Ala Ser Ala Val Val
      435      440      445

```

```

Ser Ala Gly Val Pro Pro Ala Ala Ala Ser Ser Ile Gly Ser Ser Val
450 455 460
Lys Gln Leu Tyr Lys Thr Ser Lys Ser Thr Gly Ser Asp Tyr Lys Thr
465 470 475 480
Gln Ile Ser Ala Gly Tyr Asp Ala Tyr Lys Ser Ile Asn Asp Ala Tyr
485 490 495
Gly Arg Ala Arg Asn Asp Ala Thr Arg Asp Val Ile Asn Asn Val Ser
500 505 510
Thr Pro Ala Leu Thr Arg Ser Val Pro Arg Ala Arg Thr Glu Ala Arg
515 520 525
Gly Pro Glu Lys Thr Asp Gln Ala Leu Ala Arg Val Ile Ser Gly Asn
530 535 540
Ser Arg Thr Leu Gly Asp Val Tyr Ser Gln Val Ser Ala Leu Gln Ser
545 550 555 560
Val Met Gln Ile Ile Gln Ser Asn Pro Gln Ala Asn Asn Glu Glu Ile
565 570 575
Arg Gln Lys Leu Thr Ser Ala Val Thr Lys Pro Pro Gln Phe Gly Tyr
580 585 590
Pro Tyr Val Gln Leu Ser Asn Asp Ser Thr Gln Lys Phe Ile Ala Lys
595 600 605
Leu Glu Ser Leu Phe Ala Glu Gly Ser Arg Thr Ala Ala Glu Ile Lys
610 615 620
Ala Leu Ser Phe Glu Thr Asn Ser Leu Phe Ile Gln Val Leu Val
625 630 635 640
Asn Ile Gly Ser Leu Tyr Ser Gly Tyr Leu Gln
645 650

```

<210> 79

<211> 87

<212> PRT

<213> *Chlamydia pneumoniae*

<400> 79

```

Met Ser Gln Lys Asn Lys Asn Ser Ala Phe Met His Pro Val Asn Ile
5 10 15
Ser Thr Asp Leu Ala Val Ile Val Gly Lys Gly Pro Met Pro Arg Thr
20 25 30
Glu Ile Val Lys Lys Val Trp Glu Tyr Ile Lys Lys His Asn Cys Gln
35 40 45
Asp Gln Lys Asn Lys Arg Asn Ile Leu Pro Asp Ala Asn Leu Ala Lys
50 55 60
Val Phe Gly Ser Ser Asp Pro Ile Asp Met Phe Gln Met Thr Lys Ala
65 70 75 80
Leu Ser Lys His Ile Val Lys
85

```

<210> 80

<211> 3048

<212> DNA

<213> *Chlamydia trachomatis* serovar D

<400> 80

```

atgccttttt ctttgagatc tacatcattt tgttttttag cttgtttgtg ttctatttcg 60
tatggattcg cgagctctcc tcaagtgtta acacctaatg taaccactcc ttttaagggg 120
gacgatgttt actigaatgg agaactgcgt tttgtcaatg tctatgcagg ggcagagaaac 180
ggctcaatta tctcagctaa tggcgacaa ttaacyatta ccggacaaaa ccatacatta 240
tcatttacag attctcaagg gccagttcct caaaattatg cttctatttc agcgcagagag 300
acacttactc tgaagatttt ttogagtttg atgttctcga aaaatgttcc ttgcggagaa 360
aagggaatga tctcagggaa aaccgtgagt atttcggag caggcgaaat gattttttgg 420
gataactctg tggggtattc tccctttgtc attgtgccag catgactacc aactctccca 480

```

gcaccagcagc cagctcctgc tgcctcaagc tctttatctc caacagtgtag tgatgcctgg 540
 aaagggctcta tttttctgtg agagactagt ttggagatct caggcgctcaa aaaaggggtc 600
 atgtctcgata ataactgcgg gaatttttga acagtttttc gaggtaatat taataaataat 660
 gctgtgtagt ggggtagtgg gtctgtctaca acaccaagtt ttacagcttaa aaactgttaa 720
 gggaagtttt otttccacaga taacgtagcc tctgtgtggag goggagtgtg ctacaagaaga 780
 actgtctctt tcaaaacaaa taaggaaggg atattctctcc gagggaacac agcatatogat 840
 gattgagctg tcttctgtcg tactagtogg gatcagaata cggagacagg agcggctgtga 900
 ggagtttatt gotctcoga tgattctgtg aagtttgaag gcaataaagg ttctatttgt 960
 ttgtattaca actttgcaaa aggcagaggg ggaagcatcc taacgaaaga attctctctt 1020
 gtacgagatg attcggttgt otttagtaac aatacagcag aaaagggcgg tggagctatt 1080
 tatgtctcta ctatcgatat aagcagcaat ggaggatcga ttctatttga aagaaacga 1140
 gctgcagaa gaggcgcaat ctgcgtgagt gaagcaagct ctggttcaac tggaaatctt 1200
 actttaaggg cttctgatgg ggatatttgt tttctggga atagaogag tgatogtctt 1260
 ggagagcgca ggcagcaag aatcttaagt gatggaagca ctgttttctt aaatgctccc 1320
 ggactatcga agctgatctt ttatgatctt gtagtacaaa ataattcagc agcgggttga 1380
 tcgacacact cacactcttc ttctctatg cctggtgctg tcaagattaa tcagtcoggt 1440
 aatggatctg tgattttttac cgcgagtgca ttgactctt cagaaaaact tcaagttctt 1500
 aactctactt ctactctccc agggagctctg actgtgtcag gaggggagt ttgtgtgag 1560
 gaaggagcta ccttaactac tgggaacatt acagccactc acagacttga gactttaga 1620
 tcoggagctt cgttgtctgc cgttgcaggt gctgcaata ataattatac ttgtacagta 1680
 totaagttgg ggattgtatt agaactcttt ttaactctca actataagac ggcactactg 1740
 ggtgoggatg gaacagttac ttttaacagc ggcctacttt tagacactgt gatggagat 1800
 gaggcagagg tatatgata tcocgttttt gtgggagctg tgacaactcc ttgttgactt 1860
 ctactctcta gtatgtctag taacggagtt acaaaaaatt ctgtoactat taatgatgca 1920
 gaogctgcgc actatgggta toaaggctct ggtactctat attggacgaa acgcctctgt 1980
 gctctctgat ctaaggggat ggaactctat aataccaata acactctgta tctgacatgg 2040
 agaactgctt ogaaattacg tgaatatoga cctggactct cctggactct agagactagta 2100
 cccaactctc tttgggtagc gggatctgca ttaagaactc ttactaatgg ttgaaagaa 2160
 caacttgttt ctgagatagt tggatttgta goactctctg atgtctctgg ttgtattatt 2220
 ttgaattata cgaagatga tcgggatggc tttttageta gatatggggg attccagggc 2280
 accgagcctt ccaattatga aatgggtca atattttgg tggcttttgg acaactctat 2340
 ggtagcagaa agagcagaat gtattactct aaagatgctg ggaactatgac gatgtgtctc 2400
 tgtttoggaa gaagtacgt agatattaaa ggaacagaaa cgtttatgta ttgggagaog 2460
 gcttatggct attctgtgca cagaatgcal acogagatatt ttaatgacaa aacgcagaag 2520
 ttogactatt ogaaatgca ttggcacaac aataactatt atggttttgg gggtgcogag 2580
 cataattctt tagagtactg caatctact cgtcagttcg ctgagatta tgagcttaca 2640
 gggttttatg gttttgaaat ggcgggagga tgtccagtt ctacaogaga aactggctcc 2700
 ctactagat attctgcctg oggtcaggg ctaaatatgt cgcctccaat aggaattgta 2760
 gctcatgag ttctcatgt gcgaagatct octctctatg aactgcaact agaatgtgga 2820
 tatagacaa acattttggt tgcactatca atttgcataa tggaaattat tgcataagca 2880
 gtaagacac ctatacaagg atctccogct gcacggcatg octctctctt ggaactgtcat 2940
 gatgaattgt atattctca ttttggaag ccatatgga actattctgt ggaatgctgt 3000
 cgtgcacaaa oggcacatt ttgatccat ggcttgaata gaactctt 3048

<210> 81

<211> 1038

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 81

atgcaagcagc atatttttga tggaaaacag aaacgcgtta atctaaatag caagcgtota 60
 gtgaactgca acoaggtcga tgtcaaccaa cttgttctta ttaagtacaa attggctgtg 120
 gaacattatt tgaatggctg cgaacaatac tggctcccta cagagatccc catggggaaa 180
 gacatcgaa ttggaagtcg ggtatgtctt totgaagat agcggogagat actctttgt 240
 aattttagtt ttttgcagac cgcagagagc ttgtttggga ataataattg tctagcaatt 300
 tttaaacatg taactaatcc ggaagcaga caatatcttt taagacaagc ttttgaaga 360
 goggttcaac cgcacacatt ttgttatatt tgtgagtcac tctggattga cggaaagaa 420
 attttcaatg cctataacga cgtgtgctg cgttaaggcca aagatgatt ccaagattgaa 480
 atcaactggca aggtatttga tctaataatt cgcacggact cctgttgagg tctacaggag 540
 ttgtttaaaa acttagtagg atactacatc attatggag ggattttctt ctatagtgtg 600
 tttgtgatga toctttctct ccaacagaaa aataagatga ttggtattgg agaacaatat 660

caatacatct	taagagatga	gacaatocac	ttgaacttgg	gtattgattt	gatcaacggg	720
ataaaagaag	agaaccocga	gatttggact	ccagagttac	agcaagaaat	tgtcgaaatta	780
attaaagcag	ctgtocgatt	agaaattgag	tatgcgcgaac	actgtctccc	tagaggggatt	840
ttgggattga	gagcttgat	gttccatgat	tatgtgcagc	atattgcoga	cogtoggttt	900
gaagaaatcg	gtataaaacc	tatttatcat	acgaaaaaac	cattcccttg	galgagcgaa	960
acaatagacc	ttaataaaga	gaaaaacttc	tttgaaacaa	gggttataga	atacacaact	1020
gcagcaagct	taacttgg					1038

<210> 82
 <211> 3159
 <212> DNA
 <213> *Chlamydia trachomatis* serovar D

<400> 82

atgtttacaa	ggatagttat	ggtogactta	caagaaaagc	aatgcacaa	tgttaagcgc	60
aattggaagt	ttgttccatt	cgatcggaac	cgtatttttc	aggtctttag	agcagotttt	120
cgagcaactc	gcagaattga	tgatccatag	cotttgcotg	aagatctgga	aagtccatata	180
cgctcgatga	agcgtatgat	agtttaagaa	gtttgcaaac	agattacaga	tggacaagtg	240
gttaactgtga	cgatatacga	agatatgtgt	gaagcccaac	tattgtgaa	tggtttgcga	300
gatgttgctc	gcgattatata	ttgttatcgg	galgacgcga	aagcgcatcg	gaaaaaatct	360
tggcaaaagc	tacatcgtgt	togtctgttg	gggactgttg	tacactttaa	tcctatgaaa	420
atttcogcgc	otttgaaaaa	agcttccoga	gtcaccgata	agactgaggg	gatgaatcca	480
agttctgtgc	gagaggaagt	caatgtcttg	acgcataaac	ttgtcgcgga	aataagaaag	540
ttgttgcctc	aacaggatag	acgcattgat	atcgagaaga	ttaagatat	tgttgaacag	600
caactaatgg	ttgttggcca	ttatgtcgtt	gcataaagac	atatctotta	tccgaaagct	660
cgctcgtcgt	ttcgtgataa	caagagaagc	gaogggagta	cgaaaaagac	tatagcaaga	720
gaagctgttg	aggtgcctcg	taaaagcgtg	tctactata	caatgacgca	ttccgagttg	780
ttggtcatt	ttagcgcgcg	ttgtagtctg	tttccagaaa	cgacagatgc	ggcgtctgct	840
acattatagg	otttcgcaaa	tttctattcc	ggatccaaag	agctggaagt	agtaactggc	900
tgtattatgg	cgctcgtcgc	caatattgaa	aaggagcctg	attatgcott	tgttgcgtgc	960
gagctottac	ttgacgttgt	atataaggaa	cgtttaggga	aatcgaaata	tcctgagagt	1020
ttagaacaag	caatcgcgca	tcatttccaa	cgctacatcg	cgagaaggga	tacctatcgt	1080
ctgaatgcctg	aactgaacaa	tctttttgat	ttgacgcgtg	tagccgatgc	tatggatata	1140
ttcogagatc	tacagtttlt	ttacatgggt	attccaaatc	tgtatgatcg	ttatttttaa	1200
ccacccagaag	gttgccgttt	agaaactccc	caaatttttt	ggatgcgcgt	tgtattgggg	1260
ttggcattga	atgagcaaga	caagaactct	tgggtattta	ctttttataa	tttgccttgc	1320
acattccgat	ataaccacgc	taacccaacc	ttgttcaatt	caggtatgcg	gaattctcag	1380
ttaagctott	gctatotttc	caactacaaa	gataatttgg	tcattatcta	taaggtoatt	1440
gctgataaag	ctatgctatc	taagtgggca	ggaggggatg	gtaatgattg	gacggcgagt	1500
cgctgcaacg	gggttttaat	taaaaggaaac	aattggaagaa	gtcaggggat	aattcttttt	1560
attaaggctga	caaatgatac	agcagctgca	gtgaatcaag	gtggtaaacg	caagggaagct	1620
gtatgctgtc	atttagaagt	ttggcacctc	gaactacgaag	atttcttgta	ttatgagaag	1680
aatacagagg	atgagcgtcg	acgggctcat	gatgtcaata	tagctagctg	gattccagat	1740
cttttcttca	aacggtttaca	gcataaaagg	acatggactc	tattcagccc	tgatgtgatt	1800
cogggattac	acgatgotta	tggggagaag	tttgagcgtt	tgtacgaaga	atagagcgtg	1860
aaggttgata	coggagagat	tcoggttatt	aagaaggtag	aagctgaaga	tcgtggagga	1920
aaaatgctca	goatgctttt	tgaacccgga	caccatgga	tgaactttaa	agatccatcc	1980
aacatcogtt	cgctcacaaga	tcataaaagg	gtgggtgcgt	gttccaatct	gtgtacaggag	2040
attttgttaa	actgctcggga	gacagaaact	gctgtttgta	atttaggatc	gatctactta	2100
gttcaacata	tttaggggga	tgggttagat	gaggaataaac	ttctctgagac	gatctctata	2160
gcagtcgcta	tgttggtata	cgtgattgat	attaaacttt	atccacaata	ggagagotaa	2220
gaggcgaaat	ttgctcacgc	cgctatttga	ttaggggtga	tgggtatcca	agatgccttg	2280
tataagctag	ataaagcta	tgcttcgcaa	gaagctgtag	aatttgtgta	ctacagttcca	2340
gagttgattt	cttaactatgc	gattcaagct	ttctgtctgc	tcgtcaaga	acagagcgact	2400
ttcagctctt	ataaaggatc	gaatgggatc	agaggtttgc	tcctatttga	tacagttccg	2460
tgttagcgca	actatcgagg	agaagcaaat	ctccagatgg	atacgtcaat	aagaaaagat	2520
tgggaacata	tcogtagttt	ggttaaaagc	catgggtatgc	gacattgtca	gotttatgct	2580
atagctcoga	cagcagacat	ctccaaacatt	ataaggagta	ctcaacttat	tagagccaaag	2640
tacaacacatt	tgtttgtgaa	gtctaatattg	tcoggagaagt	tcacgattcc	aaatgctgat	2700
ttaattgaga	agttgaagaa	attaggtatc	tgggatgctg	atatgttaga	tcagctgaaa	2760
tatttttgat	ggtctttatt	ggaaactcgag	cgtatacgaag	ataactttaa	acattatttc	2820

55

ttgacagctt	ttgagattga	accagaatgg	attatcgaat	gcgcgtctcg	aagacaaaaa	2880
ttgatgtgata	tggggcaatc	cctcaacott	tatcttgccc	agccagacgg	gaaaaaactg	2940
tcgaatatgt	atttaacggc	ttggaaaaaa	ggtttgaaaa	ctacgtatta	ctcgtagatct	3000
tcacatcgaa	cgaccgttga	aaaatctttt	gtagatatta	ataagagagg	aattcagcct	3060
cgttgatga	gaataaagtc	tgcttcggca	ggaattattg	ttgaagagcg	gaagaagaac	3120
cctgtctgtt	ctttggaaga	agggtgtgaa	gcattgtcag			3159

<210> 83

<211> 4593

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 83

atgagttccg	agaaagatat	aaaaagcacc	tgttctaagt	tttctttgtc	tgtagtagca	60
gctatccttg	cctctgttag	cgggttagct	agttgcgtag	atcttcatgc	tgaggagac	120
ctctgtaaat	agctggtata	tgtaggccct	caagcgggtt	tattgttaga	ccaaattcga	180
gatctattcg	ttgggtctaa	agatagtcag	gctgaaggac	agtagtaggt	aattgttaga	240
gatccaagtt	ctttccaaga	gaaagatgcg	gatactcttc	ccgggaagg	agagcaaatg	300
actttgtctc	cagtaaacca	tcocctgggt	ttccaagggt	tgaccaaca	ggatcaagct	360
tcttccaag	ggttaatttg	tagtttttao	agcagcaacc	ttgattctcc	ctgtgacgga	420
gaattctttt	taggtattgc	ttttgttggg	gatatagtag	agcctggaat	cacattact	480
gaagtgaaag	cttctttgtc	tggaagcgct	ttatatctca	cagaagatct	tatctttgaa	540
aaagattaaag	tgagtttggg	atttgcatac	tgcttctctc	tagaacaggg	gggagctgtt	600
cgagctcaaa	gtattttgat	tcattgattg	caaggaattg	aggttaaaaa	ctgtactaca	660
gcgctgaatg	ctgaggggtc	tagtgcgaa	gatactcttg	gatttggagg	agggcgtctc	720
tttgttcaagg	gttctctttc	tggaagaaaa	agctctcata	tgccctgagg	agatatggtg	780
ttgtggaatt	tgatgggggc	tatatctttt	gaaggaacaa	gcgcgaaact	tgctaatgga	840
ggagcgattg	ctgcctctgg	gaaagtgtct	tttgtcgata	atgataaaaa	gaactctttt	900
atagagaacc	gaagctttgc	tggaaggagc	attgcagcct	ctctcgatat	tgcttttcaa	960
aactcgccag	aactagtttt	caaaaggcaat	tgtagcaattg	gaacagagga	taaaaggtct	1020
ttaggtggag	gggtctatct	ttctctaggc	acogtctttt	tgcaagggaa	tcacgggata	1080
acttgtgata	agaatgagtc	tgcttcgcga	ggaggcgcca	tttttgccaa	aaattgtcag	1140
attttgaca	acagggggcc	agtggttttc	agagatagta	cagcttgctt	agggagaggc	1200
gctattgcag	ctcaagaaat	tgtttclatt	cagaaacaatc	aggctggggt	ttccttcgag	1260
ggaggtaaag	ctagtttcgg	aggggttatt	gcgtgtggat	ctttttcttc	cgaggtgggt	1320
gctctctgtt	tagggacat	tgatatttcg	aagaatttag	gcgcgatttc	gttctctcgt	1380
actttatgta	cgacctcaag	tttagacaaa	atggagtacc	agggaggagg	agctctattt	1440
ggtagaaaata	tttctcttct	tgagaatgct	ggtgtgctca	actttaaaga	caactattgt	1500
aaagactttt	ctctgaattg	gaaaattctg	ggaggaggag	cgatttttag	tactgtgtaa	1560
tggyaaaatta	ctaataattc	ogaaggaaat	ctttttacag	gaaatgcgag	agcttocaac	1620
gctcttccaa	ctcaagagga	gtttctctta	ttcagcaaaa	aagaaggcgc	accactctct	1680
tcaggatatt	ctgggggagg	agcgatttta	ggaagagaa	tagctattct	caacaacgct	1740
gcagtagtat	ttgagcaaaa	tcgtttgcag	tgacgcgaag	aagaagcgac	attattaggt	1800
tggtttggag	gagcgcctgt	tcactgggat	gatagcaact	cgattgttgg	acaactcttc	1860
tgaagatttg	gtaataatta	cgcaatggga	caaggagttc	caggagagag	tcttttatct	1920
aaaaacagtc	agttagctcg	gaatgggaag	gtcgattttt	ctcgaatat	tgctagtttg	1980
ggagagagag	ctcttcaacg	ttctgaagga	aattgtgagc	tagttgataa	cggttgatgt	2040
ctatttcagag	ataatcgagg	gagggtttat	gggggtgcta	tttctgtctt	aogtggagat	2100
gtagtctatt	cttgaaaaaa	gggtagagtt	gaattttaa	acaacatagc	aacacgtctt	2160
tatgtggaa	aaactgtaga	aaaggttgaa	gaggtagacg	cagctcctga	gcaaaaagac	2220
aataatgagc	tttcttctct	agggagagca	gaacagagtt	tttattactg	agcttaataa	2280
gctctttctg	catctgaaga	tggggattta	tcacctgagt	ctccatttct	ttctgaagaa	2340
cttgcgaaaa	gaagagagtg	tgctggagga	gctatttttg	caaaacgggt	tcgtatttga	2400
gatacccaag	agcgccttgt	attctcgaa	aactctctct	atttttatgg	cgcgccaat	2460
tttaacaggt	ctcttcgaga	agaggaata	ttagatgggc	aaatccctga	agttctgtat	2520
tcaggcaatg	caggggatgt	tggttttttc	ggaaattcct	cgaaagctga	tgagcatott	2580
cctcatcacg	gtgggggagc	catttgtact	caaaatttga	cgattttcca	gaataacagg	2640
aatgtctctg	tttatacaaa	cgtagcctgt	tcggggagag	ctttctgat	agaggaatca	2700
ggtaagtgtc	ttttagaagc	ttttggagga	gatattgttt	ttaaaggaaa	ttcttctctt	2760
agagcacaag	gatcogatgc	tatctatttt	acaggttaag	aatcgatat	tcagccctgt	2820
aatgctacgg	aaggacatgc	tattgttttc	cacgcagcat	tagtttttga	aaacttagaa	2880

gaaggaagaa	ctgctgaagt	attgttaatc	aatagtcgag	aaaatccagg	ttacactgga	2940
tcattctgat	ttttagaagc	agaaagtaaa	gttcctcaat	gtattcatgt	acaacaagga	3000
agccttgagt	tgctaaatgg	agccacatta	tgtagttatg	gttttaacaa	agatgtcgga	3060
gcgaagtggg	tattggctgc	tgagactaaa	ctgaagattt	tagattccag	aaactcgtga	3120
caacaaggcg	atgctatcac	taaacctgaa	gcagaaatcg	agtcactctc	tgaaaccagg	3180
gggtgcacatt	ctctttggat	tgccgaagat	gctcaaacaa	cagttctcat	ggttgatcac	3240
catactattt	cttagatttt	agcctccttc	tctctagtc	aaacaggagg	gcagtagaaa	3300
gctcctcagg	ttattgttcc	tgagggaagt	tatgttcgat	ctggagagct	taatttggag	3360
ttagttaaca	caacaggtag	tggtttatga	aatcatgttt	tattgaagaa	tgaggctaaa	3420
tgctccattg	tgtctttcgt	tgcttctggt	gatgaagctt	cagccgaagt	cagtaactcg	3480
tcggtttctg	atttacagat	tcatgtagta	actccagaga	ttgaagaaga	cacatccgct	3540
catatgggag	attggtctga	ggctaaaatt	caagatggaa	ctcttctcat	tagttggaat	3600
cctactggat	atcgattaga	tcctcaaaaa	gcaggggctt	tagtatttaa	tgcatatggg	3660
gaagaaaggg	ctgtcttctc	tgctctgaaa	aatgcaagct	ttgctcaata	tcctactgct	3720
cagcgtatgg	aattcgatta	ttctacaaat	gtgtggggat	tcgctcttgg	tggtttccga	3780
actctatctg	cagagaaatc	gggtgctatt	gatggatata	aaggagctta	tggtggtgct	3840
tcgtctggag	togatattca	attgatggaa	gattttgttc	taggagttag	tgagagctgc	3900
gtcttagtta	aaatggatag	tcagaagttt	gatcgaggag	ttctccggaa	gggagttggt	3960
ggtctgtgat	tacaaggatt	tttagctgga	tctgtgttct	tcaaaggaca	ctatagcctt	4020
tggaacaac	agaaagat	gaaaacgcgt	tatggagtac	taggaagagtc	gagtgctctc	4080
ggagacatctc	gaggagtagt	ggcagatgct	tattgtgaat	accgaagttt	agttgttcc	4140
gtgagaccta	cttttttatgc	tttgcatttc	aatccttatg	togaagatc	ttatgtcttc	4200
atgaattatc	ctggctttac	agaaacaagg	agagaagcgc	gttcttttga	agacgtcttc	4260
cttaacoata	tcaaccattcc	tttagggatg	aagtttgaat	tggtgttcat	aaaaggacag	4320
ttttcagagg	tgaaactctt	gggaataagt	tatgcattgg	aagcttatcg	aaaagtgaag	4380
ggagggcggg	tgacagcttt	agaagctggg	tttgatttgg	agggagctcc	aatggtacct	4440
ctagacagag	agctgggtgt	cgctctggaa	aataaatcgg	aatggagctc	ttacttcagc	4500
agactcttag	gattaaacag	tttttggga	gatttacttt	ctacagatag	taactatgga	4560
tatgaggcga	atactggatt	gogattgata	ttt			4593

<210> 84

<211> 1422

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 84

atgaaaaatta	ttcacacagc	tatcgaattt	gctccggtaa	tcaaaagccg	aggcctggga	60
gaocgcgtat	accgactagc	aaaagcttta	gcgcctaact	acacaacgga	agtggttaac	120
ctctttatacc	ctaaattatt	tactttggcc	aaagaacaag	attctttgctc	gatccaaaaa	180
ttactcttatt	tttttctg	agagcaagaa	gcaactgtct	tctctacttt	tattgaagga	240
attaaagttaa	ctctattcaa	actcgacaca	cagccagagt	tattcgagaa	tgccggaaca	300
atotacacaa	gcgatgatgc	ctctcgtttt	tgccgttttt	ctgctgctgc	ggcctctctac	360
attccaaaaa	aaggagccaa	tatcgtttcat	ttacacagatt	ggcatacagg	attagttgct	420
ggactactca	aaacaacagc	ctgctctcaa	ttacaaaaga	ttgttcttac	ctacataaat	480
ttttggttatc	gagggtatac	aaacagagaa	atattagaag	cctcctcttt	gaatgaattt	540
tatatcagcc	agtacaaact	atttcogcat	acacaaactt	gtgtgttgct	aaaggaagct	600
ttatctctgtt	cagatttctgt	gaactacggtt	tctctacat	acgccaaaag	aattcttgaa	660
tgattatccg	attacgaaat	tcaagatgac	attactgcta	gacaaatca	tctccgctgg	720
atttttaaatg	gaatcgacac	gacaatttgg	gggcctgaaa	cggatcccaa	tttagcgaaa	780
aaactacacta	agaagctttt	cgagacccct	tcaatttttt	ttgaagctaa	agccgagaat	840
aaaaaagcct	tgtagcgaag	attaggcctc	tctttagaac	actcctcctg	ogtgtgcatt	900
atttctagaaa	ttgctgagca	gaagggtcct	caatttatga	aaacggcoat	tctccatgca	960
ctagaaaacag	cttaccgcgt	cattattata	ggtacctgct	acgggaatca	attgcgatga	1020
gaatttgcga	attctccaaga	atcattatgag	aattccctcg	atgtaaagat	tcttttgact	1080
tatagtgatg	tgctggcagc	acaaaatttc	gccgcgcag	atagtactgt	cattctctct	1140
ttttttgaaac	catgtggact	acacaaaatg	attggaatgc	gttacgggac	gttaccogta	1200
tgaaagacta	caggaggact	agcagatact	gtagcaaatg	gaatcaatgg	attttctctc	1260
tttaattccgc	atgaactcta	tgaattccga	aaactgcttt	cggaaagcag	gacaacctac	1320
cgtaccacac	acgacaagtg	gcaacatatt	gtacgtgctt	gtctagattt	ttcttcagac	1380
ctagaaaactg	cgcgcaataa	atatttagaa	atttataaac	aa		1422

<210> 85
 <211> 1179
 <212> DNA
 <213> *Chlamydia trachomatis* serovar D

<400> 85
 atgaaaaaac tottgaaatc ggtattagta ttggccgttt tgagtctctg ttccctccttg 60
 caagctctgc ctgtggggaa tctctgtgaa ccaagcotta tgatcgacgg aattctgtgg 120
 gaaggtttog gggagagatc ttggcatcct tgcgcacatt ggtgtgacgc tatcagcatg 180
 cgtgttggtt actacggaga ctttgttttc gaccgtgttt tgaaaactga tgtgaaataa 240
 gattttcaga ttgggtgccaa gcttacaact gatacaggca atagtgcagc tccatccact 300
 cttacagcaa gagagaatcc tgcctacggc cgacatatgc aggatgctga gatgtttaca 360
 aatgcgcctt gcatggcatt gaattatttg gatcgttttg atgtattctg tacattagga 420
 gccacagctg gatattctaa aggaacctct gcttcttcca atttagtttg attgtttgga 480
 gataatgaaa atcaaaaaaac ggtcaaaagc gagtctgtac caaatatgag ctttgatcaa 540
 tctgtttgtg agttgtatcc agatactact tttgctgga gctgcggcgc tgcgcagact 600
 ttgtgggaat tgggatgtgc aacttttaga gcttcatccc aatatgctca atctaaacct 660
 aaagtgaaga aataaaacgt tctctgcaat gcagcagagt ttactattaa taacacctaaa 720
 gggatgttag gtaaggagtt tctcttgat cttacagcag gaacagatgc tgcgcagga 780
 actaaggagt cctctattga ttaccatgaa tggcaagcaa gtttagctct cttcttaca 840
 ctgaattagt tcactcccta caattggagt aaatggctc gagcaagctt ttagtcgat 900
 acgattctga tagccagcgc aaaaatcagc acagctattt ttgatactac caacgcttaac 960
 ccaactattg ctggagctgg cgaatgtgaa actggcgacg agggctcagc cggagacaca 1020
 atgcaaatcg ttctcttgca attgaacaag atgaaatcta gaaatctgt cggatttgca 1080
 gtgagaaaca ctatttgaga tgcagacaaa taocgcagta cagttgagac tgctgtgac 1140
 gatgagagag cagctcagct aaatgcacaa ttccgcttc 1179

<210> 86
 <211> 585
 <212> DNA
 <213> *Chlamydia trachomatis* serovar D

<400> 86
 atgggatcac tagttggaag acaggctcgc gatttttctg gtaaacgcgt tgtttgtgga 60
 gaagagaaaag aaatctctct agcagacttt cgttgtaagt atgtagtgtc cttcttttat 120
 cctaagaatt ttaoctatgt ttgtcctaca gaattgcagt cttttcaaga tagatttgta 180
 gattttgaaag agcgaggtgc agtctgtgctt ggttgctcgc ttgacgaact tgagacacat 240
 tctgtgtggc tgcctgtagc gagaatgca ggaggaaatg agggaaacga atatcctctg 300
 tttagcagacc cttcttttaa aatatcagaa gcttttggtg ttttgaatcc tgaaggaatg 360
 ctgcctttaa gagagacttt ccttatcgat aaatatgggg ttgttcgtca tgcggttattc 420
 aatgatcttc ctttagggcg ttccattgac gaggaaattg gtattttaga ttcatgtatc 480
 ttctttgaga accacgggaat ggtttgtcca gctaactgco gttctggaga gogtggaatg 540
 gtgccttctg aagagggatt aaaagaatat ttccagaaga tggat 585

<210> 87
 <211> 258
 <212> DNA
 <213> *Chlamydia trachomatis* serovar D

<400> 87
 atgagctaaa ataagaactc tgctttctat cagcctgtga acgtatccgc tgatttagct 60
 gccatcgttg gtcaggagcc tatgcctcgc acagagatca ttaagaaaaat gttgggtattc 120
 attaagaaga atggccttca agatcctaca acaaaacgta atatcaatcc cgaatgataa 180
 ttggtcaaaq tttttgggaa tgaaaaaacct atcgatatgt tccaaatgac aaaaatgggt 240
 tctcaacaca tcaattaaa 258

<210> 88
 <211> 1182
 <212> DNA
 <213> *Chlamydia trachomatis* serovar D

<400> 88
 atgtcaaaaag aaacttttca acgtaataag cctcatatca acatagggac cattggccac 60
 gttgaccatg tgaagactac gttgacagct gctattacgc gtgcgtgtgc ttggagatggg 120
 ttggtcgatt ttctgtatta tagctctatt gacaacactc ctgaagaaaa agctcgcgtg 180
 attacaatta acggttccca cgttgagtac gaacacagctc atogtcaacta cgtccacgtg 240
 gactgcocgt gtcacgtgta ctatgtttaa aacatgatca cgggtgcagc tcaaatggac 300
 gggctgtatt tagtagtttc tgcaacagac ggaagctatg ctcaacttaa agagocattt 360
 ottttggcaa gacaagtgg gggttcctac atogtgtttt ttotcaaataa aattgacatgt 420
 atttcgaag aagacgtgta attggtcgac ttagttaga tggagittgt tgagctctct 480
 gaagagaag gatcaaaagg gtgtccaact atcagaggtt cgtcctcgaa agctttggaa 540
 ggggatgctg catacataga gaaagtctga gagctaatgc agccgtcgga tgataacatc 600
 cctactccag aaagagaaat tgacaagcct ttctaatgc ctatttgagg cgtattctct 660
 atctccggac aggaactgt agtaactgga cgtattgagc gtggaattgt taaagtttcc 720
 gataaagtct agttgtgtcg tottagagat actaaagaaa cgtattgtac tgggggttaa 780
 atgttcagaa aagaactccc agaaggtcgt gcaggagaga acgttggatt gctcctcaga 840
 ttgtattgta agaactgtt ggaagaggga atggtgtttt gottgcaaaa caagtgttaa 900
 cctcatcacac tgcctcaagtg tgcgttttac gttttgcaaa aagaagaagg tggcgcacat 960
 aagccttctc tcaccaggata tagacctcaa ttcttcttcc gtacaacaga cgtcacaggt 1020
 ggtgtaactc tgctgaggg aattgagatg gtoatgcctg gggataacgt tgagtttgaa 1080
 gtcacatga ttagccctgt ggctttagaa gaaggtatga gatttgcgat tcgtgaaggt 1140
 ggtcgtacaa tcggtgctgg aactatttct aagatcattg caa 1182

<210> 89
 <211> 246
 <212> DNA
 <213> Chlamydia trachomatis serovar D

<400> 89
 atggggcgaag atcaaccgaag aaaatttctt aagaaagtat ottttgtaaa aaaacaagca 60
 gcttttgcgg gtaactttat ogaagaaatt aagaagattg agtgggtaaa taacggsagt 120
 ottaaaagat acgtcaagat tgttttgatg aaatttttg gctttgagtt ttccatctat 180
 tgtgtggatt tagctcttcg aaagtccctt tcattgttgc gtaaaagtaac aagotttttc 240
 tttggt 246

<210> 90
 <211> 1137
 <212> DNA
 <213> Chlamydia trachomatis serovar D

<400> 90
 atggtgatcc ctaaggtgga tctaggagaa agtgccgtca tgaagggtta caagcttaact 60
 togcaacttg ctaactgttc gatcttattg acttccacc atactatgg tcatgcaagt 120
 cagatgagcc aaactcttcc tactattata gaagcacaag cggaagaggg attgagcgtt 180
 gacaggggag ttgctggaca ggotctttaa aaactctgta aaaaagatg tgcctctaga 240
 aaactctgat gtaaggcttc ttttaagaaa aaggatttct ttcttgtat tacaagtga 300
 ttgttctctg gaaatcatga gcagcgttta actgcgaaaa aagagaacaa ggctcaggtt 360
 aaagagctcc gagttagtgt tcaaacgact aaaaaagcac aaataactca gtctgagaaa 420
 gaatttttgc attgtctatg taatagttaa agagaaagaa agctctctca gaataagctt 480
 gtaaatctct ctctgtctaa gagtgaagaa ttgagctcta aagaagcagc aatagctct 540
 gctcagagctt ctcttctcc agaagaaaaa cgtcaattga ttogtgagtg gttagcagaa 600
 gaaagagctg ctctgtaact tgggcgttgc gcttgtgcgg taagttagaa tcttaagaa 660
 gacggaagta ttaactctac attgcgtcat gatgcggaga aagotttgac agactgtga 720
 aaacgcaatg aatattctgt aaatgctaga gcaagacaac gacgcgtctt tcaaaaagcc 780
 aagaaagcaa agacggagaa aactgaggtc gatgagaaag ctgcagaagc gttgcgcga 840
 gctccaacca aacaggcgca taaggagcca gagaattact tgcagctac agctcttca 900
 aataactact atgttatgtc ctatctaaat gctcatcaat acogttgtga ttctcggag 960
 acggagctgg cttgctcttc ttgtgttaag aaacgcogag ctaacttcgg tattttctg 1020
 tgtactatgg tggttacogt cattgctatg atcgttagag gcttatcat ttctaatgt 1080
 acagactcta cgttgcggg ctctcgtgga acaggagagg gaggctcaac gcaacca 1137

<210> 91

<211> 1689

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 91

atgggtttatt ttagagctca tcaacctagc catagccta aaacatttcc ttgggaagtt 60
 caccatttgt tctcgataa gcatcctcaa attgctaaag ctatgcggat tacggggata 120
 gccttcgcag ctctatctct gctcgtctgt gtctgctcgc ttattgtcgt ctctgcggga 180
 ggagctgcga tctctcttgc tgtcattagt ggaattgctg taatgtctgg cctcttatcc 240
 gctgcacca tttctgttct tgcacaaagc gctttggctc aacgaaacaa aaaacacata 300
 gaagagctgc tctcgtttag taatgcgacc gagcatgtga gttacctgac ctacgacacc 360
 tctttattta atcaatggga atccttaggt gctctaata agcagttgtc tcagattgac 420
 ttaactattc aagctccgga aaaaaaacta ttaaaagaag ttcttggctc cagatacagat 480
 tccaattatc actccatcga agagatctcc gatcgttta cgaataatgt cctctcttct 540
 cgattaaagag aacattttta tggaggagaa gagcgttatg ccccttattt aagcctctct 600
 ctacttaaca agaatttgtt gctgacccaa atcacatcaa atatgattag gatgctacca 660
 aactccgggtg tgttttttll cctcaaaagc aatacaactaa gtcacgcaag cgcacacata 720
 tatacagtat taaaagtgcg tttatcctta ggagttctcg ctggagctgc tgcctttatc 780
 atctttcttc cctctagcct gctttttatc gctgttatag gagtatcttc cttagcattg 840
 ggagtgcaat ctctctctat gatctggggc attaagtatt tgcctgaaca tctctctctc 900
 aatagaagag aactagctaa agatattcaa aaaaccattg gccacagatg ctgggctctc 960
 atggctcatt aacagcatca attactatca catctacatg aaactctatt agatgaagcc 1020
 atcacacatt gatggagaga gctctctctt attgaacacg ctactcttaa ttcacacaaa 1080
 gaagatttga caaaacaata tgcataattg aaacgcagct ttaataaact ttcaataaac 1140
 gatgagcgcc tctcgtctca attagagaaa cgagcttaact tatctccaat tcttaataac 1200
 ggcgaaaatg ctaaaactaa agaactgcag ctcttagact cagaaaatga tccaattctc 1260
 gaatttccag agattataaa taaagactaa gaagctgcga ataaacagc agctgacgct 1320
 aagtcacaaat ctatacggg agacgaaacc tctgacaaaa tattctctat atggaaaccc 1380
 acaaaagaact tgcgactaga agatttctgg agagtgcatg aagcttgcaa tgaagagcaa 1440
 caagctctcc tcttagaaga ttatatgagt tataaaactc cagaatgtca agctgcaatc 1500
 caaaaagtga gtcaagaact gaaggcgcca caaaatcat tcgcagctcc agaaaagcat 1560
 gctctagaca gatcttatga atccagtgtg gccacgatgt atttagctg agcgaatcaa 1620
 gaacacaccc gctctctgaa catctctctc gaattacac aactagcaca atacctgtta 1680
 gataatcac 1689

<210> 92

<211> 1074

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 92

gtgcgtaaaa ctgtcattgt tgcctatgtt ggaggagtg attcctcggt tgttgcttat 60
 ctcttaaaaa agcaaggagg gtataatggt gtgtggctct tcatgaaaaa ttggggagag 120
 caggacgaga atggtgagtg taactgcaacc aaagatttct gcgatglaga gggagctgca 180
 gaacaaatgt caattccata ttacacagtt tctctttcta aggaataaaa agagcgagtg 240
 ttttttagat ttctaagaga atatgogaac ggcatacact ccaatcctga tgttttatgc 300
 aatgcagaaa tcaaatttga tttattacag aagaaggtac gtgagctaaa aggtgatttt 360
 tttagccacgg gacattattg toaggagggg gctgatggaa ctggtttgtc cagaggaata 420
 gaccoccatca aagaccaaag ttatttctta tgtggcactc ctaaggatgc tttatccaat 480
 gtacttttcc cactgggagg tatgtataaa acggaggtag ctgcaattgc tcaagaagct 540
 ggttttagct cccgccaaaa aaaaatagtc acaggggatt gcttacttgg taaacggcct 600
 tttaagagtt tctttagaca gtttgtagca gactctcctg gagacattat gatctttgat 660
 acacacacag tagtcggcgg acatgaagga gccattattt atacgattgg acagcgtgca 720
 ggggttaaaa taggaggaat ggaaaagcct tgttatgttc tttagcaaga tatggaaaaa 780
 aatattgttt acattgtaag ggggtgaagt catcctttac tttatgcaga agagctttta 840
 gctaaaggaac ttaattgggt tgttcccttg caggagccta tgcctctgag tctcaaaagt 900
 cggtaacagat cccctgaaga gaaatgttct gtatatcctt tggaagatgg aacggtaaaa 960
 gtgattttcg atgtccctgt gaaagctgtc accctgggac agactgtagc tttctaccag 1020
 ggggacattt gtttaggagg aggagtgatt gaagtgccta tgattcctaa gctg 1074

<210> 93

<211> 801
 <212> DNA
 <213> Chlamydia trachomatis serovar D

<400> 93
 atgtccagaa aacccggcttc taactcatcc cggaacacca aacggtctctc agacacttcc 60
 ttgggaagtc tggcccaaga ttataataaa gccgttgatc gcatgggaca ttctatcat 120
 aaggaaagta ttctccctaa tctcctttct aagctacata ttcccgctcc atcgctctctg 180
 gttgatgtag gatgtgggtca agggattttg gagaagcatt taccocaaaca tctcccttat 240
 ctaggaaatcg attcttccccc tagctctgctg cgtttttgcaa agaaaaagcgc ttccctaaaaa 300
 tcaogctgctt ttcttcatca cgtatgacgc caacoggtac cagcagatca tcatgagcag 360
 ttctcccatg ctacagcaat ccttctctct cagaatatg aaatctccaga acaagctctac 420
 gcaacacacag cgaattcttt ggctccctcaa ggtagggtgt ttattgttct caacctcca 480
 tgccttcgca tccctaggct ttcttcatg cttattgat agcctaaaaa actcttatct 540
 agaaaaatag accgctatct ctctccctg ggggttctca tctgtttgca tctgtggaga 600
 aaacattctg agcagacata ttctttccat ttccctctaa gctattgggt acaagcttta 660
 tctaactaca attctctgat tgaatgatg gaagaatgga tctccctcaa aaatctcca 720
 gggaagaggg ctgcagcaga aaatctttgt cgcaaggagt ttccgctttt ctgtgtttac 780
 tcagcattaa aaatatcaaa a 801

<210> 94
 <211> 2601
 <212> DNA
 <213> Chlamydia trachomatis serovar D

<400> 94
 atggagaat ttccagatgc agtaagcgaa gccttagaaa aggcgttttga gttagotaaa 60
 aactctaaag attctacgtg gacagaaaac catttctgca aaagtctttt gcaaaatcca 120
 gggtccctat ttgttttgggt cattaagagt gtgcaacgta attcttggttt gtttaactct 180
 gctgtggagc agcccttacc cagagaaacca actgtagtgc agggaacgcg ttgtgtactg 240
 cctctctcaa gtctacagca gttgttgctc aatgcgcctc aagaagctag aagtatgggt 300
 gacgaatata tatcagggga tcatittgta ctagcttttt ggcatcgac taaagagcct 360
 ttgtctctct ggagaaaaac tgtaaaaact acctctgaag cgttgaaaaa attaatctac 420
 aaattaagac aaggaaagtg tatggactca cctagtgtgc aagaaaaatc gaaagatta 480
 gagaataact gcaaaaattt gactgtactt gcaagagagc gcaagcttga tctgtgatt 540
 ggctcgatg gtaggaattag acgtacgata caggttcttt ctgacgcaac aaagaataat 600
 cctatgttga taggggagcc cggagtgtgg aaaaacgcaa tgcgtgaagc acttgtctct 660
 cgcatagtgc aaggggatgt tccagagagt ttcaaggaaa agcatctgta tgtactggat 720
 atgggagcctt tgaattgcagg tgcacagat cgaggagagt ttgaagagcg gttaaaaagt 780
 gttatgaagg gtgtagaagg ttctgaagcg gactgtatcc tattoattga tgaagtgc 840
 actttagtag gtagggggagc tacagatgga gctatggatg cagcgaatct attaaagcct 900
 gcttagcac gaggcacttt gcatgttatt ggccctacga ctttgatga atacocaaaa 960
 tatatagaga aagacgcggc ttgtgaacgg cgtttccagc ctatttttgc aacagaacct 1020
 tctttggaa atctgttatt caattctccg gggtttaagg gacttaattg actttttct 1080
 gggtgtcgca ttacagaagg ggctttgaat gcagctgtag ttctttctta tctgttacct 1140
 acagacccgat ttctctctga taaggcgatt gacctaattg atgaggtgc gagttaact 1200
 ctatcgcaaa taggaagttt acctctgctc attgatgaaa aggaagagga attatcagct 1260
 tgaatcgtga aacaagaagc tatataacgc gagcaagcac agccttata ggaagagct 1320
 gaagacatgc aaaaagcaat tgacoggggt aaggaagagc tggccgcttt acgcttgcgc 1380
 tggagtgaag aaaaagattt aattacagga ttaaaaagaa agagaatgc tttagaaaa 1440
 ttaaaatttg ccgaagagga agctgagcgt actgccgatt acaatccggt ggcagaaata 1500
 cgtatagttt tttgactctc tttagggaaa gaattctact tagctgagga aagtttaatt 1560
 caaagagatg ggcgcctgct tcaagaggaa gttgatgagc ggttgattgc gcaagttgt 1620
 gcaatttggc ctggaatccc tgtgcacaaa atgttggagg gagaactga agatttatg 1680
 gtgttggagg agtctttaga agaaaggttt gtgtgacac ctttgcctat tgcgcagct 1740
 agtgatctga tctgagctgc tgcagtagga ttgagtgc ttgagtac cgcagctctc tagctgagc 1800
 ttcttatttc ttggcaactc aggggttagg aaaactgagc ttgctaagcg attagcagag 1860
 cttttattta ataaggaaag agcogatgatt cgttttgaca aggtttgaaa tatggaaaa 1920
 cactctcggt ccaaatgtat aggatctctc ccagggtagt taggatata gaaagagag 1980
 agtctctcag aagctttaag aagacgacct ttctctgtt ttcttttga tgaagttaga 2040
 aaagcagata aagaagtatt taatttttta ttgcagattt ttgatgatg gattcttacc 2100

```

gatagcaaga agcgtgaaggt aaattgtaag aatgctcttt tcattatgac atcaaatatt 2160
ggttcgcaag agcttgctga ttattgtact aagaagggaa ctatogtaga caaagaagct 2220
gtgctatctg ttgttgcccc tgcgcttaaa aattatttta gtccagaatt tatcaatcgt 2280
atcgtatgaca ttctgctctt cgctcctttg actacggaaag acattgtaaa aattgtcggt 2340
attcaaatga atcggggttg tttacgtttg ctggaaagaa aaatttcgtt aacttgggat 2400
gattotttag tgctatttct cagttagcaa gggtatgaca gogcttttgg agctcgccot 2460
ctgaagcggt tgatacagca aaaagtagtg actatgtgtg cttaaagctct ttgaaaagga 2520
gatacaaaao ctggaatggc ggtggagctt actatggcaa aagatgtagt tgtgtttaaa 2580
attaaaacaa atccagctgt g

```

<210> 95

<211> 1016

<212> PRT

<213> *Chlamydia trachomatis* serovar D

<400> 95

```

Met Pro Phe Ser Leu Arg Ser Thr Ser Phe Cys Phe Leu Ala Cys Leu
Cys Ser Tyr Ser Tyr Gly Phe Ala Ser Ser Pro Gln Val Leu Thr Pro
20 25 30
Asn Val Thr Thr Pro Phe Lys Gly Asp Asp Val Tyr Leu Asn Gly Asp
35 40 45
Cys Ala Phe Val Asn Val Tyr Ala Gly Ala Glu Asn Gly Ser Ile Ile
50 55 60
Ser Ala Asn Gly Asp Asn Leu Thr Ile Thr Gly Gln Asn His Thr Leu
65 70 75 80
Ser Phe Thr Asp Ser Gln Gly Pro Val Leu Gln Asn Tyr Ala Phe Ile
85 90 95
Ser Ala Gly Glu Thr Leu Thr Leu Lys Asp Phe Ser Ser Leu Met Phe
100 105 110
Ser Lys Asn Val Ser Cys Gly Glu Lys Gly Met Ile Ser Gly Lys Thr
115 120 125
Val Ser Ile Ser Gly Ala Gly Glu Val Ile Phe Trp Asp Asn Ser Val
130 135 140
Gly Tyr Ser Pro Leu Ser Ile Val Pro Ala Ser Thr Pro Thr Pro Pro
145 150 155 160
Ala Pro Ala Pro Ala Pro Ala Ala Ser Ser Ser Leu Ser Pro Thr Val
165 170 175
Ser Asp Ala Arg Lys Gly Ser Ile Phe Ser Val Glu Thr Ser Leu Glu
180 185 190
Ile Ser Gly Val Lys Lys Gly Val Met Phe Asp Asn Asn Ala Gly Asn
195 200 205
Phe Gly Thr Val Phe Arg Gly Asn Ser Asn Asn Asn Ala Gly Ser Gly
210 215 220
Gly Ser Gly Ser Ala Thr Thr Pro Ser Phe Thr Val Lys Asn Cys Lys
225 230 235 240
Gly Lys Val Ser Phe Thr Asp Asn Val Ala Ser Cys Gly Gly Gly Val
245 250 255
Val Tyr Lys Gly Thr Val Leu Phe Lys Asp Asn Glu Gly Gly Ile Phe
260 265 270
Phe Arg Gly Asn Thr Ala Tyr Asp Asp Leu Gly Ile Leu Ala Ala Thr
275 280 285
Ser Arg Asp Gln Asn Thr Gly Thr Gly Gly Gly Gly Val Ile Cys
290 295 300
Ser Pro Asp Asp Ser Val Lys Phe Glu Gly Asn Lys Gly Ser Ile Val
305 310 315 320
Phe Asp Tyr Asn Phe Ala Lys Gly Arg Gly Gly Ser Ile Leu Thr Lys
325 330 335
Glu Phe Ser Leu Val Ala Asp Asp Ser Val Val Phe Ser Asn Asn Thr
340 345 350
Ala Glu Lys Gly Gly Gly Ala Ile Tyr Ala Pro Thr Ile Asp Ile Ser

```

	355					360				365				
Thr	Asn	Gly	Gly	Ser	Ile	Leu	Phe	Glu	Arg	Asn	Arg	Ala	Ala	Glu
	370					375				380				
Gly	Ala	Ile	Cys	Val	Ser	Glu	Ala	Ser	Ser	Gly	Ser	Thr	Gly	Asn
385						390				395				400
Thr	Leu	Ser	Ala	Ser	Asp	Gly	Asp	Ile	Val	Phe	Ser	Gly	Asn	Met
				405					410					415
Ser	Asp	Arg	Pro	Gly	Glu	Arg	Ser	Ala	Ala	Arg	Ile	Leu	Ser	Asp
			420					425				430		
Thr	Thr	Val	Ser	Leu	Asn	Ala	Ser	Gly	Leu	Ser	Lys	Leu	Ile	Phe
			435					440				445		
Asp	Pro	Val	Val	Gln	Asn	Asn	Ser	Ala	Ala	Gly	Ala	Ser	Thr	Pro
	450					455					460			
Pro	Ser	Ser	Ser	Ser	Met	Pro	Gly	Ala	Val	Thr	Ile	Asn	Gln	Ser
465					470					475				480
Asn	Gly	Ser	Val	Ile	Phe	Thr	Ala	Glu	Ser	Leu	Thr	Pro	Ser	Glu
				485					490					495
Leu	Gln	Val	Leu	Asn	Ser	Thr	Ser	Asn	Phe	Pro	Gly	Ala	Leu	Thr
			500					505					510	
Ser	Gly	Gly	Glu	Leu	Val	Val	Thr	Glu	Gly	Ala	Thr	Leu	Thr	Thr
			515					520				525		
Thr	Ile	Thr	Ala	Thr	Ser	Gly	Arg	Val	Thr	Leu	Gly	Ser	Gly	Ala
	530					535				540				
Leu	Ser	Ala	Val	Ala	Gly	Ala	Ala	Asn	Asn	Asn	Tyr	Thr	Cys	Thr
545					550					555				560
Ser	Lys	Leu	Gly	Ile	Asp	Leu	Glu	Ser	Phe	Leu	Thr	Pro	Asn	Tyr
				565					570					575
Thr	Ala	Ile	Leu	Gly	Ala	Asp	Gly	Thr	Val	Thr	Val	Asn	Ser	Gly
			580						585				590	
Thr	Leu	Asp	Leu	Val	Met	Glu	Ser	Glu	Ala	Glu	Val	Tyr	Asp	Asn
	595					600						605		Pro
Leu	Phe	Val	Gly	Ser	Leu	Thr	Ile	Pro	Phe	Val	Thr	Leu	Ser	Ser
	610					615						620		
Ser	Ala	Ser	Asn	Gly	Val	Thr	Lys	Asn	Ser	Val	Thr	Ile	Asn	Asp
625					630					635				640
Asp	Ala	Ala	His	Tyr	Gly	Tyr	Gln	Gly	Ser	Trp	Ser	Ala	Asp	Trp
				645						650				655
Lys	Pro	Pro	Leu	Ala	Pro	Asp	Ala	Lys	Gly	Met	Val	Pro	Pro	Asn
			660					665					670	
Asn	Asn	Thr	Leu	Tyr	Leu	Thr	Trp	Arg	Pro	Ala	Ser	Asn	Tyr	Gly
			675				680					685		
Tyr	Arg	Leu	Asp	Pro	Gln	Arg	Lys	Gly	Glu	Val	Pro	Asn	Ser	Leu
	690					695						700		
Trp	Val	Ala	Gly	Ser	Ala	Leu	Arg	Thr	Phe	Thr	Asn	Gly	Leu	Lys
				710					715					720
His	Tyr	Val	Ser	Arg	Asp	Val	Gly	Phe	Val	Ala	Ser	Leu	His	Ala
				725					730					735
Gly	Asp	Tyr	Ile	Leu	Asn	Tyr	Thr	Gln	Asp	Asp	Arg	Gly	Phe	Leu
			740					745				750		
Ala	Arg	Tyr	Gly	Gly	Phe	Gln	Ala	Thr	Ala	Ala	Ser	His	Tyr	Glu
			755				760					765		Asn
Gly	Ser	Ile	Phe	Gly	Val	Ala	Phe	Gly	Gln	Leu	Tyr	Gly	Gln	Thr
				770			775					780		Lys
Ser	Arg	Met	Tyr	Tyr	Ser	Lys	Asp	Ala	Gly	Asn	Met	Thr	Met	Leu
					790					795				800
Cys	Phe	Gly	Arg	Ser	Tyr	Val	Asp	Ile	Lys	Gly	Thr	Glu	Thr	Val
				805					810					815
Tyr	Trp	Glu	Thr	Ala	Tyr	Gly	Tyr	Ser	Val	His	Arg	Met	His	Thr
			820					825					830	Gln
Tyr	Phe	Asn	Asp	Lys	Thr	Gln	Lys	Phe	Asp	His	Ser	Lys	Cys	His
			835				840							845

His Asn Asn Asn Tyr Tyr Ala Phe Val Gly Ala Glu His Asn Phe Leu
 850 855 860
 Glu Tyr Cys Ile Pro Thr Arg Gln Phe Ala Arg Asp Tyr Glu Leu Thr
 865 870 875 880
 Gly Phe Met Arg Phe Glu Met Ala Gly Gly Trp Ser Ser Ser Thr Arg
 885 890 895
 Glu Thr Gly Ser Leu Thr Arg Tyr Phe Ala Arg Gly Ser Gly His Asn
 900 905 910
 Met Ser Leu Pro Ile Gly Ile Val Ala His Ala Val Ser His Val Arg
 915 920 925
 Arg Ser Pro Pro Ser Lys Leu Thr Leu Asn Met Gly Tyr Arg Pro Asp
 930 935 940
 Ile Trp Arg Val Thr Pro His Cys Asn Met Glu Ile Ala Asn Gly
 945 950 955 960
 Val Lys Thr Pro Ile Gln Gly Ser Pro Leu Ala Arg His Ala Phe Phe
 965 970 975
 Leu Glu Val His Asp Thr Leu Tyr Ile His His Phe Gly Arg Ala Tyr
 980 985 990
 Met Asn Tyr Ser Leu Asp Ala Arg Arg Gln Thr Ala His Phe Val
 995 1000 1005
 Ser Met Gly Leu Asn Arg Ile Phe
 1010 1015

<210> 96

<211> 346

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 96

Met Gln Ala Asp Ile Leu Asp Gly Lys Gln Lys Arg Val Asn Leu Asn
 5 10 15
 Ser Lys Arg Leu Val Asn Cys Asn Gln Val Asp Val Asn Gln Leu Val
 20 25 30
 Pro Ile Lys Tyr Lys Trp Ala Trp Glu His Tyr Leu Asn Gly Cys Ala
 35 40 45
 Asn Asn Trp Leu Pro Thr Gly Ile Pro Met Gly Lys Asp Ile Glu Leu
 50 55 60
 Trp Lys Ser Asp Arg Leu Ser Glu Asp Glu Arg Arg Val Ile Leu Leu
 65 70 75 80
 Asn Leu Gly Phe Phe Ser Thr Ala Glu Ser Ser Leu Val Gly Asn Asn Ile
 85 90 95
 Val Leu Ala Ile Phe Lys His Val Thr Asn Pro Glu Ala Arg Gln Tyr
 100 105 110
 Leu Leu Arg Gln Ala Phe Glu Glu Ala Val His Thr His Thr Phe Leu
 115 120 125
 Tyr Ile Cys Glu Ser Leu Gly Leu Asp Glu Lys Glu Ile Phe Asn Ala
 130 135 140
 Tyr Asn Glu Arg Ala Ala Ile Lys Ala Lys Asp Asn Phe Gln Met Glu
 145 150 155 160
 Ile Thr Gly Lys Val Leu Asp Pro Asn Phe Arg Thr Asp Ser Val Glu
 165 170 175
 Gly Leu Gln Glu Phe Val Lys Asn Leu Val Gly Tyr Tyr Ile Ile Met
 180 185 190
 Glu Gly Ile Phe Phe Tyr Ser Gly Phe Val Met Ile Leu Ser Phe His
 195 200 205
 Arg Gln Asn Lys Met Ile Gly Ile Gly Glu Gln Tyr Gln Tyr Ile Leu
 210 215 220
 Arg Asp Glu Thr Ile His Leu Asn Phe Gly Ile Asp Leu Ile Asn Gly
 225 230 235 240
 Ile Lys Glu Glu Asn Pro Glu Ile Trp Thr Pro Glu Leu Gln Gln Glu

```

                245                250                255
Ile Val Glu Leu Ile Lys Arg Ala Val Asp Leu Glu Ile Glu Tyr Ala
                260                265                270
Gln Asp Cys Leu Pro Arg Gly Ile Leu Gly Leu Arg Ala Ser Met Phe
                275                280                285
Ile Asp Tyr Val Gln His Ile Ala Asp Arg Arg Leu Glu Arg Ile Gly
                290                295                300
Leu Lys Pro Ile Tyr His Thr Lys Asn Pro Phe Pro Trp Met Ser Glu
305                310                315                320
Thr Ile Asp Leu Asn Lys Glu Lys Asn Phe Phe Glu Thr Arg Val Ile
                325                330                335
Glu Tyr Gln His Ala Ala Ser Leu Thr Trp
                340                345

```

<210> 97

<211> 1053

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 97

```

Met Phe Thr Arg Ile Val Met Val Asp Leu Gln Glu Lys Gln Cys Thr
                5                10                15
Ile Val Lys Arg Asn Gly Met Phe Val Pro Phe Asp Arg Asn Arg Ile
                20                25                30
Phe Gln Ala Leu Glu Ala Ala Phe Arg Asp Thr Arg Arg Ile Asp Asp
                35                40                45
His Met Pro Leu Pro Glu Asp Leu Glu Ser Ser Ile Arg Ser Ile Thr
                50                55                60
His Gln Val Val Lys Glu Val Val Gln Lys Ile Thr Asp Gly Gln Val
                65                70                75                80
Val Thr Val Glu Arg Ile Gln Asp Met Val Glu Ser Gln Leu Tyr Val
                85                90                95
Asn Gly Leu Gln Asp Val Ala Arg Asp Tyr Ile Val Tyr Arg Asp Asp
                100                105                110
Arg Lys Ala His Arg Lys Lys Ser Trp Gln Ser Leu Ser Val Val Arg
                115                120                125
Arg Cys Gly Thr Val Val His Phe Asn Pro Met Lys Ile Ser Ala Ala
                130                135                140
Leu Glu Lys Ala Phe Arg Ala Thr Asp Lys Thr Glu Gly Met Thr Pro
                145                150                155                160
Ser Ser Val Arg Glu Glu Ile Asn Ala Leu Thr Gln Asn Ile Val Ala
                165                170                175                180
Glu Ile Glu Glu Cys Cys Pro Gln Gln Asp Arg Arg Ile Asp Ile Glu
                185                190                195
Lys Ile Gln Asp Ile Val Glu Gln Gln Leu Met Val Val Gly His Tyr
                200                205                210
Ala Val Ala Lys Asn Tyr Ile Leu Tyr Arg Glu Ala Arg Val
                215                220                225
Arg Asp Asn Arg Glu Glu Asp Gly Ser Thr Glu Lys Thr Ile Ala Glu
                230                235                240
Glu Ala Val Glu Val Leu Ser Lys Asp Gly Ser Thr Tyr Thr Met Thr
                245                250                255
His Ser Gln Leu Leu Ala His Leu Ala Arg Ala Cys Ser Arg Phe Pro
                260                265                270
Glu Thr Thr Asp Ala Ala Leu Leu Thr Asp Met Ala Phe Ala Asn Phe
                275                280                285
Tyr Ser Gly Ile Lys Glu Ser Glu Val Val Leu Ala Cys Ile Met Ala
                290                295                300
Ala Arg Ala Asn Ile Glu Lys Glu Pro Asp Tyr Ala Phe Val Ala Ala
305                310                315                320

```

Glu Leu Leu Leu Asp Val Val Tyr Lys Glu Ala Leu Gly Lys Ser Lys
 325 330 335
 Tyr Ala Glu Asp Leu Glu Gln Ala His Arg Asp His Phe Lys Arg Tyr
 340 345 350
 Ile Ala Glu Gly Asp Thr Tyr Arg Leu Asn Ala Glu Leu Lys His Leu
 355 360 365
 Phe Asp Leu Asp Ala Leu Ala Asp Ala Met Asp Leu Ser Arg Asp Leu
 370 375 380
 Gln Phe Ser Tyr Met Gly Ile Gln Asn Leu Tyr Asp Arg Tyr Phe Asn
 385 390 395 400
 His His Glu Gly Cys Arg Leu Glu Thr Pro Gln Ile Phe Trp Met Arg
 405 410 415
 Val Ala Met Gly Leu Ala Leu Asn Glu Gln Asp Lys Thr Ser Trp Ala
 420 425 430
 Ile Thr Phe Tyr Asn Leu Leu Ser Thr Phe Arg Tyr Thr Pro Ala Thr
 435 440 445
 Pro Thr Leu Phe Asn Ser Gly Met Arg His Ser Gln Leu Ser Ser Cys
 450 455 460
 Tyr Leu Ser Thr Val Gln Asp Asn Leu Val Asn Ile Tyr Lys Val Ile
 465 470 475 480
 Ala Asp Asn Ala Met Leu Ser Lys Trp Ala Gly Gly Ile Gly Asn Asp
 485 490 495
 Trp Thr Ala Ile Arg Ala Thr Gly Ala Leu Ile Lys Gly Thr Asn Gly
 500 505 510
 Arg Ser Gln Gly Val Ile Pro Phe Ile Lys Val Thr Asn Asp Thr Ala
 515 520 525
 Val Ala Val Asn Gln Gly Gly Lys Arg Lys Gly Ala Val Cys Val Tyr
 530 535 540
 Leu Glu Val Trp His Leu Asp Tyr Glu Asp Phe Leu Glu Leu Arg Lys
 545 550 555 560
 Asn Thr Gly Asp Glu Arg Arg Arg Ala His Asp Val Asn Ile Ala Ser
 565 570 575
 Trp Ile Pro Asp Leu Phe Phe Lys Arg Leu Gln Gln Lys Gly Thr Trp
 580 585 590
 Thr Leu Phe Ser Pro Asp Asp Val Pro Gly Leu His Asp Ala Tyr Gly
 595 600 605
 Glu Glu Phe Glu Arg Leu Tyr Glu Glu Tyr Glu Arg Lys Val Asp Thr
 610 615 620
 Gly Glu Ile Arg Leu Phe Lys Lys Val Glu Ala Glu Asp Leu Trp Arg
 625 630 635 640
 Lys Met Leu Ser Met Leu Phe Glu Thr Gly His Pro Trp Met Thr Phe
 645 650 655
 Lys Asp Pro Ser Asn Ile Arg Ser Ala Gln Asp His Lys Gly Val Val
 660 665 670
 Arg Cys Ser Asn Leu Cys Thr Glu Ile Leu Leu Asn Cys Ser Glu Thr
 675 680 685
 Glu Thr Ala Val Cys Asn Leu Gly Ser Ile Asn Leu Val Gln His Ile
 690 695 700
 Val Gly Asp Gly Leu Asp Glu Glu Lys Leu Ser Glu Thr Ile Ser Ile
 705 710 715 720
 Ala Val Arg Met Leu Asp Asn Val Ile Asp Ile Asn Phe Tyr Pro Thr
 725 730 735
 Lys Glu Ala Lys Glu Ala Asn Phe Ala His Arg Ala Ile Gly Leu Gly
 740 745 750
 Val Met Gly Phe Gln Asp Ala Leu Tyr Lys Leu Asp Ile Ser Tyr Ala
 755 760 765
 Ser Gln Glu Ala Val Glu Phe Ala Asp Tyr Ser Ser Glu Leu Ile Ser
 770 775 780
 Tyr Tyr Ala Ile Gln Ala Ser Cys Leu Leu Ala Lys Glu Arg Gly Thr
 785 790 795 800
 Tyr Ser Ser Tyr Lys Gly Ser Lys Trp Asp Arg Gly Leu Leu Pro Ile

805 810 815
 Asp Thr Ile Gln Leu Leu Ala Asn Tyr Arg Gly Glu Ala Asn Leu Gln
 820 825 830
 Met Asp Thr Ser Ser Arg Lys Asp Trp Glu Pro Ile Arg Ser Leu Val
 835 840 845
 Lys Glu His Gly Met Arg His Cys Gln Leu Met Ala Ile Ala Pro Thr
 850 855 860
 Ala Thr Ile Ser Asn Ile Ile Gly Val Thr Gln Ser Ile Glu Pro Thr
 865 870 875 880
 Tyr Lys His Leu Phe Val Lys Ser Asn Leu Ser Gly Glu Phe Thr Ile
 885 890 895
 Pro Asn Val Tyr Leu Ile Glu Lys Leu Lys Lys Leu Gly Ile Trp Asp
 900 905 910
 Ala Asp Met Leu Asp Asp Leu Lys Tyr Phe Asp Gly Ser Leu Leu Glu
 915 920 925
 Ile Glu Arg Ile Pro Asp His Leu Lys His Ile Phe Leu Thr Ala Phe
 930 935 940
 Glu Ile Glu Pro Glu Trp Ile Ile Glu Cys Ala Ser Arg Arg Gln Lys
 945 950 955 960
 Trp Ile Asp Met Gly Gln Ser Leu Asn Leu Tyr Leu Ala Gln Pro Asp
 965 970 975
 Gly Lys Lys Leu Ser Asn Met Tyr Leu Thr Ala Trp Lys Lys Gly Leu
 980 985 990
 Lys Thr Thr Tyr Tyr Leu Arg Ser Ser Ala Thr Thr Val Glu Lys
 995 1000 1005
 Ser Phe Val Asp Ile Asn Lys Arg Gly Ile Gln Pro Arg Trp Met Lys
 1010 1015 1020
 Asn Lys Ser Ala Ser Ala Gly Ile Ile Val Glu Arg Ala Lys Lys Ala
 1025 1030 1035 1040
 Pro Val Cys Ser Leu Glu Glu Gly Cys Glu Ala Cys Gln
 1045 1050

<210> 98
 <211> 1531
 <212> PRT
 <213> *Chlamydia trachomatis* serovar D

<400> 98
 Met Ser Ser Glu Lys Asp Ile Lys Ser Thr Cys Ser Lys Phe Ser Leu
 5 10 15
 Ser Val Val Ala Ala Ile Leu Ala Ser Val Ser Gly Leu Ala Ser Cys
 20 25 30
 Val Asp Leu His Ala Gly Gly Gln Ser Val Asn Glu Leu Val Tyr Val
 35 40 45
 Gly Pro Gln Ala Val Leu Leu Leu Asp Gln Ile Arg Asp Leu Phe Val
 50 55 60
 Gly Ser Lys Asp Ser Gln Ala Glu Gly Gln Tyr Arg Leu Ile Val Gly
 65 70 75 80
 Asp Pro Ser Ser Phe Gln Glu Lys Asp Ala Asp Thr Leu Pro Gly Lys
 85 90 95
 Val Glu Gln Ser Thr Leu Phe Ser Val Thr Asn Pro Val Val Phe Gln
 100 105 110
 Gly Val Asp Gln Gln Asp Gln Val Ser Ser Gln Gly Leu Ile Cys Ser
 115 120 125
 Phe Thr Ser Ser Asn Leu Asp Ser Pro Arg Asp Gly Glu Ser Phe Leu
 130 135 140
 Gly Ile Ala Phe Val Gly Asp Ser Ser Lys Ala Gly Ile Thr Leu Thr
 145 150 155 160
 Asp Val Lys Ala Ser Leu Ser Gly Ala Ala Leu Tyr Ser Thr Glu Asp
 165 170 175

Leu Ile Phe Glu Lys Ile Lys Gly Gly Leu Glu Phe Ala Ser Cys Ser
 180 185 190
 Ser Leu Glu Gln Gly Gly Ala Cys Ala Ala Gln Ser Ile Leu Ile His
 195 200 205
 Asp Cys Gln Gly Leu Gln Val Lys His Cys Thr Thr Ala Val Asn Ala
 210 215 220
 Glu Gly Ser Ser Ala Asn Asp His Leu Gly Phe Gly Gly Ala Phe
 225 230 235 240
 Phe Val Thr Gly Ser Leu Ser Gly Glu Lys Ser Leu Tyr Met Pro Ala
 245 250 255
 Gly Asp Met Val Val Ala Asn Cys Asp Gly Ala Ile Ser Phe Glu Gly
 260 265 270
 Asn Ser Ala Asn Phe Ala Asn Gly Gly Ala Ile Ala Ala Ser Gly Lys
 275 280 285
 Val Leu Phe Val Ala Asn Asp Lys Lys Thr Ser Phe Ile Glu Asn Arg
 290 295 300
 Ala Leu Ser Gly Gly Ala Ile Ala Ala Ser Ser Asp Ile Ala Phe Gln
 305 310 315 320
 Asn Cys Ala Glu Leu Val Phe Lys Gly Asn Cys Ala Ile Gly Thr Glu
 325 330 335
 Asp Lys Gly Ser Leu Gly Gly Gly Ala Ile Ser Ser Leu Gly Thr Val
 340 345 350
 Leu Leu Gln Gly Asn His Gly Ile Thr Cys Asp Lys Asn Glu Ser Ala
 355 360 365
 Ser Gln Gly Gly Ala Ile Phe Gly Lys Asn Cys Gln Ile Ser Asp Asn
 370 375 380
 Glu Gly Pro Val Val Phe Arg Asp Ser Thr Ala Cys Leu Gly Gly Gly
 385 390 395 400
 Ala Ile Ala Ala Gln Glu Ile Val Ser Ile Gln Asn Asn Gln Ala Gly
 405 410 415
 Ile Ser Phe Glu Gly Gly Lys Ala Ser Phe Gly Gly Gly Ile Ala Cys
 420 425 430
 Gly Ser Phe Ser Ser Ala Gly Gly Ala Ser Val Leu Gly Thr Ile Asp
 435 440 445
 Ile Ser Lys Asn Leu Gly Ala Ile Ser Phe Ser Arg Thr Leu Cys Thr
 450 455 460
 Thr Ser Asp Leu Gly Gln Met Glu Tyr Gln Gly Gly Gly Ala Leu Phe
 465 470 475 480
 Gly Glu Asn Ile Ser Leu Ser Glu Asn Ala Gly Val Leu Thr Phe Lys
 485 490 495
 Asp Asn Ile Val Lys Thr Phe Ala Ser Asn Gly Lys Ile Leu Gly Gly
 500 505 510
 Gly Ala Ile Leu Ala Thr Gly Lys Val Glu Ile Thr Asn Asn Ser Glu
 515 520 525
 Gly Ile Ser Phe Thr Gly Asn Ala Arg Ala Pro Gln Ala Leu Pro Thr
 530 535 540
 Gln Glu Glu Phe Pro Leu Phe Ser Lys Lys Glu Gly Arg Pro Leu Ser
 545 550 555 560
 Ser Gly Tyr Ser Gly Gly Gly Ala Ile Leu Gly Arg Glu Val Ala Ile
 565 570 575
 Leu His Asn Ala Ala Val Val Phe Glu Gln Asn Arg Leu Gln Cys Ser
 580 585 590
 Glu Glu Glu Ala Thr Leu Leu Gly Cys Cys Gly Gly Ala Val His
 595 600 605
 Gly Met Asp Ser Thr Ser Ile Val Gly Asn Ser Ser Val Arg Phe Gly
 610 615 620
 Asn Asn Tyr Ala Met Gly Gln Gly Val Ser Gly Gly Ala Leu Leu Ser
 625 630 635 640
 Lys Thr Val Gln Leu Ala Gly Asn Gly Ser Val Asp Phe Ser Arg Asn
 645 650 655
 Ile Ala Ser Leu Gly Gly Gly Ala Leu Gln Ala Ser Glu Gly Asn Cys

660												665												670													
Gly	Leu	Val	Asp	Asn	Gly	Tyr	Val	Leu	Phe	Arg	Asp	Asn	Arg	Gly	Arg																						
675												680												685													
Val	Tyr	Gly	Gly	Gly	Ala	Ile	Ser	Cys	Leu	Arg	Gly	Asp	Val	Ile	Ser																						
690												695												700													
Gly	Asn	Gly	Gly	Gly	Arg	Val	Glu	Phe	Lys	Asp	Asn	Ile	Ala	Val	Leu	Arg																					
705												710												715													
Tyr	Val	Glu	Glu	Glu	Thr	Val	Glu	Lys	Val	Glu	Glu	Val	Glu	Pro	Ala	Pro																					
720												725												730													
Gly	Gln	Lys	Asp	Asn	Asn	Glu	Lys	Ser	Phe	Lys	Phe	Lys	Arg	Gly	Ala	Gly																					
735												740												745													
Ser	Phe	Ile	Thr	Ala	Ala	Asn	Gln	Ala	Leu	Phe	Ala	Lys	Arg	Val	Arg	Ala																					
750												755												760													
Asp	Leu	Ser	Pro	Glu	Ser	Ser	Ile	Ser	Ser																												

Ala Ser Ala Glu Ile Ser Asn Leu Ser Val Ser Asp Leu Gln Ile His
 1155 1160 1165
 Val Val Thr Pro Glu Ile Glu Glu Asp Thr Tyr Gly His Met Gly Asp
 1170 1175 1180
 Trp Ser Glu Ala Lys Ile Gln Asp Gly Thr Leu Val Ile Ser Trp Asn
 1185 1190 1195 1200
 Pro Thr Gly Tyr Arg Leu Asp Pro Gln Lys Ala Gly Ala Leu Val Phe
 1205 1210 1215
 Asn Ala Leu Trp Glu Glu Gly Ala Val Leu Ser Ala Leu Lys Asn Ala
 1220 1225 1230
 Arg Phe Ala His Asn Leu Thr Ala Gln Arg Met Glu Phe Asp Tyr Ser
 1235 1240 1245
 Thr Asn Val Trp Gly Phe Ala Phe Gly Gly Phe Arg Thr Leu Ser Ala
 1250 1255 1260
 Glu Asn Leu Val Ala Ile Asp Gly Tyr Lys Gly Ala Tyr Gly Gly Ala
 1265 1270 1275 1280
 Ser Ala Gly Val Asp Ile Gln Leu Met Glu Asp Phe Val Leu Gly Val
 1285 1290 1295
 Ser Gly Ala Ala Phe Leu Gly Lys Met Asp Ser Gln Lys Phe Asp Ala
 1300 1305 1310
 Glu Val Ser Arg Lys Gly Val Val Gly Ser Val Tyr Thr Gly Phe Leu
 1315 1320 1325
 Ala Gly Ser Trp Phe Phe Lys Gly Gln Tyr Ser Leu Gly Glu Thr Gln
 1330 1335 1340
 Asn Asp Met Lys Thr Arg Tyr Gly Val Leu Gly Glu Ser Ser Ala Ser
 1345 1350 1355 1360
 Trp Thr Ser Arg Gly Val Leu Ala Asp Ala Leu Val Glu Tyr Arg Ser
 1365 1370 1375
 Leu Val Gly Pro Val Arg Pro Thr Phe Tyr Ala Leu His Phe Asn Pro
 1380 1385 1390
 Tyr Val Glu Val Ser Tyr Ala Ser Met Lys Phe Pro Gly Phe Thr Glu
 1395 1400 1405
 Gln Gly Arg Glu Ala Arg Ser Phe Glu Asp Ala Ser Leu Thr Asn Ile
 1410 1415 1420
 Thr Ile Pro Leu Gly Met Lys Phe Glu Leu Ala Phe Ile Lys Gly Gln
 1425 1430 1435 1440
 Phe Ser Glu Val Asn Ser Leu Gly Ile Ser Tyr Ala Trp Glu Ala Tyr
 1445 1450 1455
 Arg Lys Val Glu Gly Gly Ala Val Gln Leu Leu Glu Ala Gly Phe Asp
 1460 1465 1470
 Trp Glu Gly Ala Pro Met Asp Leu Pro Arg Gln Glu Leu Arg Val Ala
 1475 1480 1485
 Leu Glu Asn Asn Thr Glu Trp Ser Ser Tyr Phe Ser Thr Val Leu Gly
 1490 1495 1500
 Leu Thr Ala Phe Cys Gly Gly Phe Thr Ser Thr Asp Ser Lys Leu Gly
 1505 1510 1515 1520
 Tyr Glu Ala Asn Thr Gly Leu Arg Leu Ile Phe
 1525 1530

<210> 99

<211> 474

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 99

Met Lys Ile Ile His Thr Ala Ile Glu Phe Ala Pro Val Ile Lys Ala
 5 10 15
 Gly Gly Leu Gly Asp Ala Leu Tyr Gly Leu Ala Lys Ala Leu Ala Ala
 20 25 30
 Asn His Thr Thr Glu Val Val Ile Pro Leu Tyr Pro Lys Leu Phe Thr

5 10 15
 Val Val Cys Gly Glu Glu Lys Glu Ile Ser Leu Ala Asp Phe Arg Gly
 20 25 30
 Lys Tyr Val Val Leu Phe Phe Tyr Pro Lys Asp Phe Thr Tyr Val Cys
 35 40 45
 Pro Thr Glu Leu His Ala Phe Gln Asp Arg Leu Val Asp Phe Glu Glu
 50 55 60
 Arg Gly Ala Val Val Leu Gly Cys Ser Val Asp Asp Ile Glu Thr His
 65 70 75 80
 Ser Arg Trp Leu Ala Val Ala Arg Asn Ala Gly Ile Glu Gly Thr
 85 90 95
 Glu Tyr Pro Leu Leu Ala Asp Pro Ser Phe Lys Ile Ser Glu Ala Phe
 100 105 110
 Gly Val Leu Asn Pro Glu Gly Ser Leu Ala Leu Arg Ala Thr Phe Leu
 115 120 125
 Ile Asp Lys Tyr Gly Val Val Arg His Ala Val Ile Asn Asp Leu Pre
 130 135 140
 Leu Gly Arg Ser Ile Asp Glu Glu Leu Arg Ile Leu Asp Ser Leu Ile
 145 150 155 160
 Phe Phe Glu Asn His Gly Met Val Cys Pro Ala Asn Trp Arg Ser Gly
 165 170 175
 Glu Arg Gly Met Val Pro Ser Glu Glu Gly Leu Lys Glu Tyr Phe Gln
 180 185 190
 Thr Met Asp
 195

<210> 102

<211> 86

<212> PRT

<213> *Chlamydia trachomatis* serovar D

<400> 102

Met Ser Gln Asn Lys Asn Ser Ala Phe Met Gln Pro Val Asn Val Ser
 5 10 15
 Ala Asp Leu Ala Ala Ile Val Gly Ala Gly Pro Met Pro Arg Thr Glu
 20 25 30
 Ile Ile Lys Lys Met Trp Asp Tyr Ile Lys Lys Asn Gly Leu Gln Asp
 35 40 45
 Pro Thr Asn Lys Arg Asn Ile Asn Pro Asp Asp Lys Leu Ala Lys Val
 50 55 60
 Phe Gly Thr Glu Lys Pro Ile Asp Met Phe Gln Met Thr Lys Met Val
 65 70 75 80
 Ser Gln His Ile Ile Lys
 85

<210> 103

<211> 394

<212> PRT

<213> *Chlamydia trachomatis* serovar D

<400> 103

Met Ser Lys Lys Glu Thr Phe Gln Arg Asn Lys Pro His Ile Asn Ile Gly
 5 10 15
 Thr Ile Gly His Val Asp His Gly Lys Thr Thr Leu Thr Ala Ala Ile
 20 25 30
 Thr Arg Ala Leu Ser Gly Asp Gly Leu Ala Asp Phe Arg Asp Tyr Ser
 35 40 45
 Ser Ile Asp Asn Thr Pro Glu Glu Lys Ala Arg Gly Ile Thr Ile Asn
 50 55 60

Ala Ser His Val Glu Tyr Glu Thr Ala Asn Arg His Tyr Ala His Val
 65 70 75 80
 Asp Cys Pro Gly His Ala Asp Tyr Val Lys Asn Met Ile Thr Gly Ala
 85 90 95
 Ala Gln Met Asp Gly Ala Ile Leu Val Val Ser Ala Thr Asp Gly Ala
 100 105 110
 Met Pro Gln Thr Lys Glu His Ile Leu Leu Ala Arg Gln Val Gly Val
 115 120 125
 Pro Tyr Ile Val Val Phe Leu Asn Lys Ile Asp Met Ile Ser Glu Glu
 130 135 140
 Asp Ala Glu Leu Val Asp Leu Val Glu Met Glu Leu Val Glu Leu Leu
 145 150 155 160
 Glu Glu Lys Gly Tyr Lys Gly Cys Pro Ile Ile Arg Gly Ser Ala Leu
 165 170 175
 Lys Ala Leu Glu Glu Gly Asp Ala Ala Tyr Ile Glu Lys Val Arg Glu Leu
 180 185 190
 Met Gln Ala Val Asp Asp Asn Ile Pro Thr Pro Glu Arg Glu Ile Asp
 195 200 205
 Lys Pro Phe Leu Met Pro Ile Glu Asp Val Phe Ser Ile Ser Gly Arg
 210 215 220
 Gly Thr Val Val Thr Gly Arg Ile Glu Arg Gly Ile Val Lys Val Ser
 225 230 235
 Asp Lys Val Gln Leu Val Gly Leu Arg Asp Thr Lys Glu Thr Ile Val
 245 250 255
 Thr Gly Val Glu Met Phe Arg Lys Glu Leu Pro Glu Gly Arg Ala Gly
 260 265 270
 Glu Asn Val Gly Leu Leu Leu Arg Gly Ile Gly Lys Asn Asp Val Glu
 275 280 285
 Arg Gly Met Val Val Cys Leu Pro Asn Ser Val Lys Pro His Thr Gln
 290 295 300
 Phe Lys Cys Ala Val Tyr Val Leu Gln Lys Glu Glu Gly Gly Arg His
 305 310 315
 Lys Pro Phe Thr Gly Tyr Arg Pro Gln Phe Phe Arg Thr Thr
 325 330 335
 Asp Val Thr Gly Val Val Thr Leu Pro Glu Gly Ile Glu Met Val Met
 340 345 350
 Pro Gly Asp Asn Val Glu Phe Glu Val Gln Leu Ile Ser Pro Val Ala
 355 360 365
 Leu Glu Glu Gly Met Arg Phe Ala Ile Arg Glu Gly Gly Arg Thr Ile
 370 375 380
 Gly Ala Gly Thr Ile Ser Lys Ile Ile Ala
 385 390

<210> 104

<211> 82

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 104

Met Gly Gln Asp His Arg Arg Lys Phe Leu Lys Lys Val Ser Phe Val
 5 10 15
 Lys Lys Gln Ala Ala Phe Ala Gly Asn Phe Ile Glu Glu Ile Lys Lys
 20 25 30
 Ile Glu Trp Val Asn Lys Arg Asp Leu Lys Arg Tyr Val Lys Ile Val
 35 40 45
 Leu Met Asn Ile Phe Gly Phe Gly Phe Ser Ile Tyr Cys Val Asp Leu
 50 55 60
 Ala Leu Arg Lys Ser Leu Ser Leu Phe Gly Lys Val Thr Ser Phe Phe
 65 70 75 80
 Phe Gly

<210> 105
 <211> 379
 <212> PRT
 <213> Chlamydia trachomatis serovar D

<400> 105
 Met Val Ile Pro Lys Val Asp Leu Gly Glu Ser Ala Val Met Met Gly
 5 10 15
 Tyr Lys Leu Thr Ser Gln Leu Ala Met Leu Ser Ile Leu Leu Thr Phe
 20 25 30
 Thr His Thr Met Gly His Ala Ser Gln Met Ser Gln Thr Leu Pro Thr
 35 40 45
 Ile Ile Glu Ala Gln Ala Glu Glu Ala Leu Gln Ala Asp Arg Gly Val
 50 55 60
 Ala Gly Gln Ala Leu Lys Lys Leu Arg Lys Lys Arg Cys Ala Ser Arg
 65 70 75 80
 Lys Ser Ala Cys Lys Ala Ser Phe Lys Lys Lys Asp Phe Phe Ser Cys
 85 90 95
 Ile Thr Asn Gly Leu Phe Ser Gly Asn His Glu Gln Arg Leu Thr Ala
 100 105 110
 Lys Lys Glu Asn Lys Ala Arg Gly Lys Glu Pro Arg Val Val Gln
 115 120 125
 Thr Thr Lys Lys Arg Gln Ile Thr Gln Ser Glu Lys Glu Phe Phe Asp
 130 135 140
 Trp Leu Cys Asn Ser Lys Arg Glu Arg Lys Leu Lys Lys Lys Pro
 145 150 155 160
 Val Asn Thr Ser Ser Leu Ala Lys Ser Glu Glu Leu Ser Pro Lys Glu Ala
 165 170 175
 Ala Ile Ala Ala Arg Ala Ser Leu Ser Pro Glu Glu Lys Arg Gln
 180 185 190
 Leu Ile Arg Glu Trp Leu Ala Glu Glu Lys Thr Ala Arg Lys Ser Gly
 195 200 205
 Arg Ala Ala Cys Ala Val Ser Glu Asn Leu Lys Arg Asp Gly Ser Ile
 210 215 220
 Thr Ser Thr Leu Arg Tyr Asp Ala Glu Lys Ala Leu Thr Thr Arg Val
 225 230 235 240
 Lys Arg Asn Glu Asn Ser Val Asn Ala Arg Ala Arg Gln Arg Ala Ala
 245 250 255
 Leu Gln Lys Ala Lys Lys Ala Lys Thr Glu Lys Pro Glu Ala Asp Glu
 260 265 270
 Lys Ala Ala Glu Ala Val Ala Ala Ala Pro Thr Lys Gln Ala His Lys
 275 280 285
 Glu Pro Glu Asn Tyr Phe Ala Ala Thr Ala Ser Thr Asn Asn Thr Asn
 290 295 300
 Val Met Ser Tyr Leu Asn Ala His Gln Tyr Arg Cys Asp Ser Ser Glu
 305 310 315 320
 Thr Asp Trp Pro Cys Ser Ser Cys Val Thr Lys Arg Arg Ala Asn Phe
 325 330 335
 Gly Ile Ser Val Cys Thr Met Val Val Thr Val Ile Ala Met Ile Val
 340 345 350
 Gly Ala Val Ile Ile Ser Asn Ala Thr Asp Ser Thr Val Ala Gly Ser
 355 360 365
 Ser Gly Thr Gly Gly Gly Gly Ser Thr Gln Pro
 370 375

<210> 106
 <211> 563

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 106

```

Met Val Tyr Phe Arg Ala His Gln Pro Arg His Thr Pro Lys Thr Phe
                    5                               10           15
Pro Leu Glu Val His His Ser Phe Ser Asp Lys His Pro Gln Ile Ala
                20                25                30
Lys Ala Met Arg Ile Thr Gly Ile Ala Leu Ala Ala Leu Ser Leu Leu
                35                40                45
Ala Val Val Ala Cys Val Ile Ala Val Ser Ala Gly Gly Ala Ala Ile
                50                55                60
Pro Leu Ala Val Ile Ser Gly Ile Ala Val Met Ser Gly Leu Leu Ser
                65                70                75                80
Ala Ala Thr Ile Ile Cys Ser Ala Lys Lys Ala Leu Ala Gln Arg Lys
                85                90                95
Gln Lys Gln Leu Glu Glu Ser Leu Pro Leu Asp Asn Ala Thr Glu His
                100               105               110
Val Ser Tyr Leu Thr Ser Asp Thr Ser Tyr Phe Asn Gln Trp Glu Ser
                115               120               125
Leu Gly Ala Leu Asn Lys Gln Leu Ser Gln Ile Asp Leu Thr Ile Gln
                130               135               140
Ala Pro Glu Lys Lys Leu Leu Lys Glu Val Leu Gly Ser Arg Tyr Asp
                145               150               155               160
Ser Ile Asn His Ser Ile Glu Glu Ile Ser Asp Arg Phe Thr Lys Met
                165               170               175
Leu Ser Leu Leu Arg Leu Arg Glu His Phe Tyr Arg Gly Glu Glu Arg
                180               185               190
Tyr Ala Pro Tyr Leu Ser Pro Pro Leu Leu Asn Lys Asn Arg Leu Leu
                195               200               205
Thr Gln Ile Thr Ser Asn Met Ile Arg Met Leu Pro Lys Ser Gly Gly
                210               215               220
Val Phe Ser Leu Lys Ala Asn Thr Leu Ser His Ala Ser Arg Thr Leu
                225               230               235               240
Tyr Thr Val Leu Lys Val Ala Leu Ser Leu Gly Val Leu Ala Gly Val
                245               250               255
Ala Ala Leu Ile Ile Phe Leu Pro Pro Ser Leu Pro Phe Ile Ala Val
                260               265               270
Ile Gly Val Ser Ser Leu Ala Leu Gly Met Ala Ser Phe Leu Met Ile
                275               280               285
Arg Gly Ile Lys Tyr Leu Leu Glu His Ser Pro Leu Asn Arg Lys Gln
                290               295               300
Leu Ala Lys Asp Ile Gln Lys Thr Ile Gly Pro Asp Val Leu Ala Ser
                305               310               315               320
Met Val His Tyr Gln His Gln Leu Leu Ser His Leu His Glu Thr Leu
                325               330               335
Leu Asp Glu Ala Ile Thr Ala Arg Trp Ser Glu Pro Phe Phe Ile Glu
                340               345               350
His Ala Asn Leu Lys Ala Lys Ile Glu Asp Leu Thr Lys Gln Tyr Asp
                355               360               365
Ile Leu Asn Ala Ala Phe Asn Lys Ser Leu Gln Gln Asp Glu Ala Leu
                370               375               380
Arg Ser Gln Leu Glu Lys Arg Ala Tyr Leu Phe Pro Ile Pro Asn Asn
                385               390               395               400
Asp Glu Asn Ala Lys Thr Lys Glu Ser Gln Leu Leu Asp Ser Glu Asn
                405               410               415
Asp Ser Asn Ser Glu Phe Gln Glu Ile Ile Asn Lys Gly Leu Glu Ala
                420               425               430
Ala Asn Lys Arg Arg Ala Asp Ala Lys Ser Lys Phe Tyr Thr Glu Asp
                435               440               445
Glu Thr Ser Asp Lys Ile Phe Ser Ile Trp Lys Pro Thr Lys Asn Leu

```

450
 Ala Leu Glu Asp Leu Trp Arg Val His Glu Ala Cys Asn Glu Glu Gln
 465
 Gln Ala Leu Leu Leu Glu Asp Tyr Met Ser Tyr Lys Thr Ser Glu Cys
 485
 Gln Ala Ala Leu Gln Lys Val Ser Gln Glu Leu Lys Ala Ala Gln Lys
 500
 Ser Phe Ala Val Leu Glu Lys His Ala Leu Asp Arg Ser Tyr Glu Ser
 515
 Ser Val Ala Thr Met Asp Leu Ala Arg Ala Asn Gln Glu Thr His Arg
 530
 Leu Leu Asn Ile Leu Ser Glu Leu Gln Gln Leu Ala Gln Tyr Leu Leu
 545
 Asp Asn His

<210> 107
 <211> 358
 <212> PRT
 <213> Chlamydia trachomatis serovar D

<400> 107
 Met Arg Lys Thr Val Ile Val Ala Met Ser Gly Gly Val Asp Ser Ser Ser
 5
 Val Val Ala Tyr Leu Leu Lys Lys Gln Gly Glu Tyr Asn Val Val Gly
 20
 Leu Phe Met Lys Asn Trp Gly Glu Gln Asp Glu Asn Gly Glu Cys Thr
 35
 Ala Thr Lys Asp Phe Arg Asp Val Glu Arg Ile Ala Glu Gln Leu Ser
 50
 Ile Pro Tyr Tyr Thr Val Ser Phe Ser Lys Glu Tyr Lys Glu Arg Val
 65
 Phe Ser Arg Phe Leu Arg Glu Tyr Ala Asn Gly Tyr Thr Pro Asn Pro
 85
 Asp Val Leu Cys Asn Arg Glu Ile Lys Phe Asp Leu Leu Gln Lys Lys
 100
 Val Arg Glu Leu Lys Gly Asp Phe Leu Ala Thr Gly His Tyr Cys Arg
 115
 Gly Gly Ala Asp Gly Thr Gly Leu Ser Arg Gly Ile Asp Pro Asn Lys
 130
 Asp Gln Ser Tyr Phe Leu Cys Gly Thr Pro Lys Asp Ala Leu Ser Asn
 145
 Val Leu Phe Pro Leu Gly Gly Met Tyr Lys Thr Glu Val Arg Arg Ile
 165
 Ala Gln Glu Ala Gly Leu Ala Thr Ala Thr Lys Lys Asp Ser Thr Gly
 180
 Ile Cys Phe Ile Gly Lys Arg Pro Phe Lys Ser Phe Leu Glu Gln Phe
 195
 Val Ala Asp Ser Pro Gly Asp Ile Ile Asp Phe Asp Thr Gln Gln Val
 210
 Val Gly Arg His Glu Gly Ala His Tyr Tyr Thr Ile Gly Gln Arg Arg
 225
 Gly Leu Asn Ile Gly Gly Met Glu Lys Pro Cys Tyr Val Leu Ser Lys
 245
 Asn Met Glu Lys Asn Ile Val Tyr Ile Val Arg Gly Glu Asp His Pro
 260
 Leu Leu Tyr Arg Gln Glu Leu Leu Ala Lys Glu Leu Asn Trp Phe Val
 275
 Pro Leu Gln Glu Pro Met Ile Cys Ser Ala Lys Val Arg Tyr Arg Ser
 290

77

Pro Asp Glu Lys Cys Ser Val Tyr Pro Leu Glu Asp Gly Thr Val Lys
 305 310 315 320
 Val Ile Phe Asp Val Pro Val Lys Ala Val Thr Pro Gly Gln Thr Val
 325 330 335
 Ala Phe Tyr Gln Gly Asp Ile Cys Leu Gly Gly Val Ile Glu Val
 340 345 350
 Pro Met Ile His Gln Leu
 355

<210> 108
 <211> 267
 <212> PRT
 <213> Chlamydia trachomatis serovar D

<400> 108
 Met Ser Arg Lys Pro Ala Ser Asn Ser Ser Arg Asn Thr Lys Arg Ser
 5 10 15
 Ser Asp Thr Ser Trp Glu Val Ile Ala Gln Asp Tyr Asn Lys Ala Val
 20 25 30
 Asp Arg Asp Gly His Phe Tyr His Lys Glu Val Ile Leu Pro Asn Leu
 35 40 45
 Leu Ser Lys Leu His Ile Ser Arg Ser Ser Ser Leu Val Asp Val Gly
 50 55 60
 Cys Gly Gln Gly Ile Leu Glu Lys His Leu Pro Lys His Leu Pro Tyr
 65 70 75 80
 Leu Gly Ile Asp Leu Ser Pro Ser Leu Leu Arg Phe Ala Lys Lys Ser
 85 90 95
 Ala Ser Ser Lys Ser Arg Arg Phe Leu His His Asp Met Thr Gln Pro
 100 105 110
 Val Pro Ala Asp His His Glu Gln Phe Ser His Ala Thr Ala Ile Leu
 115 120 125
 Ser Leu Gln Asn Met Glu Ser Pro Glu Gln Ala Ile Ala His Thr Ala
 130 135 140
 Asn Leu Leu Ala Pro Gln Gly Arg Leu Phe Ile Val Leu Asn His Pro
 145 150 155 160
 Cys Phe Arg Ile Pro Arg Leu Ser Ser Trp Leu Tyr Asp Glu Pro Lys
 165 170 175
 Lys Leu Leu Ser Arg Lys Ile Asp Arg Tyr Leu Ser Pro Val Ala Val
 180 185 190
 Pro Ile Val Val His Pro Gly Glu Lys His Ser Glu Thr Thr Tyr Ser
 195 200 205
 Phe His Phe Pro Leu Ser Tyr Trp Val Gln Ala Leu Ser Asn His Asn
 210 215 220
 Leu Leu Ile Asp Ser Met Glu Glu Trp Ile Ser Pro Lys Lys Ser Ser
 225 230 235 240
 Gly Lys Arg Ala Arg Ala Glu Asn Leu Cys Arg Lys Glu Phe Pro Leu
 245 250 255
 Phe Leu Phe Ile Ser Ala Leu Lys Ile Ser Lys
 260 265

<210> 109
 <211> 867
 <212> PRT
 <213> Chlamydia trachomatis serovar D

<400> 109
 Met Glu Lys Phe Ser Asp Ala Val Ser Glu Ala Leu Glu Lys Ala Phe
 5 10 15

Glu Leu Ala Lys Asn Ser Lys His Ser Tyr Val Thr Glu Asn His Leu
 20 25 30
 Leu Lys Ser Leu Leu Gln Asn Pro Gly Ser Leu Phe Cys Leu Val Ile
 35 40 45
 Lys Asp Val His Gly Asn Leu Gly Leu Thr Ser Ala Val Asp Asp
 50 55 60
 Ala Leu Arg Arg Glu Pro Thr Val Val Glu Gly Thr Ala Val Ala Ser
 65 70 75 80
 Pro Ser Pro Ser Leu Gln Gln Leu Leu Leu Asn Ala His Gln Glu Ala
 85 90 95
 Arg Ser Met Gly Asp Glu Tyr Leu Ser Gly Asp His Leu Leu Leu Ala
 100 105 110
 Phe Trp Arg Ser Thr Lys Glu Pro Phe Ala Ser Trp Arg Lys Thr Val
 115 120 125
 Lys Thr Thr Ser Glu Ala Leu Lys Glu Leu Ile Thr Lys Leu Arg Gln
 130 135 140
 Gly Ser Arg Met Asp Ser Pro Ser Ala Glu Glu Asn Leu Lys Gly Leu
 145 150 155 160
 Glu Lys Tyr Cys Lys Asn Leu Thr Val Leu Ala Arg Glu Gly Lys Leu
 165 170 175
 Asp Pro Val Ile Gly Arg Asp Glu Glu Ile Arg Arg Thr Ile Gln Val
 180 185 190
 Leu Ser Arg Arg Thr Lys Asn Asn Pro Met Leu Ile Gly Glu Pro Gly
 195 200 205
 Val Gly Lys Thr Ala Ile Ala Glu Gly Leu Ala Leu Arg Ile Val Gln
 210 215 220
 Gly Asp Val Pro Glu Ser Leu Lys Glu Lys His Leu Tyr Val Leu Asp
 225 230 235 240
 Met Gly Ala Leu Ile Ala Gly Ala Lys Tyr Arg Gly Glu Phe Glu Glu
 245 250 255
 Arg Leu Lys Ser Val Leu Lys Gly Val Glu Ala Ser Glu Gly Glu Cys
 260 265 270
 Ile Leu Phe Ile Asp Glu Val His Thr Leu Val Gly Ala Gly Ala Thr
 275 280 285
 Asp Gly Ala Met Asp Ala Ala Asn Leu Leu Lys Pro Ala Leu Ala Arg
 290 295 300
 Gly Thr Leu His Cys Ile Gly Ala Thr Thr Leu Asn Glu Tyr Gln Lys
 305 310 315 320
 Tyr Ile Glu Lys Asp Ala Ala Leu Glu Arg Arg Phe Gln Pro Ile Phe
 325 330 335
 Val Thr Glu Pro Ser Leu Glu Asp Ala Val Phe Ile Leu Arg Gly Leu
 340 345 350
 Arg Glu Lys Tyr Glu Ile Phe His Gly Val Arg Ile Thr Glu Gly Ala
 355 360 365
 Leu Asn Ala Ala Val Val Leu Ser Tyr Arg Tyr Ile Thr Asp Arg Phe
 370 375 380
 Leu Pro Asp Lys Ala Ile Asp Leu Ile Asp Glu Ala Ala Ser Leu Ile
 385 390 395 400
 Arg Met Gln Ile Gly Ser Leu Pro Leu Pro Ile Asp Glu Lys Glu Arg
 405 410 415
 Glu Leu Ser Ala Leu Ile Val Lys Gln Glu Ala Ile Lys Arg Glu Gln
 420 425 430
 Ala Pro Ala Tyr Gln Glu Glu Ala Glu Asp Met Gln Lys Ala Ile Asp
 435 440 445
 Arg Val Lys Glu Glu Leu Ala Ala Leu Arg Leu Arg Trp Asp Glu Glu
 450 455 460
 Lys Gly Leu Ile Thr Gly Leu Lys Glu Lys Lys Asn Ala Leu Glu Asn
 465 470 475 480
 Leu Lys Phe Ala Glu Glu Glu Ala Glu Arg Thr Ala Asp Tyr Asn Arg
 485 490 495
 Val Ala Glu Leu Arg Tyr Ser Leu Ile Pro Ser Leu Glu Glu Glu Ile

500 505 510
 His Leu Ala Glu Glu Ala Leu Asn Gln Arg Asp Gly Arg Leu Leu Gln
 515 520 525
 Glu Glu Val Asp Glu Arg Leu Ile Ala Gln Val Val Ala Asn Trp Thr
 530 535 540
 Gly Ile Pro Val Gln Lys Met Leu Glu Gly Glu Ser Glu Lys Leu Leu
 545 550 555 560
 Val Leu Glu Glu Ser Leu Glu Glu Arg Val Val Gly Gln Pro Phe Ala
 565 570 575
 Ile Ala Ala Val Ser Asp Ser Ile Arg Ala Ala Arg Val Gly Leu Ser
 580 585 590
 Asp Pro Gln Arg Pro Leu Gly Val Phe Leu Phe Leu Gly Pro Thr Gly
 595 600 605
 Val Gly Lys Thr Glu Leu Ala Lys Ala Leu Ala Glu Leu Leu Phe Asn
 610 615 620
 Lys Glu Glu Ala Met Ile Arg Phe Asp Met Thr Glu Tyr Met Glu Lys
 625 630 635 640
 His Ser Val Ser Lys Leu Ile Gly Ser Pro Gly Tyr Val Gly Tyr
 645 650 655
 Glu Glu Gly Gly Ser Leu Ser Glu Ala Leu Arg Arg Arg Pro Tyr Ser
 660 665 670
 Val Val Leu Phe Asp Glu Ile Glu Lys Ala Asp Lys Glu Val Phe Asn
 675 680 685
 Ile Leu Leu Gln Ile Phe Asp Asp Gly Ile Leu Thr Asp Ser Lys Lys
 690 695 700
 Arg Lys Val Asn Cys Lys Asn Ala Leu Phe Ile Met Thr Ser Asn Ile
 705 710 715 720
 Gly Ser Gln Glu Leu Ala Asp Tyr Cys Thr Lys Lys Gly Thr Ile Val
 725 730 735
 Asp Lys Glu Ala Val Leu Ser Val Val Ala Pro Ala Leu Lys Asn Tyr
 740 745 750
 Phe Ser Pro Glu Phe Ile Asn Arg Ile Asp Asp Ile Leu Pro Phe Val
 755 760 765
 Pro Leu Thr Thr Glu Asp Ile Val Lys Ile Val Gly Ile Gln Met Asn
 770 775 780
 Arg Val Ala Leu Arg Leu Leu Glu Arg Lys Ile Ser Leu Thr Trp Asp
 785 790 795 800
 Asp Ser Leu Val Leu Phe Leu Ser Glu Gln Gly Tyr Asp Ser Ala Phe
 805 810 815
 Gly Ala Arg Pro Leu Lys Arg Leu Ile Gln Gln Lys Val Thr Met
 820 825 830
 Leu Ser Lys Ala Leu Leu Lys Gly Asp Ile Lys Pro Gly Met Ala Val
 835 840 845
 Glu Leu Thr Met Ala Lys Asp Val Val Phe Lys Ile Lys Thr Asn
 850 855 860
 Pro Ala Val
 865

<210> 110

<211> 1170

<212> DNA

<213> Chlamydia pneumoniae

<400> 110

atgaaaaaac tcttaaagtc ggcgttatta tcgcgcgcac ttgctgggttc tgttggctcc 60
 ttacaagcct tgcctgtagg gaacccttct gatccaaagt tattaattga tggtaacaata 120
 tgggaagggtg ctgcaggaga tccttgcgat ccttgcgcga cttggtgcga cgtattagc 180
 ttacgtgctg gatattacgg agactatggt ttgcaccgta tcttaaaagt agatgcacct 240
 aaaaacatttt ctatggggagc caagcctact ggatccgctg ctgcaaacata tactactgcc 300
 gtagatagac ctaaccggcg ctacaataag catttacacg atgcagagtg gttcactaat 360

gcaggcttca	tgccttaaa	catttgggat	gcctttgatg	ttttctgtac	tttagagagt	420
tctaagtggt	acatttagag	aaactctaca	gcgttcaaat	tcgtttggtt	attcogaggt	480
aaaggtacta	ctgtaaatgc	aaatgaacta	ccaaacgttt	ctttaagtaa	cgaggttggt	540
gaactttaca	cacagacctc	tttctcttgg	agcgtagcgg	ctcgtggagc	cttatgggaa	600
tgccggttgg	caactttggg	agctgaattc	caatatgcac	agtccaaacc	taaagttgaa	660
gaacttaatg	tgtatgttaa	cgatcgcgaa	ttctctgtaa	acaaaccgaa	gggctataaa	720
ggcgttgatg	tccctctgcc	aacagacgct	ggcgttagcaa	cagctactgg	acaaagtctc	780
gcgaccatca	attatcatga	atggcgaagta	ggagcctctc	tatcttacag	actaaactct	840
ttagtgcacat	acaattggagt	acaatggtct	cgagcaactt	ttgatgctga	taacatccgc	900
attgtctcagc	caaaactacc	tacagctggt	ttaaacttaa	ctgcatggaa	ccctcttcta	960
ctaggaaatg	ccacagcaatt	gtctactact	gattcgtttc	cagactctcat	gcaaatgttt	1020
tcctgtcaga	tcacaagtt	taaactctaga	aaagcttgtg	gagttactgt	aggagctact	1080
ttagttgatg	ctgataaagt	gtcaacttact	gcagaagctc	gtttaattaa	cgagagagct	1140
gctocagtat	ctgtgcagtt	cgattcttaa				1170

<210> 111

<211> 2601

<212> DNA

<213> *Chlamydia pneumoniae*

<400> 111

atggagaaat	tttcogatgc	tgtctctgaa	gcttttagaga	aggcttttoga	acttgctataa	60
tcttcgaaac	atacctatgt	cacagaaaaa	acactattac	tggttttatt	agaaaataca	120
gagctctctc	ttatttttgt	aattaaggac	attcatggga	acccctggtt	gctcaatcag	180
cgacttaaaag	atcgctctctc	acgagagcgg	actgtagtgt	agggagaggt	ggatcctaaa	240
cctctctcgg	gtttcaaaac	ctctctctagg	gatgcacaaac	agaggcgaaa	gacattagga	300
gtgaataaca	ttctctggaga	ctactctgcty	cttgccttttt	ggagttcaaa	caaaagagcct	360
tttaattctct	gggaagcaac	aacaaaagt	agtttttaag	actcttaagaa	tcgtgattac	420
aaaatacagc	ggagaaatcg	tatgattgtc	ccagcggcty	aaagtaattt	tcagggtttta	480
gaaaagtatt	gtaaaaattt	aaacagcatc	gctctggaag	gttaactcgg	ctctgtgctc	540
ggtagagatg	aagaaattcg	tagaacatcc	caagtgcctt	cccttagaag	taaaaaaac	600
ccatgcttta	ttggtgagcc	gggtgtaggg	aaaactgcta	tagcagaagg	attagctctt	660
agggttatcc	agggtgatgt	tcttgaatct	ctcaaaagta	aacagcttta	tgtcttagat	720
atggagagctt	tgattgcagg	agctaagtat	cgaggtgagt	ttgaagaaag	actaaagagt	780
gttttaaaag	atgtgaatc	tggagatggc	gagcaacatta	cttttatgtg	tgaagtcac	840
acotctgttg	gagcaggagc	tactgatgga	gctatggatg	ctgcgaatct	tttaaagcct	900
gcatttagcaa	gagggacgct	acactgtatt	ggcgcgcaga	ctttgaatga	gtatcagaag	960
tatatgtaaa	aagatgctgc	tttggaaagt	cgatttccgc	ctatttttgt	gacagagcct	1020
tcttttgagg	atcgtctgtt	tattctctgt	ggactaaag	aaaaaatatga	aattttccat	1080
ggagtcagga	ttacagaggg	ggctttgaa	gcgcagctcc	tactttctca	tgcttatctc	1140
ccagatcgct	ttcttccaga	taaggctatc	gatttgatag	atgaagcggc	aaagttaatt	1200
cgactgcaaa	ttgttgatct	tctcttctc	attgatgaaa	aggagagaga	gctgtgcgt	1260
ttgatcgtta	agcaagaggc	tataaaacgc	gagcaatctc	cttctctatca	agaaggggct	1320
gatgctatgc	agaagcttat	agatgctttg	agagaggaat	tagcatctct	acgtttgggt	1380
ttgggtgaag	agaagaagtt	gatttccggg	ctcaaggaaa	aaaagaattc	cttggaaagt	1440
atgaaaattt	ctcaagagga	ggcggagcgt	gttgacagct	ataactgtgt	acgtgagcct	1500
cgttatagtt	taattcccca	acttgaagaa	gaatcaaac	aggatgaagc	ctctttaaat	1560
caaaagagata	acogtctctc	tcagaagaaa	gttgacgagc	gattgatggc	gcaagtggtta	1620
gcataattga	caaggattcc	gttgcaaaaa	atgtcagaag	gggaagcgtg	gaaactctga	1680
attcttgaag	aaactctaga	agaaactgtg	gtaggacagc	cttttgcagt	ctctgoggtt	1740
agtgatctta	ttcgtgctgc	acgtgtaggt	ttaaatgato	ctcaacgtcc	cttaggaagt	1800
tttttatctt	ttagggccaac	aggggttagga	aaaaaccgag	ttgcaaaagc	ctttgcagtc	1860
ctctcttttca	ataaagagga	agctatggct	cgcttcgata	tgctcagagta	tatggaaaag	1920
attcattctt	ccaaagcttat	aggatcttct	ccagggatg	tggtttatga	ggaaggttgg	1980
agctctttctg	aggtcctctg	acgacgtccc	tattcagtag	ttctctttga	tgatgatag	2040
aaagcagata	agaagttct	aaatatctct	ttcacaggtt	ttgatgatgg	gattctaacg	2100
gtggggaaaa	aacgcaaggt	aaattgttaa	aatgccttgt	ttatcatgac	atcaaatata	2160
ggctctccag	aacttgcaga	ttattgttca	aaaaaaggaa	gtgagcttac	gaaagaaagc	2220
attcttctctg	tagtctctcc	agtattgaaa	agatacttga	gcctcgaatt	tatgaaccca	2280
attgatgaga	tactctctct	tgttccatta	acgaagaag	atatogtgaa	aatagttgac	2340
attcaaatgc	gaaggattgc	ccagagatta	aaggcaagcg	ggatcaattt	atcttgggag	2400

```

gattctgtaa tattatttct tagtgaacag ggttatgaca gtgctttcgg agccgcgcct 2460
ttaaaccgtt tgatccaaca aaaagttgtg atcttgcttt ctaaggcttt gcttaaagga 2520
gatattaac cgtatacatc gattgagittg acgatggcaa aagaggtgct cgtattttaaa 2580
aaagtggaaa ctccctotta g                                2601

```

<210> 112

<211> 389

<212> PRT

<213> Chlamydia pneumoniae

<400> 112

```

Met Lys Lys Leu Leu Lys Ser Ala Leu Leu Ser Ala Ala Phe Ala Gly
      5      10      15
Ser Val Gly Ser Leu Gln Ala Leu Pro Val Gly Asn Pro Ser Asp Pro
      20      25      30
Ser Leu Leu Ile Asp Gly Thr Ile Trp Glu Gly Ala Ala Gly Asp Pro
      35      40      45
Cys Asp Pro Cys Ala Thr Trp Cys Asp Ala Ile Ser Leu Arg Ala Gly
      50      55      60
Phe Tyr Gly Asp Tyr Val Phe Asp Arg Ile Leu Lys Val Asp Ala Pro
      65      70      75
Lys Thr Phe Ser Met Gly Ala Lys Pro Thr Gly Ser Ala Ala Ala Asn
      85      90      95
Tyr Thr Thr Ala Val Asp Arg Pro Asn Pro Ala Tyr Asn Lys His Leu
      100      105      110
His Asp Ala Glu Trp Phe Thr Asn Ala Gly Phe Ile Ala Leu Asn Ile
      115      120      125
Trp Asp Arg Phe Asp Val Phe Cys Thr Leu Gly Ala Ser Asn Gly Tyr
      130      135      140
Ile Arg Gly Asn Ser Thr Ala Phe Asn Leu Val Gly Leu Phe Gly Val
      145      150      155
Lys Gly Thr Thr Val Asn Ala Asn Glu Leu Pro Asn Val Ser Leu Ser
      165      170      175
Asn Gly Val Val Glu Leu Tyr Thr Asp Thr Ser Phe Ser Trp Ser Val
      180      185      190
Gly Ala Arg Gly Ala Leu Trp Glu Cys Gly Cys Ala Thr Leu Gly Ala
      195      200      205
Glu Phe Gln Tyr Ala Gln Ser Lys Pro Lys Val Glu Glu Leu Asn Val
      210      215      220
Ile Cys Asn Val Ser Gln Phe Ser Val Asn Lys Pro Lys Gly Tyr Lys
      225      230      235
Gly Val Ala Phe Pro Leu Pro Thr Asp Ala Gly Val Ala Thr Ala Thr
      245      250      255
Gly Thr Lys Ser Ala Thr Ile Asn Tyr His Glu Trp Gln Val Gly Ala
      260      265      270
Ser Leu Ser Tyr Arg Leu Asn Ser Leu Val Pro Tyr Ile Gly Val Gln
      275      280      285
Trp Ser Arg Ala Thr Phe Asp Ala Asp Asn Ile Arg Ile Ala Gln Pro
      290      295      300
Lys Leu Pro Thr Ala Val Leu Asn Leu Thr Ala Trp Asn Pro Ser Leu
      305      310      315
Leu Gly Asn Ala Thr Ala Leu Ser Thr Thr Asp Ser Phe Ser Asp Phe
      325      330      335
Met Gln Ile Val Ser Cys Gln Ile Asn Lys Phe Lys Ser Arg Lys Ala
      340      345      350
Cys Gly Val Thr Val Gly Ala Thr Leu Val Asp Ala Asp Lys Trp Ser
      355      360      365
Leu Thr Ala Glu Ala Arg Leu Ile Asn Glu Arg Ala Ala His Val Ser
      370      375      380
Gly Gln Phe Arg Phe

```

385

<210> 113

<211> 866

<212> PRT

<213> Chlamydia pneumoniae

<400> 113

Met Glu Lys Phe Ser Asp Ala Val Ser Glu Ala Leu Glu Lys Ala Phe
5 10 15
Glu Leu Ala Lys Ser Ser Lys His Thr Tyr Val Thr Glu Asn His Leu
20 25 30
Leu Leu Ala Leu Leu Glu Asn Thr Glu Ser Leu Phe Tyr Leu Val Ile
35 40 45
Lys Asp Ile His Gly Asn Pro Gly Leu Leu Asn Thr Ala Val Lys Asp
50 55 60
Ala Leu Ser Arg Glu Pro Thr Val Val Glu Gly Glu Val Asp Pro Lys
65 70 75 80
Pro Ser Pro Gly Leu Gln Thr Leu Leu Arg Asp Ala Lys Gln Glu Ala
85 90 95
Lys Thr Leu Gly Asp Glu Tyr Ile Ser Gly Asp His Leu Leu Leu Ala
100 105 110
Phe Trp Ser Ser Asn Lys Glu Pro Phe Asn Ser Trp Lys Gln Thr Thr
115 120 125
Lys Val Ser Phe Lys Asp Leu Lys Asn Leu Ile Thr Lys Ile Arg Arg
130 135 140
Gly Asn Arg Met Asp Ser Pro Ser Ala Glu Ser Asn Phe Gln Gly Leu
145 150 155 160
Glu Lys Tyr Cys Lys Asn Leu Thr Ala Leu Ala Arg Glu Gly Lys Leu
165 170 175
Asp Pro Val Ile Gly Arg Asp Glu Glu Ile Arg Arg Thr Ile Gln Val
180 185 190
Leu Ser Arg Arg Thr Lys Asn Asn Pro Met Leu Ile Gly Glu Pro Gly
195 200 205
Val Gly Lys Thr Ala Ile Ala Glu Gly Leu Ala Leu Arg Leu Ile Gln
210 215 220
Gly Asp Val Pro Glu Ser Leu Lys Gly Lys Gln Leu Tyr Val Leu Asp
225 230 235 240
Met Gly Ala Leu Ile Ala Gly Ala Lys Tyr Arg Gly Glu Phe Glu Glu
245 250 255
Arg Leu Lys Ser Val Leu Lys Asp Val Glu Ser Gly Asp Gly Glu His
260 265 270
Ile Ile Phe Ile Asp Glu Val His Thr Leu Val Gly Ala Gly Ala Thr
275 280 285
Asp Gly Ala Met Asp Ala Ala Asn Leu Leu Lys Pro Ala Leu Ala Arg
290 295 300
Gly Thr Leu His Cys Ile Gly Ala Thr Thr Leu Asn Glu Tyr Gln Lys
305 310 315 320
Tyr Ile Glu Lys Asp Ala Ala Leu Glu Arg Arg Phe Gln Pro Ile Phe
325 330 335
Val Thr Glu Pro Ser Leu Glu Asp Ala Val Phe Ile Leu Arg Gly Leu
340 345 350
Arg Glu Lys Tyr Glu Ile Phe His Gly Val Arg Ile Thr Glu Gly Ala
355 360 365
Leu Asn Ala Ala Val Leu Leu Ser Tyr Arg Tyr Ile Pro Asp Arg Phe
370 375 380
Leu Pro Asp Lys Ala Ile Asp Leu Ile Asp Glu Ala Ala Ser Leu Ile
385 390 395 400
Arg Met Gln Ile Gly Ser Leu Pro Leu Pro Ile Asp Glu Lys Glu Arg
405 410 415

Glu Leu Ala Ala Leu Ile Val Lys Gln Glu Ala Ile Lys Arg Glu Gln
 420 425 430
 Ser Pro Ser Tyr Gln Glu Glu Ala Asp Ala Met Gln Lys Ser Ile Asp
 435 440 445
 Ala Leu Arg Glu Glu Leu Ala Ser Leu Arg Leu Gly Trp Asp Glu Glu
 450 455 460
 Lys Lys Leu Ile Ser Gly Leu Lys Glu Lys Lys Asn Ser Leu Glu Ser
 465 470 475 480
 Met Lys Phe Ser Glu Glu Glu Ala Glu Arg Val Ala Asp Tyr Asn Arg
 485 490 495
 Val Ala Glu Leu Arg Tyr Ser Leu Ile Pro Gln Leu Glu Glu Glu Ile
 500 505 510
 Lys Gln Asp Glu Ala Ser Leu Asn Gln Arg Asp Asn Arg Leu Leu Gln
 515 520 525
 Glu Glu Val Asp Glu Arg Leu Ile Ala Gln Val Val Ala Asn Trp Thr
 530 535 540
 Gly Ile Pro Val Gln Lys Met Leu Glu Gly Glu Ala Glu Lys Leu Leu
 545 550 555 560
 Ile Leu Glu Glu Ser Leu Glu Glu Arg Val Val Gly Gln Pro Phe Ala
 565 570 575
 Val Ser Ala Val Ser Asp Ser Ile Arg Ala Ala Arg Val Gly Leu Asn
 580 585 590
 Asp Pro Gln Arg Pro Leu Gly Val Phe Leu Phe Leu Gly Pro Thr Gly
 595 600 605
 Val Gly Lys Thr Glu Leu Ala Lys Ala Leu Ala Asp Leu Leu Phe Asn
 610 615 620
 Lys Glu Glu Ala Met Val Arg Phe Asp Met Ser Glu Tyr Met Glu Lys
 625 630 635 640
 His Ser Ile Ser Lys Leu Ile Gly Ser Ser Pro Gly Tyr Val Gly Tyr
 645 650 655
 Glu Glu Gly Gly Ser Leu Ser Glu Ala Leu Arg Arg Arg Pro Tyr Ser
 660 665 670
 Val Val Leu Phe Asp Glu Ile Glu Lys Ala Asp Lys Glu Val Leu Asn
 675 680 685
 Ile Leu Leu Gln Val Phe Asp Asp Gly Ile Leu Thr Asp Gly Lys Lys
 690 695 700
 Arg Lys Val Asn Cys Lys Asn Ala Leu Phe Ile Met Thr Ser Asn Ile
 705 710 715 720
 Gly Ser Pro Glu Leu Ala Asp Tyr Cys Ser Lys Lys Gly Ser Glu Leu
 725 730 735
 Thr Lys Glu Ala Ile Leu Ser Val Val Ser Pro Val Leu Lys Arg Tyr
 740 745 750
 Leu Ser Pro Glu Phe Met Asn Arg Ile Asp Glu Ile Leu Pro Phe Val
 755 760 765
 Pro Leu Thr Lys Glu Asp Ile Val Lys Ile Val Gly Ile Gln Met Arg
 770 775 780
 Arg Ile Ala Gln Arg Leu Lys Ala Arg Arg Ile Asn Leu Ser Trp Asp
 785 790 795 800
 Asp Ser Val Ile Leu Phe Leu Ser Glu Gln Gly Tyr Asp Ser Ala Phe
 805 810 815
 Gly Ala Arg Pro Leu Lys Arg Leu Ile Gln Gln Lys Val Val Ile Leu
 820 825 830
 Leu Ser Lys Ala Leu Leu Lys Gly Asp Ile Lys Pro Asp Thr Ser Ile
 835 840 845
 Glu Leu Thr Met Ala Lys Glu Val Leu Val Phe Lys Lys Val Glu Thr
 850 855 860
 Pro Ser
 865

<211> 1179
 <212> DNA
 <213> Homo sapiens

<400> 114
 taactctccc ctctctctctt aaaaaagagg ggagcctttt ttccctacaa agatcgccta 60
 gcttttctct gaagaatctc atcaagagat atttgcattt tccacggat aaaggcatcc 120
 caagggaagc ctggaatcac ttcatattct cccgttgcta gcattcgaca agggaaacca 180
 aagattaaat cttccogtaa tccataggga ttgtggtccg aacacactcc ggaagaaaaa 240
 catctctcct cttttggctg atattattgat cgaagcgcct ctgctaagag tcgtgtcgca 300
 gaagctgcgc aagacttccc tcgtgcttcg attactgcac taccacgact ctgtacagaa 360
 ggaccataaa tattctctaa ccaatcacga tccgctatcg tctctgcgat aggcgcgcta 420
 ttaatcagag ctctgcgtaaa atcaggcact ttgtttggcg agtgatttcc ccaaaccaca 480
 acttgcgtata cagccgataaa aggtacttct gctctatgag ataacatgct atgcatacga 540
 ttctggtcca atcgtagcat cgcatagaag ttcttttcca ataatctggg agcatgattc 600
 attgctatcc agcaattggt attcacaggg ttcccaaaaa caaaattctt tgcattcccg 660
 ttggctgttg ttgtcaaaag ttttctctgc gtacgaaaaa tctcccatct ttctttttaga 720
 agatcccttc ttctcattcc tgggcctcta ggaactgacc ctataaggaa tgcgcgcatca 780
 atgcacataa aagcatcatg caatgatgtc gttactcgca cagctgttaa taaagggaag 840
 gcaccatcat ctatgctccat ggcacaccca gataaagccc ttctgttctc aggaatatcg 900
 tagatacgca gatcgatgcc acaatcaagg ccaaaaaaat ctccatgagc cagagaaaaa 960
 agaaagctat aggcctattg cctctgttctc cctgttactg ctacactaac tgtttgagaa 1020
 accataaagc accctctctt taccttttaca aaacgcacat actctcaca ctacgtttgc 1080
 aactaacaaa ttttggctcc aacatacgtt tggatgataa aagaatcaag tacctagatt 1140
 ccttagtaaa agcttttggc aaaaaaaagg tcatctatt 1179

<210> 115
 <211> 772
 <212> DNA
 <213> Homo sapiens

<400> 115
 gcaaaactgc tgacaaagct ggagacggaa ctacaacagc tactgtttct gctgaagcta 60
 ttatatacaga aggattacgc aatgtaacag ctggagcaaa tccaatggac ctcaaacgag 120
 gtattgataa agctgttaag gttgtgtgtg atcaaatcag aaaaatcagc aaacctgttc 180
 agcatcatata agaaattgct caagttgcgaa caattttctg taataatgat gcagaaatcg 240
 ggaatctgat tgctgaagca atggagaaaag ttggtaaaaa cggctctatc actgttgaag 300
 aagcaaaaag atttgaiaacc gttttggatg ttgttgaagg aatgaatttc aatagaggtt 360
 accctctctag ctactctcgca acaaatccag aaactcaaga atgtgtatta ggaagcgtct 420
 ttgttctaat ctacgataag aaaatttctg ggatcaaaag tttccttctc gttttacaac 480
 aagttgctga atccggccgt cotcttctta ttatagcaga agacattgaa ggcgaagctt 540
 taqctacttt ggtcgtgaac agaattcgtg gaggattcgc ggtttgcgca gttaaaagcc 600
 cagcgttttg agatagaaga aaagctatgt tggaaagacat cgtatcttca actggcgttc 660
 aactcattag cgaagagttg ggcataaagt tagaaaagcg taacttagct atgttaggta 720
 aagctaaaaa agttatcgtt tctaaggaag acacgaccat cgtcgaagga at 772

<210> 116
 <211> 487
 <212> DNA
 <213> Homo sapiens

<400> 116
 gcagctcctgc caaagccaca agctcctgtc gcacaaacac ggcattttta aaagagccat 60
 cagattttctc ctctaatttt taacgcagtc tcccaacagc tgaataaacc tgaggaagaa 120
 agacgtcctct tggagctcgc atacttcaaa ggcgcggcta agcaggcagc tgctgcaaa 180
 gaaaaaaagg cttctgaaca ggaagtatcc aaaaagaaga aagaagcttc taactctgtg 240
 ggaagagaac agagttatgc tcgtcgtgct gtgaatgcca tcaatttcag tgtaagaaga 300
 caaatagaag agcaacagaa aaccatttcc aatccaggaa atgcacagac tcttctcgtg 360
 aagaagaatc cacatacatc cggagaacct gttatccaaa cgttacaaga ctgttctcag 420

85

gatcaagaag aagagaaaa agttctagag cgattaaaca aacgtttctct gacgtgtcag 480
gatctta 487

<210> 117

<211> 1014

<212> DNA

<213> Homo sapiens

<400> 117

ctcgtgccga atcttctaac aagagaacaa gctcctttct ttcttttcta aacaagggttc 60
agcgctttct attaaaagaa accctattca gaccctatgc agcacatagt ttataaaaa 120
ttttttctat taacagagga aaaataacct attgataaac agagcggtac aaggagatgc 180
aaataaagct gctttaggat cottacctag attctagaaa atggttgcat gaatttgaa 240
aaacaaacta attaaaaatt aaaactgaaa aaaatagttt aaacaaacaa ctgagagata 300
ttttttcatg ggcgtaaaa atacggcga aaatagact gacttgttgg aaagtatcca 360
atctatcaaa ttagaaaaa tgcggaagg taagcgcca gactccaata aagcagaaaa 480
aatggtctta atgaaaaaaa gcaagccgc tgctaaaaaa gctaaagctg ctgctaagaa 540
gcgtgttcgc gctacaaaaa cagtggctaa aaaagcttgc acaaaaagaa ttgtgtctac 600
taagacaaag gtcaaaccaa caaaaaaagc cgctctctaa acaaaagtta aaacagcgaa 660
aaaaactcgc tcaacaaaaa aataatattt tagcgcttct ttctttttat agaggcgcat 720
ttatacaaca gggccctctt tctctttctc attgatccct tctctttttt ttgttatcct 780
ttcgtttctc gcaaaaggca gtcccttgca ataaaagta acctcacac ctcttttggg 840
ggaaaaacct ttcactttct ttaggattca agttgctctc ctgctatgtt aactgttaac 900
atgttggtgt cgtggaggc tgttcattct cctaaaatga atatgcatcc ttctaaaaa 960
caaaagagct tgcgtcccat aatttatttg caccctctat cccatcccaa aata 1014

<210> 118

<211> 287

<212> DNA

<213> Homo sapiens

<400> 118

atgcgaataa agctgcttta ggatccttac ctgatttcta gaaaatgggt gcatgaattt 60
gaacaaacaa actaattaaa aattaaaact gaaaaaata gtttaaaaaa acaactagag 120
gatatttttt catggcgcta aaagatcagg caaaaaaaat gactgacttg ttggaaagta 180
tccaaacaaa ttgtcttaa gcagaaaaag gaataaagc cgcagcaaca agagtctcta 240
cagaatctat caaattagaa aagatcgga aggtatatgt taaagag 287

<210> 119

<211> 1002

<212> DNA

<213> Homo sapiens

<400> 119

catatgcac accatccca tccatgagt attcgaccta ctaatgggag tggaaatgga 60
taccgtctta ttaactctc taacgataat caatacggct ttgtgcaatc gacctctggg 120
octaatcaag gaggccatc ggtatctctc cgaggaggat ttoagggat atgcgtaaga 180
atagcgattt tattcgttaa ctgtttctct cgtaatagag gcaactactac tacgccatct 240
cgaactgtta tcaactcagg agatatttat caatcgacta ttcttggaac agggactcaa 300
octattgtct ctacaggaga taagaaatta gatagcgcaa tattcaagc atttttgtct 360
gcgcagaata aacagacttt ggtacacat attcaagta agctaggttc tatggaggga 420
caactctctc aagattataa agctgggtgc tatagtgcgc taagattgat gctgtttact 480
ccagcgaaa ctactgtgag tagcgagcgg gaactgcaa cgtgcgttac gggcgggat 540
ctctgggaac aggcgcagg agatcttgct acaaatggga atacagatgg gcttatgta 600
atggctaac tatctgtggg agggaaagat gtgcctgcgg ggcatttaag agaatacatg 660
gatactgtaa agggtacgtt tactgatgag aacgaggcta cagatcctac ggtatagcc 720
attttagatt tagcagcaaa aatcgatgcg acggaattct ctagtctgtg ttcagggcaa 780
gtcaatttta attatagga aaattatgga caagtctgtt tagaaaaaga ggagatgaac 840
cttctgtgtt tagaagatca aattgggcaa gatctctaac gtgttcaaga taactcaaaa 900

86

gagttacaaa aactgttaga aaatgctcga aaaacagatc ctgagttata ttcccaaaa 960
 ctaactgtca taacttcttc tgttttctta gactaaggat cc 1002

<210> 120
 <211> 1218
 <212> DNA
 <213> Homo sapiens

<400> 120
 atgcataacc atccaccatca cgtgagtagc ataagcccta taggggggaa ttctgggcca 60
 gagggatttt ctatgtgcac tcgaggcgat gagattgatg atgtaccaga tagtgaagag 120
 ggagagcttag aagagcgctt ttcggatcat gcagagtcta tcattaccga gagctcgaa 180
 acgctgttcc gtactacttc ttcatcaggg gtcagtgaag atcttcacga acacgttagc 240
 ttggaggaat ctccacgaca acgaggttcc ctggacgga tccgtgatgc agtagcttct 300
 atttggaaag gtctgtgtgc acgaaggaat gaaaactatg atgtgaaaaa agcagaagag 360
 cagcaaggga ttgtgcaata tctgcaggat tcgaaaatgc ctgctttaac gcgtgctat 420
 cgccattctcc gtgctttcaa ttctgcacgc ttacgtacga ttctgtagtt ttctgctacc 480
 atttttctgt tcttaaggga tgcgtattat cgacattgta cacgttctgg gatcaacttt 540
 tgtggagctg ataaagactc tttagaagtt ctgtgtgcgg tgggtttgtg ttgtcgatgt 600
 gcacacgtac gctcttttga acatgcoggt ggggaattac aagatcgatt agtaataaat 660
 gatgctccgg tgacaggtgc ggggagaact ctgtgtgatg atgctgtaga cgataatgaa 720
 tcgatttttaa atacagagaa caactggcct caacatgtca tgatagggtt ttctcggtgt 780
 ctgcttcaat tatgtgcgac tcttataaat ggcactcttc aagaattgtt caagtctgtt 840
 gttcgtttag aaaaagaaga ccttcttcca gattattctc aagctttatt attagcaggg 900
 ataatagatc gcttggcgga gaaagccctc atggctgcaa agtatgtttt ggaatcattg 960
 cgtgttcgaa ctctggagct cataggagaa ctcatattc tgcatttgtt tctctctgta 1020
 ttggaaggttg gcgcggagg cgtattccct cctgtgaatg agcagctcgt tgtgcaaat 1080
 gtttaatgcaa acgtagaacg attgcattcc acttctgctc atgagcaca agcttatttt 1140
 cgtatgatcg aaggttttgt aaccaatttc tttttctac ctgacgagga agatccttt 1200
 tcggttggga atatctaa 1218

<210> 121
 <211> 726
 <212> DNA
 <213> Homo sapiens

<400> 121
 catatgcac accatcacca tcacacaaag catgaaaac gcattctgtg tatccaagag 60
 acttacgatt tagctaagtc gtattctttg ggtgaagcga tagatatatt aaacagtgat 120
 cctactgtgc ttctcgatca aacggttgat gtgtctgta aattagggat cgatccaaga 180
 aagatgtatc agcaaatctg tggttcgttt tctttacctc acggtacagg taagtgtttg 240
 cgaatttttag tttttctgct tggagataag gctgcagagg ctattgaagc aggaagcgag 300
 ttgttgtgta gcgcagactt ggtagaaaaa atcaaaaggt gatgggttga ctctgatgtt 360
 tgggttgcca ctcccgatat gatgagagag gtgcgaaaagc taggaaaagt tttaggtcca 420
 agaaaccctta tgctacgccc taaagccgga actgtaacaa cagatgtgtt taaaactatt 480
 gcggaactgc gaaaaggttaa aattgaattt aaagctgac gcagctgtgt atgcaacgtc 540
 ggagttgcga agctttcttt cgatagtgcg caaatcaaa aaaatgttga agcgttgtgt 600
 cgagccttag ttaaagctaa gccgcgaact gctaaaggac aatatttagt taatttcaat 660
 atttctcaga ccatggggcc aggggttacc gtggatacta gggagttagt tgcgttataa 720
 gaattc 726

<210> 122
 <211> 330
 <212> PRT
 <213> Homo sapiens

<400> 122
 Met His His His His His Met Ser Ile Arg Pro Thr Asn Gly Ser

5 10 15
 Gly Asn Gly Tyr Pro Ser Ile Asn Pro Ser Asn Asp Asn Gln Tyr Gly
 20 25 30
 Leu Val Gln Ser Thr Ser Gly Pro Asn Tyr Gly Gly His Thr Val Ser
 35 40 45
 Ser Arg Gly Gly Phe Gln Gly Ile Cys Val Arg Ile Ala Asp Leu Phe
 50 55 60
 Arg Asn Cys Phe Ser Arg Asn Arg Gly Thr Thr Thr Pro Ser Arg
 65 70 75 80
 Thr Val Ile Thr Gln Ala Asp Ile Tyr His Pro Thr Ile Ser Gly Gln
 85 90 95
 Gly Ala Gln Pro Ile Val Ser Thr Gly Asp Lys Lys Leu Asp Ser Ala
 100 105 110
 Ile Ile Gln Ala Asp Leu Arg Ala Gln Asn Lys Gln Thr Leu Ala Thr
 115 120 125
 His Ile Gln Ser Lys Leu Gly Ser Met Glu Gly Gln Ser Pro Gln Asp
 130 135 140
 Tyr Lys Ala Gly Ala Tyr Ser Ala Leu Arg Leu Met Leu Phe Thr Pro
 145 150 155 160
 Gly Glu Thr Thr Val Ser Ser Glu Arg Glu Arg Gln Ala Cys Val Thr
 165 170 175
 Gly Arg Asp Leu Trp Glu Gln Ala Ala Gly Asp Leu Ala Thr Asn Gly
 180 185 190
 Asn Thr Asp Gly Leu Met Leu Met Ala Asn Leu Ser Val Gly Gly Lys
 195 200 205
 His Val Pro Ala Gly His Leu Arg Glu Tyr Met Asp Thr Val Lys Gly
 210 215 220
 Thr Phe Thr Asp Glu Asn Glu Ala Thr Asp Pro Thr Val Asp Ala Ile
 225 230 235 240
 Leu Asp Leu Ala Ala Lys Ile Asp Ala Thr Glu Phe Ser Ser Pro Gly
 245 250 255
 Ser Gly Gln Val Ile Leu Asn Tyr Ile Gly Asn Tyr Gly Gln Val Val
 260 265 270
 Leu Glu Asn Glu Glu Met Asn Leu Leu Val Leu Glu Asp Gln Asn Gly
 275 280 285
 Asn Pro Gln Arg Val Gln Asp Asn Ser Lys Glu Leu Gln Lys Leu
 290 295 300
 Leu Glu Asn Ala Arg Lys Thr Asp Pro Glu Leu Tyr Phe Gln Thr Leu
 305 310 315 320
 Thr Val Ile Thr Ser Ser Val Phe Leu Asp
 325 330

<210> 123

<211> 405

<212> PRT

<213> Homo sapiens

<400> 123

Met His His His His His Val Ser Ser Ile Ser Pro Ile Gly Gly
 5 10 15
 Asn Ser Gly Pro Glu Gly Phe Ser Ser Ala Ser Arg Gly Asp Glu Ile
 20 25 30
 Asp Asp Val Pro Asp Ser Glu Glu Gly Glu Leu Glu Glu Arg Val Ser
 35 40 45
 Asp His Ala Glu Ser Ile Ile Thr Glu Ser Ser Glu Thr Leu Phe Arg
 50 55 60
 Thr Thr Ser Ser Ser Gly Val Ser Glu Asp Leu Gln His Val Ser
 65 70 75 80
 Leu Glu Glu Ser Pro Arg Gln Arg Gly Phe Leu Gly Arg Ile Arg Asp
 85 90 95

[illegible]

```

      100      105      110
Gly Trp Val Asp Phe Asp Val Ala Val Ala Thr Pro Asp Met Met Arg
      115      120      125
Glu Val Gly Lys Leu Gly Lys Val Leu Gly Pro Arg Asn Leu Met Pro
      130      135      140
Thr Pro Lys Ala Gly Thr Val Thr Thr Asp Val Val Lys Thr Ile Ala
      145      150      155      160
Glu Leu Arg Lys Gly Lys Ile Glu Phe Lys Ala Asp Arg Ala Gly Val
      165      170      175
Cys Asn Val Gly Val Ala Lys Leu Ser Phe Asp Ser Ala Gln Ile Lys
      180      185      190
Glu Asn Val Glu Ala Leu Cys Ala Ala Leu Val Lys Ala Lys Pro Ala
      195      200      205
Thr Ala Lys Gly Gln Tyr Leu Val Asn Phe Thr Ile Ser Ser Thr Met
      210      215      220
Gly Pro Gly Val Thr Val Asp Thr Arg Glu Leu Ile Ala Leu
      225      230      235

<210> 125
<211> 713
<212> DNA
<213> Chlamydia trachomatis

<400> 125
ataacaatcc ctcccaatca tegtgtgaag tacaaggagg agccatctat gccaaaacct 60
ctttgtctat tggatcttcc gatctggaaa cctctctatat ttctcgggg aacagtgtct 120
ccactgggaa atctcaaaaca acagggcmaa tagcgggagg agcgatctac tccctactg 180
ttacatggaa ttgtctcgcg acattctcta acaatacagc ctctatagct acacggaga 240
cttctcttga agatggtatc tcaggaaatt ctattaaaga taccattgga ggagccatg 300
cagggcagac cattacccta tctggagctt ctgcattttc agggaaatag gctgatttag 360
gagctgcaat aggaactcta gctaagtcaa atacaccocag tgcaactagc ggatctocaa 420
atagcattac agaaaaaatt aotttagaaa agggttcttt tttttttgaa agaaaccaag 480
ctaataaagc tggagcgatt tactctccta gogtttccat taaagggaaat aatatatcct 540
tcaatcaaaa tacatccact catgatggaa gogctatcta ctttacaata gatgctacga 600
ttgagtcctt aggatctggt ctttttacag gaataaacgt tacagctaca caagctagtt 660
ctgcaacatc tggacaaaat acaataactg ccaactatgg ggcagccatc ttt 713

<210> 126
<211> 780
<212> DNA
<213> Chlamydia trachomatis

<400> 126
cctctctcct actcaggagt tttaaaagaa aacgcaccgt ttttacgttt cctcacacaa 60
ttaactaaca agcatactca ttctggattt cattgcctcc taaaatttct agtcaaatcc 120
gaaagaaagg gacactcgag cgtctctctc ctaaaaatct tgtttttctc ctgcttcoga 180
gttaatacgc ggctgtctca taaccacacac taacatgatg aaacctctac gtttcgggta 240
tttcttttgc acaatctatt ttaatttgyt acaggcagcg tttgctaaaag aacggaatc 300
ttctcccgag tgcagaata attggaagaa agtcaccocac acggatcaac tccctgaaaa 360
catcattcat gctgatgatg ctgtttatca ctctggttat gtacagggtc tcaattgata 420
gcatttctta gatagctgct gccaggtcat cgttgaaaac caaactgott acttattttc 480
tctctctaca gatgatgtta cgcgcacacg cattatcaac ctaattaaag accttccatt 540
catctactcc gtgaaaatct gccaaagcat ctatcaaaac tgtcatcatc aaggccctca 600
tggaagagct tctcttcag aacaacgttc ttctgtaca aaggtctygt gaaaagaagc 660
tatttgggta ccacagaata ccatctatt ctgcctcttt gtacgagata ctctcaagc 720
aactaatagt gcagggtatc gttttaacga cgaagtcgta ggaaaacggt ttggctctgc 780

<210> 127
<211> 433

```

<212> DNA

<213> Chlamydia trachomatis

<400> 127

```

ctttaaagat  togtgtctct  tttggtacta  cgagagaagt  togtgtgaaa  tggcggttatg  60
ttcctgaagg  tgtaggagat  ttggctaacc  tagctccttc  tatcaggggct  ccacagtttac  120
agaaaatcga  gagaagcttt  ttccctaaga  aagatgatgc  gtttcatcgg  tctagtctgc  180
tattctactc  tccaatgggtt  ccgcattttt  gggcagagct  tcgcaatcat  tatgcaacga  240
gtgggttgaa  aagcgggtac  aatattggga  gtacogatgg  gtttctccct  gtcattgggc  300
gtgttatatg  ggagtcggag  ggtcttttcc  gcgcttatat  ttcttgggtg  actgatgggg  360
atggtaagag  ccataaagta  ggattttctaa  gaattctctac  atatagtgtg  caggacatgg  420
aagatttga  tcc  433

```

<210> 128

<211> 803

<212> DNA

<213> Chlamydia trachomatis

<400> 128

```

aatctattaat  taatagcaag  cttgaaacta  aaaaacctaat  ttattttaaag  ctcaaaaataa  60
aaaagagtttt  taaaatggga  aattctgggt  tttatttga  taacactgaa  aactcggtct  120
ttgtgtatata  tatcaaaagt  gggcaaatga  cagagcgctg  caaggaccag  caataatctac  180
ttgggacacac  atccacacot  gtgcgagcca  aaatgacagc  ttctgatgga  atatttttaa  240
ctactctccaa  taattctaat  accaatgctt  ctattacacat  tgggttggtat  ggggaaaaag  300
ctaacacgct  tattctagaa  aagttgggag  atcaaatctt  tgatgaaatt  gctgatacta  360
ttgtgtatag  tacagtccaa  gatattttag  acaaaatcaa  aacagacccct  tctctaggtt  420
ttgttgaaagc  ttttaacaaa  ttctcaatca  ctaataaaat  tcaatgcaac  ggggtattct  480
ctccacagtaa  caattgaaat  ttattaggag  gaactgaaat  aggaaaatto  acagtcaaac  540
ccaaaagcttc  tgggagcatg  ttcttagtct  cagcagatat  tattgcatca  agaatggaag  600
ggcgcggtgtg  tctagctttg  gtacgagaag  gtgattctaa  gcctcgcgcg  attagtgtatg  660
gatactcatc  aggcatttct  aatttatgta  gtctaagaac  cagttattact  aatacaggat  720
tgaactccagc  aacgtattca  ttaactgtag  goggtttaga  aagcggtgtg  gtatgggtta  780
atgocctttc  taactctgtg  cgg  803

```

<210> 129

<211> 842

<212> DNA

<213> Chlamydia trachomatis

<400> 129

```

tgggaatgtc  gaagaaatag  attacgtttc  cgtatctata  ggacgcoggt  tgaatacaga  60
aaatatttgc  ttggataaag  ctgggtgtat  ttgtgatgaa  cgcggagatca  tccatcacga  120
tgccacaatg  cgcacaaaag  taacctaacat  ttatgctatt  ggagatatca  caggaaaaatg  180
gcacacttgc  catgtagctt  ctcatcaag  aatcatttga  gcacgggaata  tagdtggcca  240
taaaaggagaa  atcgattact  ctgcggtccc  ttctgtgato  ttacotttcc  ctgaagtctgc  300
ttcagtaggc  ctctcccaaa  cagcagctca  acaacaaaaa  atccocgtca  aagtaacaaa  360
attccatttt  cagcgtattg  gaaaagcggt  cgcaatgggc  gaggccgatg  gatttgacgc  420
cattatcagc  catgagacta  ctacgcagat  cctaggagct  tatgtgattg  gccctcatgc  480
ctcatcaactg  atttccgaaa  ttaccctagc  agtttgtaat  gaactgactc  ttcoctgtat  540
ttaacgaact  atccacgcac  atccaaacct  agcagaagtt  tgggctgaaa  gtgcgttgtt  600
agctcgtgat  accccattac  atatgcccc  tgctaaaaaa  tgaccgatc  agaattctct  660
actcctaaaa  aatctatacc  cgcagatttc  cctaaatggc  tacgcagaaa  acctocctta  720
ggcggggtat  ttgctcaaac  tgataatact  atcaaaaata  aagggtctcc  tacagtctgt  780
gagggaagcct  ctgtctcgaa  tcgcacccat  tgttgtctca  gacatacagc  tacatatcta  840
gc  842

```

<210> 130

<211> 813

<212> DNA

<213> Chlamydia trachomatis

<400> 130

aaaataacttt	gagctgcaca	agctccccc	tgtttagag	aagaacatga	tgcaaatcc	60
aatccaccct	taattctttt	aaagataaga	ctctctgtag	aataaaagc	cgctccagac	120
aaagaagctt	tcaactgcagt	taaagtgtatt	ccagccttac	tactatcccc	aaacaaagca	180
atacttaaaa	aagattctoc	gtcacgagga	gaatcaaggt	tgctgctcgt	aaactacaa	240
attaacccctt	gggaagagac	ttgatctcgt	tgggtccacac	cttggaanaac	taaggagattg	300
gttactgaga	acaaagtaact	ttgctctacc	ttacccgggaa	gagtatccgc	atctttctct	360
tggaaagaa	ttggatctcc	tacaattaac	ctatactgtc	cttcagcctg	actattctta	420
gaccacaacga	atagatctcg	aatttggctc	aaacaataaaa	ccgcttgagg	gcctacatat	480
accagctcat	ttacagactg	tctccagcga	tgaagatcta	cgcaactagc	taacccgcta	540
acagaggcaa	ggatagctgc	tactacagac	aaagaaaact	tagaacaggt	gcttlttata	600
tctttctcgg	aactcatttc	aaacctgcga	aatagcaact	ttttgacaaa	ctagcgtacc	660
gaacaaatcg	gtcccaacac	gogtctgcgc	latgalttca	caagacacaa	acgcacata	720
gacaagctcc	agagacgaca	ttagagcttt	agaccgtgga	atgtacaatg	ctgactgctt	780
tttgagaaa	atttttata	aagaacaggc	act			813

<210> 131

<211> 1947

<212> DNA

<213> Chlamydia trachomatis

<400> 131

tcttttgcct	atagagcaat	cttttatcat	tgggtctgat	ccaccagact	attttcttta	60
gagtagagatt	ctactacccc	atccatggca	ttcaacctct	catcagtaaa	cactttatta	120
gagtggttta	cttgcccatc	atcgatgata	tctttctgaag	tcttttaatac	cttttatacat	180
aagatccatc	tctccggaga	acagtgctct	tctatggata	aaatttccat	cgagataatt	240
acgcatccca	aaatagcagg	aatacctaga	tagatggcat	ttacaaaoga	agctgcogaa	300
actaggaata	tcaaacgagt	aatcactaaa	agtagtccta	tcaccaactaa	tcocacotta	360
aatgcagctgg	aagatagaag	attcgatata	cgctctttca	gigttaatgg	tgcagaacta	420
gttggaatat	octgtgcoga	attggaagat	ccagctcctt	gaacaaagg	tacagtgtcc	480
atatttttaca	ttoctttttt	ggttgtgagc	agggagtota	cacaacacat	tatttttttc	540
aaaaaacctg	ctagaatatg	ctctgagacc	gaaatgaac	tcttttattt	tcactatagat	600
aaaaaaaaaa	agccgcccag	gaatccctgg	acggcacota	cacatcgata	aaatcaaaaga	660
ttaatatagat	tgtgtattct	ctgtatcaga	aactggaaaca	gtcaatgtat	cggaagaaga	720
aatgccttcc	ccacagcat	ctccagctga	tactgctttc	aaigttaacg	aaaactctac	780
agttttcttta	gaacctaatc	taggttaacga	atcgaatact	actgtatttc	ctgttaactgt	840
tcttttagtt	ggctccagaga	aggtatcagg	ttgcagttct	ttagagaatt	taagcatttaa	900
agaaacattt	gtatcttctg	cagaacctct	gttggtgaca	caaataccgt	aaacagtatt	960
tctctctaca	caaacagggt	cacaagatc	tactacgcac	atatgagtag	cagcaactcc	1020
tttccagtaa	gttgtctgct	ctgcgcaaga	agtacaagta	ccacagctcag	agcagctctt	1080
cacaacaacaa	tattttgtga	attgtccagg	agttttgtct	cttactagaa	ctttataactg	1140
tagagactct	ccaggattca	gttctttcac	agtcacaact	actttattac	aagaaatttg	1200
agctccctgca	gttccagaag	ctgtgactcc	ggggaagaaga	gtgtcttcaa	cgagacactc	1260
tgcacaacaca	agatctccag	gattggaaac	ggagatcaca	tattctacag	gotttacaac	1320
ataagaccaa	tctgtctcgt	caatactttac	ttgtacgcaa	ggctcatiga	tcacagttgt	1380
tacgcttctg	gtatttttat	gtctctccaa	gtaagaaccc	gttgctatat	tgttagcacg	1440
accagcttca	agcggacaaa	actctacagt	aattgttctg	tgtctccag	ctgtgcatac	1500
tccaggagta	aactgcagta	cacgctgtcc	agaagagtga	gcgttaacca	ctggaaacag	1560
atttccaaca	acaacggtac	gagctattgc	tgttctcttg	tctactacat	taatttttga	1620
aactactggg	caacgcacac	aagcatcttc	tgggctctct	tgtttaaac	agatagcagc	1680
ttgtccacat	ttgttaacog	aacggatctc	tggacaagcg	catactgttg	cgctgtataa	1740
cgagcaacct	tcttttaag	gttttaccac	tacagtaatt	ttactctttt	cgctgtgtcc	1800
taagcgtgtca	attttccaaa	ctagcttaac	atcagcagta	ggagttgtcg	ctgtactact	1860
ggttacgaac	tctgtctcac	atggttaatt	ctgagtaatt	ataacatcaa	cacaatccct	1920
tttactctga	cgagtaattt	caatagg				1947

<210> 132

<211> 1278

<212> DNA

<213> Chlamydia trachomatis

<400> 132

```

gataacaaaa aaaaagccgc caggaatccc tggacggcac ctacacatcg ataaaaatcaa 60
agattaataag atgtgtgtat tctctgtatc agaacttgga acagtcacatg tatcggaaga 120
aagaatcgctt tccccacgag catctccagc tgatactgct ttcaatggtta cagaaaaatc 180
tacaagttctt ttagaaccta atctaggtaa cgaatcgaaat actactgtat tgcctgtaat 240
cgttccttita gttggtccag agaaggtatc aggttgcaat tcttttagaga atttaacgat 300
taaaagaaca tttgtatctt ctgcagaacc tctgtgtgtg acacaaatca ggtaaacagt 360
attttctcct acacaaacag ggtcacaaat atctactaag cacatatgag tagcagcaac 420
tcttttccag taagtgtgtg ctctcgcgca agaagtacaa gtaccacagt cagagcagt 480
cttcacaaca acattatttg tgaattgttc aggagtttgt gctcttaact gaactttata 540
ctgtagagac tctccaggat tcaatttctt cacagtccaa actactttat tacaagaat 600
ttgagctcct gcagcttcaa gaactgtgac tccgggagaa agagtgtctt caacgacgac 660
atctcgcaac acaagatctc caggatttga aacggagatc acatatttca caggcttaca 720
aacataagac caatctgctc ctgcaatact tacttgtacg caaggctcat tgatcacagt 780
tgttacgctt gctgtatttt tatgtctctc acagtaagaa acogttgtca tattgtgtac 840
acgacaacgt ttaagcggac aaaactctac agtaatttgt tccagaagag tgagcgtaac catctggaac 900
atctocaaaga gtaaacgcca gtacacgctg tccgggagaa agagtgtctt tggttcaact cattaat 1020
aggattttca acacaacagt tacgagctat tgctgttctt tctgttctca cattaat 1080
gtaaactact gggcaacgca acaacgcatc ctctgggcat tctgttttaa cacagatagc 1140
asggtgttcca cattttgtta ccgaacggat ctctggacaa gcgcatactg ttgcagctgt 1200
aacgaagcaa cctcttttaa gaggttttac ccatacagta attttactct ttgcgctgt 1260
tctaagcggc tcaattttcc aaactagctt accatcagca ttaggagttg tctgtgtgac 1320
actcgtacg aactctgc

```

<210> 133

<211> 916

<212> DNA

<213> Chlamydia trachomatis

<400> 133

```

atggcgacaaa ttttaacgatt acoggacaaa accatacatt atcatttaca gattctcaag 60
ggcaggttct tcaaaattat gccttcattt cagcaggaga gacacttact ctgaaagatt 120
tttcgagttt gatgttctcg aaaaatgttt ctgcgggaga aaagggaatg atctcagaga 180
aaaccggtgag tatttccgga gcaggcggaag tgattttttg ggataactct gtggggtatt 240
ctcctttgtc tatttgcgca gcactcgaac caactctctc agcaccagca ccagctctcg 300
ctgcctcaag ctcttttatc ccaacagtta gtgagtctcg gaagggtct attttttctg 360
tagagactag ttttgagatc tcaggcgcca aaaaagggtt catgttcgat aataatgcgc 420
ggaaattttg aacagttttt cgaggtaata gtaataataa tgcggtagtg gggggtagtg 480
ggctcgtctac aacaccaagt tttacagtta aaactgttaa agggaaagt tctttcacag 540
ataacagtagc ctctgtgga ggcggagtag tctcaaaagg aactgtgctt tcaaaagaca 600
atgaagaggag catattcttc cgagggaaca cagcatacga tgaatttagg attctgtctg 660
ctactagctc ggtacgaat acgagacagc gaggcggtgg aggagttatt tgcctccac 720
atgattctgt aaagtgttaa ggcataaagg gttctattgt ttttgattac aactttgcaa 780
aaggcagagg cggaaagcat ctaacgaaaag aattctctct ttagcagat gattcgtgtg 840
tcttttagtaa caatacagca gaaaaaggcg gtggagctat ttatgtctct aogtatcgat 900
ataagcacga atggag

```

<210> 134

<211> 751

<212> DNA

<213> Chlamydia trachomatis

<220>

<221> misc_feature

<222> {1}...{751}

<223> n = A,T,C or G

<400> 134
 agcctctggc gaaggagagc cataaaaaagt gctaccagc ggagaacaa taaaatctcc 60
 ctgagcagcg aactcaacttt ctttctctctc gatactctct ttaacaatag gattcccaag 120
 gttttgatctt gaggataaagt ttgaaatoc agaaaacagt ctgttatcat aaaagactgg 180
 ctctctgaata ctctgggaact tatccctttc taactctaac tccaaacott cactgttgat 240
 aacaatggcg ttccagtgcc gaattcggca cgaggctott tttacgagg atctogagtc 300
 aagaagcctt gagccttcaa ttcttgcttc atgtctcttt tctcttgag aacagctcta 360
 gctaaaccca atcgagttag aataacctga ctttgaaccc ctctccact tactcggaata 420
 atcaaatcga aactgttagc atcaaccgag attctgagcg gagctaaagt ggttgctctt 480
 tgaacttcaa gagggaata ttgctctaaa gtctttccat ttacgtcaat ttctctatc 540
 ccagaacgaa gacgaacga cactcgtctt cttctgcctg ttgcaacaga ctcttgctatc 600
 atattctttg tcaacaaatta ccccaaatca cgcgtctaaa acaattgggt tgatagcttc 660
 ataotgtgct taagaactac ctttcaaaac ttctaaagat ttcatttgac gtcttccaa 720
 ttttggttta ggcaacattc nttaacagca t 751

<210> 135
 <211> 410
 <212> DNA
 <213> Chlamydia trachomatis

<400> 135
 ataactcaga ctctctctca totggagata ggcctggaga ctctgaagaa ctgactgaga 60
 cagaagctgg ttctacaaca gaactctcta ctttaattag aggaggctgt actatggag 120
 aaactgttaa gattgagaac ttctctggcg aaggaattct tcttggaac aaagtatag 180
 ataacaacac agaaagctcc ttctccaaat ctgacgtctt cggaggtgcy gctatgcta 240
 aaactgtgtt taatctcgat agcgggaact ctgacgaac tgcacottc tccgggaata 300
 ctgtctcttc tcaatctaca acaggtcagg ttgctggagg agctatctac tctctactg 360
 taacattgc tactctgta gtattttcta aaaactctgc aacaaacatt 410

<210> 136
 <211> 2719
 <212> DNA
 <213> Chlamydia trachomatis

<400> 136
 ctogtgcgca aaagctttct gctctacca agagattcgt tttttaaatt cttcattctc 60
 tctaaagagat ttagtttctt tcgcagaa ca attgatagat actcogtaog ttgggggtgg 120
 ccggtgcatt cataaacagc ttctctgtaa tgggtgatag tgttcggggt atattcaact 180
 actttacca gtcacaggaa gaaatatccc tcgcaatgct agagatcaat acagagactg 240
 ttctccagta aaagattttct cgtctctacc tataggagga ctatctctcc taagaagaagc 300
 aagcagcgga caaatcaacc atgttatgat gaaaactctcg gagcatgaat tcatctatgc 360
 tgogtaaaaa atagggaag tagaaaaagt aatcctagga aatagggtct tctttaaagg 420
 gaattctatt tgcctattag gtgaacgcgc tatagaagct gtttttggcg tctctaaaaa 480
 tagaaaagcc ttcttttgaa agaaggtctt tctgaaacgc actccaat atggacaagc 540
 aatagcttat cgtttggaga attggaact cttaagact tcttaacgac cgtatttttt 600
 acgctctttc ttacgaggat ctgcagtoaa gaagccttga gcttcaatt ctgtctctac 660
 gtctcttttc ttctgcagaa cagctctagc taaacccaat cgagtagcaa taaotgaoc 720
 ttgaacccct cctccactta ctccggaat caaatcgaaa ctgttgacat caccgagcat 780
 ttgagcgga gtaagatgg ttgctcttg aactcaaga gggaatat tt gctctaaagt 840
 otttccattt aogtcaattt ttccattccc agaaggaaga cgaaagctag aaacagcgt 900
 otttctctcg ottgttgcaa cagactottg tatcatatto ttgtcaca ataccocaa 960
 attagcgtc taaaaacatt ggtttgatag ctctcaactg tgcgtaagaa ctaccttca 1020
 aaactcttaa agatttcaatt tgacgtcttc caagttttgt tttaggcaac attcctttaa 1080
 cagcatgctc gataacataa gcaggcttcc gogcaatcat gttttcaaaa ggaaactctt 1140
 cactccaga aataaagcct gtgtaaatag gatacacttt ctgagttcct ttggtccag 1200
 tcaaacgcac ttctcagca ttgatcaca tgacacacat tccactcgt acgtgagag 1260
 taaaagtcac ctatgctta cctctcagga tctctgaaac tctggaagt aatctcccta 1320
 aggtcttccc ttcagatta actacataac aggtcttgtt tgcgtctcc gaagccttag 1380

ctagggtcgt	tttctgatct	tttctttttt	ccataaacta	aatcacctta	tcagaggtaa	1440
tgatltataat	tttgatgatt	atttttttca	aacaaaagcg	agctgtattt	gccttctaaa	1500
gaatttagaa	aagaaaaaat	ttcaaaaaag	tctctttttt	ttttgcctto	aaaaacagcc	1560
ttcacctctc	atacttcttt	cgaaaaaata	tttttaggaa	gttcttgaat	catgatttac	1620
ataataaaaa	aaaatgtllag	ctgcacacag	ctaaatttaa	aaaggtgcta	ccagacgcta	1680
aaagctggctc	caacgtaatta	atatcataat	cagaaaagag	aaacttcgga	ttatccaaca	1740
tgaactgatg	aaaaggaaatt	gtagaatgca	cccacacaa	atggaaactct	tttaagcttc	1800
ttttcataat	ggctatcgct	tctcttcgat	tctttctott	tgtagattacc	ttlagcaatca	1860
tggaatcata	ataaggaggt	atgcataaac	cactgtagca	agcccgctct	actcgacgac	1920
caggacccctg	aggaggagga	taataatcta	atctacacgg	ggaaggagta	aagtatttaa	1980
ttgatctctc	tcgattgatt	cggcattgaa	tcacgtgccc	tttaaaactct	atatctcttt	2040
cgctccaagg	cagttttttc	cccttagcga	cactaatctg	agcctttaac	aaatgatcat	2100
ctgtcaccttc	ttctcgtaata	gtatgttcca	cttgatagcg	cgatttcato	ttcatgaat	2160
aaaaacgctt	ctccttatct	aacagaaatt	ctactgttcc	aacagagaaa	taccggcacc	2220
tcgagcctaa	atccactgct	acttttccaa	cttlagctcg	catttctgga	gttaaaatag	2280
gacttggagt	ctcttctatt	aatttttggc	gacgctttg	tactgacaa	ctcgttctcc	2340
aagatacacg	taatttccgt	gcttatctcc	aattacttga	acttctaact	gtcttgatt	2400
ttcaataaat	ttttcaatat	acacgtcagg	attattaaat	ccogctctgt	cttcagcccg	2460
agcgggcaga	agaagccctat	agaatttgc	ttttctctca	acaatcogta	ttctcgtctc	2520
accgctctcca	gcacacagctt	tgatgaogat	ggggaatcog	atctttctgt	caattctaat	2580
cccttccacc	tcactctcca	ctacaccttc	agatccaggg	attacagggc	acttaactct	2640
tttagccaac	tgcttagctg	cgactttatc	tcccatagtc	gctatcgact	cagcactagg	2700
accgataaat	ctcgtgcgcg					2719

<210> 137

<211> 2354

<212> DNA

<213> *Chlamydia trachomatis*

<400> 137

gtgcaagatg	ggacgagttt	gaagtttaat	actagocacat	aaotctocctt	ctggagggttt	60
aggagagagc	cccttttatta	gggctctctt	tttttgtgtg	tgaggaaagc	tagcgtctaa	120
ctaaaagtct	ctaagttaagg	atgttttttag	gggaaatagc	gatittcagt	gttgagaagc	180
ttagtacaaa	gacataaaac	aaggctaaaga	aaaacttttc	ttagcctgtg	ttctccaagc	240
atcgccctata	gaagactaat	cttccagcgt	tgccctatgg	ctcagcttca	actggcgtctt	300
ttcgttaatg	ctaaaggagtt	taacagcaag	cttgttctct	tttttgacaa	agccacagat	360
attgtctact	ttttgttttag	acaattccaga	aatatgacag	agccctcttt	ttctcgggag	420
gacttctaac	aatactccaa	atgtttgcgat	agatgtaaca	cgcccatatt	aaactttacc	480
gacttccaact	tctccagttta	atccttctgat	aagtttcttia	gctttgttaa	tcgattttctg	540
gtggctttgca	gctatgttaa	tgacgcgcctc	atcattgatg	tcaacttgcg	cacocagaagc	600
ctcgataaatt	tgacggattt	gttttctctc	gggaccaaag	accgttgcga	tttttgaggt	660
attgatctgc	atagtttcaa	tcggcgggagc	atatttagaa	acagttccct	tagggggagcg	720
cagaacctgtg	gcataaagat	taaggatatg	actacgacct	tgtttagctt	gcgcagagcg	780
ttgtctccata	atcttatgatg	tgatttccctc	tatcttgata	ttcatttgga	aagctgttaat	840
accttttagct	gttccgggcta	ctttaaagtc	catatctctc	agatgatctt	ctatcagaga	900
aatatacgac	aagatgatgg	cttgatctcg	atctaagatt	aagcccatag	caatacctgc	960
cacggggagct	ttgataggaa	ctccagcctc	catgagtgca	agacagctcc	cacatacggga	1020
tgccatggag	gaagatccat	tagactcagt	aatattagat	tctaggcgaa	tgataaagg	1080
gaatcgcgat	gtctcaggaa	gaacatgact	taaaagctttc	tcagctaatt	ttccatgtcc	1140
aaatttcaagt	cttctctggg	aaccaattct	gcacactctt	ctacgcgaga	aaggagggaa	1200
gaataactgt	agatagaagc	gagcggctcc	atctccatct	agatcttoga	atcgtctgtc	1260
gcatttttgc	ctctccaaagc	tacatacggc	catgctttgc	gtctctccgc	gagtaataaa	1320
cacacttcog	tggtttctctg	gaagaaaaag	agctctctat	gaaatggggc	gaatctctgt	1380
ggtgtgttctg	ccatctacac	gaataccaag	atcttggtata	agagctcgca	tttgatttga	1440
ttttgtctgc	ttaaatgcag	cccttaacgtt	caacaaagaa	aaatcactgt	tttcttctgt	1500
aaaccaagtta	gcaataacgg	attcctctaa	ttcttctcag	gcttgctcta	gagctttctt	1560
atctctaaaa	gacaaatgctt	tttgcgaatt	ttctctaaaa	aaactctgaa	ctacattttg	1620
tcagctctct	ggcatatcaa	gacggcoga	gaattctttt	tggttgcgga	tagctttctg	1680
ccatgcttca	atgacatcgc	atattttagc	tatatagatt	tgcccaaaaa	caatagcttc	1740
tagaacttgc	ttctctgtta	aaaagtgcga	atgtctctca	atcattaaaa	ctgcagaagc	1800

```

tgtctctgcc atgacgagat ccagcctgga ggcacttaac tcatctctgg ttgggttaat 1860
gacccacttt cctcgcagca gcccaacgcg tacaccgcga acgatacaat ttgaggaaac 1920
ctctgagata gctaaagcgg cagaagctcc gcaaatagct agaggatcag gtaaatgttt 1980
cccgctgtaa gaccaaacgt aggacaagac ttgaatatct tgcattgagtc tattaggaaa 2040
cgacggcagc aaagagcgat ccattagccg agaaacaaga attctctctc cggagggcgg 2100
tctctcaagt tttagaaatc ctccagaggt tcttctctcg gaggaataat tctcttgata 2160
gtctactctg aaaggcagaa aatcgacagc ctctgacaag gaggtgcac acgctgaaga 2220
aaaaacccaa gtctcgttca ttttgacgag aacagcccca ctggcctggc gagctatttt 2280
ccctgtctcg aaaattaatg ttttattttt gtctaaogca acagaaaaag tctcaaaagc 2340
catggagttg tctct 2354

```

<210> 138

<211> 898

<212> DNA

<213> Chlamydia trachomatis

<400> 138

```

tcatcttgct tcatatttcc ggtatagaag atcatctagg agatattggac tttaaagtag 60
ccggaacacg taaaggtatt acagctttcc aaatggatat caagatagag ggaatcaact 120
ataagattat ggagcaagct ctacggcaag ctatacaagg cggtagtcat atccttaact 180
ttatgacaca ggttctggcc tcccttaagg gaactgtttc taaatatgct cggcgatgtg 240
aaactatgca gatcaatacc tcaaaaatcg caacggtcat tggccccgga ggaatacaaa 300
tcctgtcaaat tatcgagcgt tctgggtgag aagttagacat caatgatgac cgggtcaata 360
acatgactgc aagcaacccaa gaatcgatta acaaaagctaa agaacttato gaaggattaa 420
ctggaggaagt tgaagtcggt aaagtattata atggccgtgt tacatctato gcaacatttg 480
gagatttcgt agaagtcctc ccaggaaaaa aaggccctctg tcatatttct gaattgtcta 540
aacaaaagt agacaatata tctggcttgg tcaagaagg agacaagctt gctgttaaac 600
tcttagactat taacgaataa ggcagttgga agctgagcca tagggcaag ctyggaagatt 660
agctctctat agcgactatg ttgagaacaa aggtataaga atttccctta tagcttgtt 720
tattgtcttg taactaagct tctcaacact gaaaatcgct atttccctta aaaaactaat 780
taactagaga caattagtta gacgttagct ttcctcacac acaaaaaaag agagcctcaa 840
taaaagggct ctctctctaa cctcagaag ggaagttatg tgctagtatt aaacttca 898

```

<210> 139

<211> 660

<212> FRT

<213> Chlamydia trachomatis

<400> 139

```

Met His His His His His Met Glu Ser Gly Pro Glu Ser Val Ser
      5      10      15
Ser Asn Gln Ser Ser Met Asn Pro Ile Ile Asn Gly Gln Ile Ala Ser
      20      25      30
Asn Ser Glu Thr Lys Glu Ser Thr Lys Ala Ser Glu Ala Ser Pro Ser
      35      40      45
Ala Ser Ser Ser Val Ser Ser Trp Ser Phe Leu Ser Ser Ala Lys Asn
      50      55      60
Ala Leu Ile Ser Leu Arg Asp Ala Ile Leu Asn Lys Asn Ser Ser Pro
      65      70      75      80
Thr Asp Ser Leu Ser Gln Leu Glu Ala Ser Thr Ser Thr Ser Thr Val
      85      90      95
Thr Arg Val Ala Ala Lys Asp Tyr Asp Glu Ala Lys Ser Asn Phe Asp
      100      105      110
Thr Ala Lys Ser Gly Leu Glu Asn Ala Lys Thr Leu Ala Glu Tyr Glu
      115      120      125
Thr Lys Met Ala Asp Leu Met Ala Ala Leu Gln Asp Met Glu Arg Leu
      130      135      140
Ala Asn Ser Asp Pro Ser Asn Asn His Thr Glu Glu Val Asn Asn Ile
      145      150      155      160
Lys Lys Ala Leu Glu Ala Gln Lys Asp Thr Ile Asp Lys Leu Asn Lys

```

Leu	Val	Thr	Leu	65	Gln	Asn	Gln	Asn	Lys	170	Ser	Leu	Thr	Glu	Val	Leu	Lys
Thr	Thr	Asp	Ser	180	Ala	Asp	Gln	Ile	185	Pro	Ala	Ile	Asn	Ser	Gln	Leu	Lys
Ile	Asn	Lys	Asn	195	Ser	Ala	Asp	200	Ile	Ile	Ala	Lys	Asp	Leu	205	Glu	Gln
Asn	Ile	Ser	Tyr	210	Glu	Ala	215	Leu	Thr	Asn	Ala	Gly	Glu	Val	Ile	Lys	240
Ala	Ser	Ser	Glu	225	Ala	Gly	Ile	Lys	Leu	Gly	Gln	Ala	Leu	Gln	Ser	Ile	240
Val	Asp	Ala	Gly	230	Asp	Gln	Ser	Gln	Ala	Ala	Val	Leu	Gln	Ala	Gln	Gln	255
Asn	Asn	Ser	Pro	245	Asn	Ile	Ala	Ala	Thr	Lys	Glu	Leu	Ile	Asp	Ala	Ala	270
Ala	Glu	Thr	Lys	260	Val	Asn	Glu	Leu	Lys	Gln	Glu	His	Thr	Gly	Leu	Thr	285
Asp	Ser	Pro	Leu	275	Val	Lys	Lys	Ala	Glu	Glu	Gln	Ile	Ser	Gln	Ala	Gln	300
Lys	Asp	Ile	Gln	290	Glu	Ile	Lys	Pro	Ser	Gly	Ser	Asp	Ile	Pro	Ile	Val	315
Gly	Pro	Ser	Gly	305	Ser	Ala	Ala	Ser	Ala	Gly	Ser	Ala	Ala	Gly	Ala	Leu	330
Lys	Ser	Ser	Asn	325	Asn	Ser	Gly	Arg	Ile	Ser	Leu	Leu	Leu	Asp	Val	Val	340
Asp	Asn	Glu	Met	335	Ala	Ala	Ile	Ala	Leu	Gln	Gly	Phe	Arg	Ser	Met	Ile	350
Glu	Gln	Phe	Asn	340	Val	Asn	Asn	Pro	Ala	Thr	Ala	Lys	Glu	Leu	Gln	Ala	365
Met	Glu	Ala	Gln	355	Leu	Thr	Ala	Met	Ser	Asp	Gln	Leu	Val	Gly	Ala	Asp	380
Gly	Glu	Leu	Pro	370	Ala	Glu	Ile	Gln	Ala	Ile	Lys	Asp	Ala	Leu	Ala	Gln	395
Ala	Leu	Lys	Gln	385	Pro	Ser	Ala	Asp	Gly	Leu	Ala	Thr	Ala	Met	Gly	Gln	410
Val	Ala	Phe	Ala	390	Ala	Ala	Lys	Val	Gly	Gly	Gly	Ser	Ala	Gly	Thr	Ala	425
Gly	Thr	Val	Gln	405	Met	Asn	Val	Lys	Gln	Leu	Tyr	Lys	Thr	Ala	Phe	Ser	440
Ser	Thr	Ser	Ser	410	Ser	Ser	Ser	Tyr	Ala	Ala	Ala	Leu	Ser	Asp	Gly	Tyr	455
Ala	Tyr	Lys	Thr	420	Leu	Asn	Ser	Leu	Tyr	Ser	Glu	Ser	Arg	Ser	Gly	Val	470
Gln	Ser	Ala	Ile	435	Ser	Gln	Thr	Ala	Asn	Pro	Ala	Leu	Ser	Arg	Ser	Val	485
Ser	Arg	Ser	Gly	440	Ile	Glu	Ser	Gln	Gly	Arg	Ser	Ala	Asp	Ala	Ser	Gln	495
Arg	Ala	Ala	Glu	450	Thr	Ile	Val	Arg	Asp	Ser	Gln	Thr	Leu	Gly	Asp	Val	505
Tyr	Ser	Arg	Leu	460	Gln	Val	Leu	Asp	Ser	Leu	Met	Ser	Thr	Ile	Val	Ser	515
Asn	Pro	Gln	Ala	470	Asn	Gln	Glu	Ile	Met	Gln	Lys	Leu	Thr	Ala	Ser	Ala	525
Ile	Ser	Lys	Ala	480	Pro	Gln	Phe	Gly	Tyr	Pro	Ala	Val	Gln	Asn	Ser	Ala	535
Asp	Ser	Leu	Gln	490	Lys	Phe	Ala	Gln	Leu	Glu	Arg	Glu	Arg	Glu	Val	Asp	545
Gly	Glu	Arg	Ser	500	Leu	Ala	Glu	Ser	Gln	Glu	Asn	Ala	Phe	Arg	Lys	Gln	555
G25	Pro	Ala	Phe	510	Ile	Gln	Gln	Val	Leu	Val	Asn	Ile	Ala	Ser	Leu	Phe	560
				520													575
				530													585
				540													595
				550													605
				560													615
				570													

Gly Tyr Leu Ser
660

<210> 140
<211> 598
<212> PRT
<213> Chlamydia trachomatis

<400> 140
Met His His His His His His Met Ser Ile Arg Gly Val Gly Gly Asn
5 10 15
Gly Asn Ser Arg Ile Pro Ser His Asn Gly Asp Gly Ser Asn Arg Arg
20 25 30
Ser Gln Asn Thr Lys Gly Asn Asn Lys Val Glu Asp Arg Val Cys Ser
35 40 45
Leu Tyr Ser Ser Arg Ser Asn Glu Asn Arg Glu Ser Pro Tyr Ala Val
50 55 60
Val Asp Val Ser Ser Met Ile Glu Ser Thr Pro Thr Ser Gly Glu Thr
65 70 75 80
Thr Arg Ala Ser Arg Gly Val Leu Ser Arg Phe Gln Arg Gly Leu Val
85 90 95
Arg Ile Ala Asp Lys Val Arg Arg Ala Val Gln Cys Ala Trp Ser Ser
100 105 110
Val Ser Thr Ser Arg Ser Ser Ala Thr Arg Ala Ala Glu Ser Gly Ser
115 120 125
Ser Ser Arg Thr Ala Arg Gly Ala Ser Ser Gly Tyr Arg Glu Tyr Ser
130 135 140
Pro Ser Ala Ala Arg Gly Leu Arg Leu Met Phe Thr Asp Phe Trp Arg
145 150 155 160
Thr Arg Val Leu Arg Gln Thr Ser Pro Met Ala Gly Val Phe Gly Asn
165 170 175
Leu Asp Val Asn Glu Ala Arg Leu Met Ala Ala Tyr Thr Ser Glu Cys
180 185 190
Ala Asp His Leu Glu Ala Lys Glu Leu Ala Gly Pro Asp Gly Val Ala
195 200 205
Ala Ala Arg Glu Ile Ala Lys Arg Trp Glu Lys Arg Val Arg Asp Leu
210 215 220
Gln Asp Lys Gly Ala Ala Arg Lys Leu Leu Asn Asp Pro Leu Gly Arg
225 230 235 240
Arg Thr Pro Asn Tyr Gln Ser Lys Asn Pro Gly Glu Tyr Thr Val Gly
245 250 255
Asn Ser Met Phe Tyr Asp Gly Pro Gln Val Ala Asn Leu Gln Asn Val
260 265 270
Asp Thr Gly Phe Trp Leu Asp Met Ser Asn Leu Ser Asp Val Val Leu
275 280 285
Ser Arg Glu Ile Gln Thr Gly Leu Arg Ala Arg Ala Thr Leu Glu Glu
290 295 300
Ser Met Pro Met Leu Glu Asn Leu Glu Glu Arg Phe Arg Arg Leu Gln
305 310 315 320
Glu Thr Cys Asp Ala Ala Arg Thr Glu Ile Glu Glu Ser Gly Trp Thr
325 330 335
Arg Glu Ser Ala Ser Arg Met Glu Gly Asp Glu Ala Gln Gly Pro Ser
340 345 350
Arg Val Gln Gln Ala Phe Gln Ser Phe Val Asn Glu Cys Asn Ser Ile
355 360 365
Glu Phe Ser Phe Gly Ser Phe Gly Glu His Val Arg Val Leu Cys Ala
370 375 380
Arg Val Ser Arg Gly Leu Ala Ala Ala Gly Glu Ala Ile Arg Arg Cys
385 390 395 400
Phe Ser Cys Cys Lys Gly Ser Thr His Arg Tyr Ala Pro Arg Asp Asp

```

                                405                410                415
Leu Ser Pro Glu Gly Ala Ser Leu Ala Glu Thr Leu Ala Arg Phe Ala
                                420                425                430
Asp Asp Met Gly Ile Glu Arg Gly Ala Asp Gly Thr Tyr Asp Ile Pro
                                435                440                445
Leu Val Asp Asp Trp Arg Arg Gly Val Pro Ser Ile Glu Gly Glu Gly
                                450                455                460
Ser Asp Ser Ile Tyr Glu Ile Met Met Pro Ile Tyr Glu Val Met Asn
                                465                470                475                480
Met Asp Leu Glu Thr Arg Arg Ser Phe Ala Val Gln Gln Gly His Tyr
                                485                490                495
Gln Asp Pro Arg Ala Ser Asp Tyr Asp Leu Pro Arg Ala Ser Asp Tyr
                                500                505                510
Asp Leu Pro Arg Ser Pro Tyr Pro Thr Pro Pro Leu Pro Pro Arg Tyr
                                515                520                525
Gln Leu Gln Asn Met Asp Val Glu Ala Gly Phe Arg Glu Ala Val Tyr
                                530                535                540
Ala Ser Phe Val Ala Gly Met Tyr Asn Tyr Val Val Thr Gln Pro Gln
                                545                550                555
Glu Arg Ile Pro Asn Ser Gln Gln Val Glu Gly Ile Leu Arg Asp Met
                                565                570                575
Leu Thr Asn Gly Ser Gln Thr Phe Arg Asp Leu Met Lys Arg Trp Asn
                                580                585                590
Arg Glu Val Asp Arg Glu
                                595

```